

Active Bacterial Core Surveillance (ABCs) Report Emerging Infections Program Network Haemophilus influenzae, 2017



Haemophiius injiuenzae, 201

ABCs Areas
California (3 county San Francisco Bay area); Colorado (5 county Denver area); Connecticut; Georgia; Maryland; Minnesota; New Mexico; New York (15 county Rochester and Albany areas); Oregon; Tennessee (20 urban counties)

ABCs Population

The surveillance areas represent 44,593,843 persons. Source: National Center for Health Statistics bridged-race vintage 2017 postcensal file

ABCs Case Definition

Invasive *Haemophilus influenzae* (Hi) disease: isolation of Hi from normally sterile site in a resident of a surveillance area in 2017.

ABCs Methodology

ABCs personnel routinely contacted all microbiology laboratories serving acute care hospitals in their area to identify cases. Standardized case report forms that include information on demographic characteristics, clinical syndrome, and outcome of illness were completed for each identified case. Serotyping was done on Hi isolates at CDC and state laboratories. Regular laboratory audits assessed completeness of active surveillance and detected additional cases.

All rates of invasive Hi disease were calculated using population estimates for 2017 from the bridged-race vintage 2017 postcensal file. For national estimates, race- and age-specific rates of disease were applied from the aggregate surveillance areas to the race- and age-specific distribution of the 2017 U.S. population. Cases with missing data, excluding ethnicity, were multiply imputed using sequential regression imputation methods. ¶

Reported ABCs Profiles

Race	No.	(Rate*)
White	766	(2.4)
Black	215	(2.6)
Other	42	(1.1)
Total	1023	(2.3)

^{*} Per 100,000 population for ABCs areas

¶ Surveillance Note

Missing race (n=75) data were multiply imputed using sequential regression imputation methods.

	Ca	ses	De	aths
Syndrome	No.	(% [*])	No.	(Rate [†])
Meningitis	70	(6.8)	1	(1.5)
Bacteremia without focus	272	(26.6)	31	(12.4)
Pneumonia with bacteremia	574	(56.1)	69	(12.1)

^{*} Percent of cases

Serotype

]	В	Non-B		Non-Type [†]		Unknown	
Age (years)	No.	(Rate*)	No.	(Rate*)	No. (Rate*)	No.	(Rate*)
< 1	2	(0.38)	19	(3.60)	26	(4.92)	3	(0.57)
1	2	(0.38)	15	(2.81)	8	(1.50)	0	(0.00)
2-4	1	(0.06)	12	(0.74)	10	(0.62)	1	(0.06)
5-17	1	(0.01)	6	(0.08)	18	(0.25)	1	(0.01)
18-34	0	(0.00)	7	(0.07)	52	(0.50)	7	(0.07)
35-49	1	(0.01)	25	(0.29)	62	(0.71)	14	(0.16)
50-64	1	(0.01)	72	(0.82)	131	(1.49)	13	(0.15)
65-74	1	(0.03)	35	(0.88)	112	(2.81)	21	(0.53)
75-84	1	(0.05)	33	(1.76)	132	(7.04)	18	(0.96)
<u>≥</u> 85	0	(0.00)	21	(2.53)	123	(14.81)	15	(1.81)
Total	10	(0.02)	245	(0.55)	674	(1.51)	93	(0.21)

Per 100,000 population for ABCs areas

National Estimates of Invasive Disease

Cases: 7,550 (2.32/100,000) Deaths: 910 (0.28/100,000)

Healthy People 2020 Update

Invasive Haemophilus influenzae type B disease

Objective: Decrease the incidence of invasive *Haemophilus influenzae* type B disease to 0.27 cases per 100,000 persons less than 5 years of age

Age (year)	2020 Objective	2017 Rate*
< 5	0.27/100,000	0.19/100,000

^{*} Per 100,000 U.S. population < 5 years

For more information, visit our web site:

http://www.cdc.gov/abcs

Citation

Centers for Disease Control and Prevention. 2017. Active Bacterial Core Surveillance Report, Emerging Infections Program Network, *Haemophilus influenza* 2017. Available via the internet:

http://www.cdc.gov/abcs/reports-findings/survreports/hi17.pdf

Jan 2019 File - Final March 07, 2019

[†] Deaths per 100 cases with known outcome

[†] Non-typeable isolates