

# Ecology, Epidemiology, and Prevention of Lyme Disease in the United States



**Paul Mead, MD, MPH**

*Chief, Epidemiology and Surveillance Activity*

Division of Vector-Borne Diseases

National Center for Emerging and Zoonotic Infectious Diseases

Centers for Disease Control and Prevention




U.S. Department of  
Health and Human Services  
Centers for Disease  
Control and Prevention

# The Essentials

- ❑ **Lyme disease is a multisystem vector-borne zoonosis caused by the spirochete *Borrelia burgdorferi***
- ❑ **Small mammals and birds are reservoirs**
- ❑ **Lyme disease is transmitted in North America by 2 species of black-legged ticks**
  - *Ixodes scapularis*
  - *Ixodes pacificus*



# From Ticks to Humans: Transmission of *B. burgdorferi*



- ❑ Nymphs are most active in late spring and early summer
- ❑ Nymphs play a major role in transmission to humans
- ❑ Deer are immune to infection by *B. burgdorferi*, but support tick populations

# National Surveillance for Lyme Disease


- ❑ **Lyme disease became nationally notifiable in 1991**
- ❑ **Confirmed case definition for surveillance purposes**
  - Erythema migrans with exposure in an endemic area, OR
  - Erythema migrans with laboratory evidence but no exposure, OR
  - Noncutaneous manifestation (e.g., arthritis, carditis, neuritis) with laboratory evidence of infection
- ❑ **Probable case definition added in 2008 to capture patients with a broader array of clinical features**

# Surveillance Challenges and Caveats

- ❑ **Verifying cases can be time-consuming**
- ❑ **Current magnitude of underreporting is unknown**
  - Estimates of “10 fold” underreporting are obsolete
- ❑ **Cases are reported according to county of residence, not county of exposure**



# In the United States Lyme Disease is Regional, but Spreading




1998

1 dot per case placed randomly in county of patient residence; may not reflect county of exposure



U.S. Department of  
Health and Human Services  
Centers for Disease  
Control and Prevention

# In the United States Lyme Disease is Regional, but Spreading




2008

1 dot per case placed randomly in county of patient residence; may not reflect county of exposure



U.S. Department of  
Health and Human Services  
Centers for Disease  
Control and Prevention

# Reported Lyme Disease Cases United States, 1991–2009



U.S. Department of  
Health and Human Services  
Centers for Disease  
Control and Prevention



# Top 7 Notifiable Diseases United States, 2009

Rank	Disease	U.S.
1	Chlamydia	1,244,180
2	Gonorrhea	301,174
3	Salmonellosis	49,192
4	Syphilis	44,828
5	Novel influenza A	43,696
6	Lyme disease	38,468
7	AIDS	36,870

Rank	Disease	New England
1	Chlamydia	39,246
2	Lyme disease	9,205
3	Gonorrhea	5,470
4	Salmonellosis	2,244
5	Varicella	1,729
6	Giardiasis	1,660

New England = CT, ME, MA, NH, RI, VT



U.S. Department of  
Health and Human Services  
Centers for Disease  
Control and Prevention

# Lyme Disease: Current Challenges

- ❑ **Clinical diagnosis and treatment**
- ❑ **Laboratory diagnostics**
- ❑ **Public health practice**
- ❑ **Prevention**
  - Personal protection in the absence of vaccine
  - Environmental management for tick control
  - Community-based interventions



# Personal Protection in the Absence of Vaccine



- Avoid tick habitat**
- Wear protective clothing**
- Use insect repellents**
- Check for ticks daily**
- Bathe promptly after exposure**



# Studies of Selected Personal Protective Measures

Use of insect repellents		Check for ticks		Reference
Effect	P value	Effect	P value	
<b>OR 0.6</b>	<b>NS</b>	<b>OR 0.5</b>	<b>0.02</b>	2009 Connally
<b>OR 0.8</b>	<b>0.05</b>	<b>OR 1.0</b>	<b>NS</b>	2008 Vázquez
<b>OR 0.7</b>	<b>0.02</b>	<b>OR 0.6</b>	<b>0.001</b>	2001 Smith G
<b>OR 1.2</b>	<b>NS</b>	<b>OR 1.2</b>	<b>NS</b>	
<b>OR 1.0</b>	<b>NS</b>	<b>OR 0.5</b>	<b>NS</b>	1998 Orloski
<b>–</b>	<b>NS</b>	<b>–</b>	<b>NS</b>	1996 Klein
<b>OR 1.5</b>	<b>NS</b>	<b>OR 0.8</b>	<b>NS</b>	1995 Ley
<b>RR 0.5</b>	<b>NS</b>	<b>RR 1.1</b>	<b>NS</b>	1988 Smith P <sup>1</sup>
<b>RR 0.7</b>	<b>NS</b>	<b>RR 0.8</b>	<b>NS</b>	

<sup>1</sup> Risk presented as inverse

OR, Odds ratio  
RR, Relative risk  
NS, Not significant




U.S. Department of Health and Human Services  
Centers for Disease Control and Prevention

# Bathing as Primary Prevention

## □ Prospective case control study of 364 Connecticut patients with Lyme disease diagnosed 2005–2007


Behavior	Adjusted OR (95% CI)
Wearing repellent while in yard	0.59 (0.35–1.03)
Checking for ticks within 36 hrs	0.55 (0.32–0.94)
Bathing within 2 hrs	0.42 (0.23–0.78)



# Environmental Management for Tick Control

## □ Landscaping to create “tick-safe zones”

- Clear brush and leaf litter
- 3-foot barrier of wood chips can reduce questing ticks in lawn by 50%
- Use deer-resistant plantings
- Install deer fencing



# Chemical Tick Control

- ❑ **A single, springtime application of pesticide can reduce questing tick populations by 68–100%**



Stafford III, KC. Tick Management Handbook (Bulletin 1010) 2007.  
Connecticut Agricultural Experiment Station, New Haven, CT  
<http://www.ct.gov/caes/lib/caes/documents/publications/bulletins/b1010.pdf>



**U.S. Department of  
Health and Human Services**  
Centers for Disease  
Control and Prevention

# Community-based Interventions


- ❑ **USDA “4-poster” stations treat deer with topical pesticide and reduce tick carriage**
- ❑ **Obstacles include concerns about pesticides and the spread of chronic wasting disease**



Photo credit by Scott Bauer, ARS



# Sharp, Community-wide Reductions in Deer Populations May Decrease Lyme Disease Cases Bridgeport, Connecticut



# Summary

- ❑ **Lyme disease is an important public health problem**
- ❑ **The number of cases continues to grow**
- ❑ **An array of prevention interventions are available**
- ❑ **Currently, there is no single, widely-accepted prevention method**



# CDC Lyme Disease Prevention Strategies

## Education, education, education

- Assure that current prevention options are widely known and adopted
- Use fewer but better targeted messages



U.S. Department of Health and Human Services  
Centers for Disease Control and Prevention

# CDC Lyme Disease Prevention Strategies

- ❑ **Improve current, and develop and validate new prevention methods**
  - Placebo-controlled trial of 1,600 households is under way to validate benefits of pesticide applications
  - Natural products from plant extracts
  - Rodent-targeted vaccines
  - Deer-based interventions



# Clinical Manifestations and Treatment of Lyme Disease



**Allen C. Steere, MD**

*Division of Rheumatology, Allergy and Immunology*

Massachusetts General Hospital

Harvard Medical School



U.S. Department of  
Health and Human Services  
Centers for Disease  
Control and Prevention

# Overview

- ❑ **How it all began**
- ❑ **Clinical manifestations**
  - Active infection
  - Postinfectious syndromes
- ❑ **Treatment**
  - What, when, and how long?
- ❑ **What's ahead**



Lyme, Connecticut



U.S. Department of  
Health and Human Services  
Centers for Disease  
Control and Prevention


# How it All Began

- ❑ **October 1975: Two mothers contacted health officials about arthritis cases in their communities (Lyme and Old Lyme, CT )**
- ❑ **January 1977: First description of “Lyme arthritis”**
  - Patients had an arthropod-transmitted illness
  - 1/4 of the children or their parents recalled an expanding skin lesion before the onset of arthritis

## LYME ARTHRITIS

AN EPIDEMIC OF OLIGOARTICULAR ARTHRITIS IN CHILDREN AND ADULTS IN THREE CONNECTICUT COMMUNITIES

ALLEN C. STEERE, STEPHEN E. MALAWISTA, DAVID R. SNYDMAN, ROBERT E. SHOPE, WARREN A. ANDIMAN, MARTIN R. ROSS, and FRANCIS M. STEELE



# Clinical Manifestations of Lyme Disease

## □ Stage 1: Localized infection

- Erythema migrans – a slowly expanding skin lesion, sometimes with partial central clearing
- Often with flu-like symptoms: Headache, stiff neck, myalgias, arthralgias, or fever, but no gastro-intestinal or respiratory symptoms
- About 1 in 5 patients lack this initial skin lesion, and the illness begins with flu-like symptoms or a later disease manifestation





# Clinical Manifestations of Lyme Disease

## □ Stage 2: Early disseminated infection

- Neuroborreliosis: About 15% of untreated patients
- Most commonly
  - Meningitis
  - Cranial neuropathy
  - Motor or sensory radiculoneuropathy
- Cardiac involvement: About 5% of untreated patients
  - Atrio-ventricular (AV) nodal block
  - Myopericarditis

# Clinical Manifestations of Lyme Disease

## □ Stage 3: Late persistent infection

- Arthritis – 60% of untreated patients
- Intermittent attacks in one or a few joints, especially the knee, sometimes becoming chronic
- Late subtle encephalopathy or polyneuropathy, accompanied by abnormal cerebrospinal fluid (CSF) or electromyogram (EMG)
- Late in the illness, the infection is usually quite localized, and systemic symptoms are minimal, if present at all
- Even without antibiotics, the immune system seems to win out eventually, and symptoms resolve



# Treatment of Early Lyme Disease

## Guidelines of the Infectious Diseases Society of America

### □ What

- Doxycycline or amoxicillin
- Cefuroxime or erythromycin  
(in case of allergy to doxycycline or amoxicillin)
- All taken by mouth

### □ How long

- 14–21 days

# Treatment of Later Manifestations of Lyme disease

## Guidelines of the Infectious Diseases Society of America

### □ **Early or late neuroborreliosis: 2–4 weeks**

- Ceftriaxone or cefotaxime, intravenously (IV)
- Na-penicillin G, IV

### □ **Heart involvement: 4 weeks**

- Generally, start with IV therapy
- When clinical picture improves, complete course with oral therapy

### □ **Joint involvement: 4–8 weeks**

- Oral regimens 4–8 weeks
- Some patients require IV antibiotics for 4 weeks for successful treatment of the infection

# Then and Now

- ❑ **Key clinical challenge today: How to diagnose and treat syndromes that may follow standard courses of antibiotic therapy for Lyme disease**
  - Distinguishing these symptoms from other illnesses
  - Most researchers think that these syndromes result from other factors than active infection
  - Strong feeling on the part of advocacy groups that these persistent symptoms result from persistent infection and require months or years of antibiotics



# Reasons for Persistent Signs or Symptoms after Antibiotic Treatment

## □ Neuroborreliosis

- Neurologic recovery (e.g., facial palsy) may be incomplete

## □ Antibiotic-refractory Lyme arthritis

- Proliferative synovitis may persist for months or several years after 1–2 months of oral antibiotics and 1 month of IV antibiotics
- Autoimmunity may play a role in the course of Lyme disease

# Reasons for Persistent Signs or Symptoms after Antibiotic Treatment

## □ Pain, neurocognitive, and/or fatigue symptoms

- In a small percentage of cases, these symptoms may begin after recommended courses of antibiotics for Lyme disease.
- CSF and EMG testing shows normal results
- The majority of patients now diagnosed with “chronic Lyme disease” have pain and fatigue symptoms, but lack evidence of past or present *B. burgdorferi* infection
  - Sigal LH, et al. Am J Med 1990;88:577-81
  - Steere, A et al. JAMA 1993;269:1812-16
  - Carrington, RM et al. Ann Intern Med 1998;128:354-62
- Amplification of sensory signals in the brain may be an important mechanism

# Antibiotic Therapy for Persistent Symptoms after Standard Antibiotic Treatment for Lyme Disease

## □ Pain, neurocognitive, and/or fatigue symptoms after Lyme disease

- Four double-blind, placebo-controlled trials have been conducted
- No sustained benefit from additional oral or IV antibiotic therapy has been shown
- Severe adverse reactions have been reported
  - Klempner, MS et al. N Engl J Med 2001;345:85-92
  - Krupp, LB et al. Neurology 2003;60:1923-30
  - Fallon, BA et al. Neurology 2008:992-1003



# Summary

## □ Lyme disease

- Multisystem infection
- Typically occurs in stages with different clinical manifestations at each stage

## □ Infection can be treated effectively with antibiotics

- Effective treatment is tailored to the disease manifestation
- Early disease can usually be treated effectively with oral antibiotics, but organ system involvement may require intravenous therapy



# Summary

## ❑ Post-infectious syndromes

- Incomplete recovery of nerve function
- Persistent synovitis after apparent killing of spirochete with antibiotics
- Pain, neurocognitive, and fatigue symptoms

❑ **Currently, there is no evidence for sustained benefit from further courses of antibiotic therapy, but there is potential for substantial harm because of adverse effects, particularly from IV antibiotics**

# What's Ahead?

- ❑ **Search for evidence of active *B. burgdorferi* after IDSA-recommended courses of antibiotic therapy**
- ❑ **Understand the role of autoimmunity in Lyme disease**
- ❑ **Understand and treat effectively centralized pain syndromes, not just in Lyme disease, but in the many conditions in which this may occur**

# Laboratory Testing for Lyme Disease



**Adriana Marques, MD**

*Laboratory of Clinical Infectious Diseases*

National Institute of Allergy and Infectious Diseases

National Institutes of Health



U.S. Department of  
Health and Human Services  
Centers for Disease  
Control and Prevention

# Disclosure Statement

I will not discuss off-label use and/or investigational use of drugs/devices.

I am a co-inventor on a patent application for the VOVO LIPS test for Lyme disease, in which one of the antigens is based on the IR6 peptide.



**U.S. Department of  
Health and Human Services**  
Centers for Disease  
Control and Prevention

# Overview

- ❑ **Current recommendations for laboratory tests for Lyme disease in the United States**
- ❑ **Facts and challenges**
- ❑ **Progress to improve laboratory testing**
- ❑ **Research needs and what's ahead**



# Methods for Laboratory Diagnosis of Lyme Disease

## ❑ Direct: Detection of *causative organism*

- Culturing *B. burgdorferi* from clinical specimens
- PCR detection of *B. burgdorferi* DNA from clinical specimens

## ❑ Indirect: Detection of *immune response to the causative organism*

- Detection of antibodies against *B. burgdorferi*

# Direct Methods: Detection of Causative Organism

## ❑ *B. burgdorferi* is more easily detected

- By culture and/or PCR: Skin and blood samples during the early stages of the disease (erythema migrans, when the diagnosis is mostly clinical)
- In the synovial fluid of patients with Lyme arthritis


## ❑ For other presentations, it is very difficult to confirm the presence of the bacteria

**No direct detection methods  
have been reviewed and approved by the FDA**



# Indirect Methods: Detection of Immune Response to the Causative Organism

- ❑ Serologic assays: Detecting antibodies to *B. burgdorferi*
- ❑ Current CDC recommendations: 2-tier algorithm




ELISA, Enzyme-linked immunosorbent assay  
IFA, Indirect immunofluorescence assay



U.S. Department of  
Health and Human Services  
Centers for Disease  
Control and Prevention

# Indirect Methods: Detection of Immune Response to the Causative Organism



# Facts and Challenges

## □ Facts

- The current algorithm works well when used as recommended
- Serological testing is not required for patients with erythema migrans
  - Patients who present very early in their illness are more likely to have a negative result
  - Less than 50% of the patients with erythema migrans lesions (stage 1) are positive at presentation
- Laboratory tests are most helpful in patients with stage 2 and stage 3 of Lyme disease



# Facts and Challenges

## ❑ Challenges: Appropriate use of tests

- About 3.4 million Lyme serology tests are performed annually in the United States (compared to 38,000 reported cases in 2009)
- Tests are being used in situations where they are not recommended
  - To rule out Lyme disease in populations with a low probability of having the disease
  - To test patients with suspected erythema migrans
  - To test people bitten by ticks
- Insufficiently validated tests and interpretation criteria are being used




# VlsE: A New Diagnostic Marker

- ❑ **VlsE (variable major protein-like sequence, expressed)**
  - An outer surface lipoprotein of *B. burgdorferi*
  - C6 peptide: Derived from its invariable region 6
- ❑ **Addition of VlsE to both 1<sup>st</sup> and 2<sup>nd</sup> tier tests has improved their performance**
- ❑ **C6 ELISA**
  - Shown to be more sensitive for patients with erythema migrans than standard 2-tiered testing, and is more specific than whole cell sonicate ELISA
  - FDA-approved as a 1<sup>st</sup> tier test; under study as a “stand-alone test”



**U.S. Department of  
Health and Human Services**  
Centers for Disease  
Control and Prevention

# Serological Testing and Duration of Illness Patients with a Single Erythema Migrans Lesion



Adapted from Wormser, GP et al. Clin Vaccine Immunol 2008;15:1519-22  
ELISA, Enzyme-linked immunosorbent assay  
WCS, Whole cell sonicate



U.S. Department of  
Health and Human Services  
Centers for Disease  
Control and Prevention

# Use of Laboratory Tests

## ❑ Current algorithm

- Works well when used as recommended
- Can be improved for patients with early stages of the disease, especially early neurological disease

## ❑ Sensitivity of the test increases with the duration of the infection

- Erythema migrans (stage 1): Treatment is indicated, no tests are necessary
- Stage 2 and 3: Tests are helpful

## ❑ In a patient with low probability of Lyme disease

- Negative ELISA test rules out the disease
- Positive ELISA test is more likely to be a false positive



# Use of Laboratory Tests

- ❑ **Current serologic assays do not distinguish between active and inactive infection**
  - Antibodies can persist after successful antibiotic therapy, including IgM antibodies
- ❑ **Positive IgM response alone does not distinguish clearly between Lyme disease and other conditions**
  - Positive IgM results for *B. burgdorferi* occur in
    - >50% of parvovirus B19 infections
    - Human granulocytic anaplasmosis, Epstein-Barr virus, and other infections
    - Autoimmune diseases



# What's Ahead

- ❑ **Improve direct methods for detecting *B. burgdorferi***
- ❑ **Improve current serology diagnostic testing algorithm**
  - Simplicity: A single test or test procedure
  - Objectivity: Quantitative data, independent of who reads the results
  - Greater sensitivity in early disease
  - Independence from disease duration
  - Avoiding using IgM Western blot
  - Decreased cost
- ❑ **Develop tests that can help follow response to therapy:  
Biomarkers for active infection**



# Lyme Disease in Minnesota: Trends and Challenges



**Ruth Lynfield, MD**

*State Epidemiologist and Medical Director*

Minnesota Department of Health



U.S. Department of  
Health and Human Services  
Centers for Disease  
Control and Prevention

# Overview

## ❑ Epidemiology of Lyme disease in Minnesota

## ❑ Challenges


- Prevention
- Laboratory diagnostics
- Adverse consequences of prolonged courses of antibiotics
- Legislation

## ❑ Way forward



# Lyme Disease in Minnesota, 2009


- ❑ **Confirmed cases: 1,065, 8<sup>th</sup> in the US**
- ❑ **Incidence: 20.2/100,000 population, 12<sup>th</sup> in the US**
  - Incidence varies throughout the state
  - Cass county: >100/100,000 population
    - Higher than overall incidence in CT  
78/100,000 in 2009



# Lyme Disease Cases United States, 2005–2009

	Minnesota	New England/ Mid-Atlantic	United States
<b>Median age</b>	<b>39 years</b>	<b>43 years</b>	<b>43 years</b>
<b>Range</b>	<b>Infant–98 years</b>	<b>Infant–109 years</b>	<b>Infant–109 years</b>
<b>Age distribution</b>	<b>33% &lt;18 years</b>	<b>25% &lt;18 years</b>	<b>25% &lt;18 years</b>
<b>Sex</b>	<b>62% male</b>	<b>56% male</b>	<b>54% male</b>

# Reported versus Confirmed Cases of Lyme Disease Minnesota, 1996–2010



# Reported versus Confirmed Cases of Lyme Disease Minnesota, 1996–2010

## ❑ Increase in reported cases: Perception

- Some may be due to increased awareness among the public and health care providers, increased compliance with reporting requirements, or improved surveillance


## ❑ Increase in reported cases: True increase in Lyme disease

- Lyme disease had been endemic and well-known in Minnesota for 15 years prior to this increase
- No new approaches to testing or reporting occurred during this period
- Data indicate ticks have spread into areas that border Minnesota's endemic areas






# Confirmed Lyme Disease Cases Minnesota, 1986–2010 (N =12,085 )



# Minnesota Biomes



<http://www.dnr.state.mn.us/biomes/index.html>






U.S. Department of  
Health and Human Services  
Centers for Disease  
Control and Prevention

# Lyme Disease Cases by County of Residence Minnesota, 1996–2010

1996–2000

2001–2005

2006–2010



**Incidence rate (cases/100,000 person-years)**



No cases



>10



10–100



100–160

# Lyme Disease: Challenges at the State Level

- Prevention**
- Laboratory diagnostics**
- Adverse consequences of prolonged courses of antibiotics**
- Legislation**

**CHALLENGES**  
CHALLENGES?



# Prevention Challenges

## Use of Personal Protection Measures in Reported Tick-Borne Disease Cases, Minnesota, 2008

Persons with Lyme disease, human anaplasmosis, and babesiosis who self-reported in the month prior to onset (No = 980)	%
Checked for ticks	73
Wore long pants	72
Used repellent	42
Wore light-colored clothing	39
Checked for ticks and used repellent	37
Avoided the woods	13



# Prevention Challenges

## Minnesota Department of Health Strategies

### □ Personal protection

- Provide information on the MDH website
- Provide phone consultations
- Reach out to the community
  - Give talks, especially to high-risk groups (e.g., loggers, foresters)
  - Give lectures to health care providers and others
  - Conduct interviews with the media

### □ Environmental tick control

- Provide information on the MDH website
  - In May 2009, the tick-borne disease web page had 40,000 hits; 3<sup>rd</sup> most frequently read MDH site
- Offer Metropolitan Mosquito Control District consultations to Minneapolis-St. Paul metropolitan area landowners

# Laboratory Diagnostic Challenges

## ❑ Lyme disease testing for clinical diagnosis

- Overuse of Lyme disease tests
  - Testing patients with EM with illness duration of <2–3 weeks (unnecessary and lower sensitivity of antibody test)
- Lyme disease testing: Misinterpretation
  - A positive IgM and a negative IgG >30 days into an illness is not indicative of Lyme disease

# Laboratory Diagnostic Challenges

## Minnesota Department of Health Strategies

### ❑ **Lyme disease testing**

- Send State Health Advisories electronically through the MDH Health Alert Network to local public health agencies and clinics
- Provide information on the MDH website
  - When to test patients for Lyme disease
  - How to interpret test results
  - Links to CDC and Infectious Disease Society of America diagnosis/treatment information

### ❑ **Give lectures to healthcare providers**

### ❑ **Publish an article on Lyme disease in MN Medicine**

- Kemperman, M et al. Minnesota Medicine. 2008; 91:37-41



# Adverse Consequences of Prolonged Courses of Antibiotics for Lyme Disease

- ❑ **Adverse effects range from mild to severe**
- ❑ **Severe adverse effects include**
  - Bloodstream infections in persons with central venous catheters receiving parenteral antibiotic therapy
    - Septic thrombosis and death due to *Candida*
  - Venous thrombosis
  - Severe allergic reactions
  - Cholecystitis
  - *Clostridium difficile* infection

# Adverse Consequences of Long-term Use of Antibiotics for Presumed Lyme Disease

## Minnesota Experience

### ❑ History and clinical presentation

- History of depression
- Fatigue, insomnia, achy joints, memory loss

### ❑ Laboratory testing for Lyme disease

- IFA: Indeterminate
- IgM Western blot: Positive
- IgG Western blot: Negative

### ❑ Treatment

- Doxycycline, 5 weeks; cefuroxime and telithromycin, 2–4 months
- Developed diarrhea 5 weeks into course; emergency colectomy

### ❑ Postmortem diagnosis: Fulminant *C. difficile*


#### Death Due to Community-Associated *Clostridium difficile* in a Woman Receiving Prolonged Antibiotic Therapy for Suspected Lyme Disease

TO THE EDITOR—*Clostridium difficile* infections can occur outside the hospital in association with antibiotic use and can result in fulminant colitis and death. In December 2009, the Minnesota Department of Health investigated a death due to *C. difficile* of a 52-year-old woman with no recent hospitalizations.

# Adverse Consequences of Antibiotics for Presumed Lyme Disease

## Minnesota Experience

- ❑ **2 nonfatal *C. difficile* cases reported to MDH with onsets in March 2007 and November 2010 in patients given prolonged courses of antibiotics for treatment of presumed Lyme disease**
  - Neither *C. difficile* case was reported to MDH as Lyme disease



# Lyme Disease Legislation at the State Level

- ❑ **Many states have passed physician protection and/or health insurance coverage bills for prolonged antibiotic treatment of patients with Lyme disease**



**U.S. Department of  
Health and Human Services**  
Centers for Disease  
Control and Prevention

# Lyme Disease Legislation in Minnesota

## ❑ **Minnesota: Physician protection bill brought before Health Committees in 2010 (HF2597; SF1631/2584)**

*“Board of Medical Practice limited from bringing a disciplinary action against a physician for prescribing, administering, or dispensing long-term antibiotic therapy for chronic Lyme disease.”*

## ❑ **Prior to bill becoming law, a compromise with the Minnesota Board of Medical Practice was reached**

<http://www.state.mn.us/portal/mn/jsp/home.do?agency=BMP>

*“MN Board of Medical Practice voluntarily will engage in a moratorium for a time period not to exceed 5 years, or the time at which double-blind, peer reviewed studies have resolved the issues, whichever is first, on the investigation, disciplining, or issuance of Corrective Action.”*



# Lyme Disease in Minnesota Summary

- ❑ **Incidence of Lyme disease is increasing in Minnesota**
  - Due to expansion of ticks into areas bordering endemic areas
- ❑ **Accurate surveillance is important, but is resource intensive**
- ❑ **Information about Lyme disease must be made available to the public and health care providers**
  - Prevention
  - Diagnosis
  - Adverse effects associated with prolonged courses of antibiotics
- ❑ **Concern about persistent non-specific symptoms that some individuals attribute to active Lyme disease is increasingly becoming a political issue**



# Lyme Disease in the United States

- ❑ **Improve understanding of reasons for increase in Lyme disease incidence**
- ❑ **Develop and effectively implement available preventative strategies**
- ❑ **Improve laboratory diagnostics**
  - Accurate and sensitive diagnostics for early illness
  - Improved laboratory tests for direct detection of the causative agent
  - Biomarkers indicative of active infection that can help follow response to therapy

**WAY FORWARD**



**U.S. Department of  
Health and Human Services**  
Centers for Disease  
Control and Prevention

# Lyme Disease in the United States

- ❑ **Improve understanding of prevalence and etiology of persistent symptoms**
  - In individuals following antibiotic treatment for Lyme disease
  - In individuals with no evidence of having had Lyme disease
- ❑ **Educate public, health care providers, and legislators**

**WAY FORWARD**



U.S. Department of  
Health and Human Services  
Centers for Disease  
Control and Prevention





Photo credit of Minnesota forest: M Kemperman, Minnesota Department of Health



**U.S. Department of Health and Human Services**  
Centers for Disease Control and Prevention