

HIV Surveillance **Report**

Volume 25

Diagnoses of HIV Infection in the United States and Dependent Areas, 2013

National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention
Division of HIV/AIDS Prevention



The *HIV Surveillance Report* is published annually by the Division of HIV/AIDS Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, Georgia.

Data are presented for diagnoses of HIV infection reported to CDC through June 2014.

The *HIV Surveillance Report* is not copyrighted and may be used and reproduced without permission. Citation of the source is, however, appreciated.

Suggested citation

Centers for Disease Control and Prevention. <i>HIV Surveillance Report, 2013</i> ; vol. 25. http://www.cdc.gov/hiv/library/reports/surveillance/ . Published February 2015. Accessed [date].
--

On the Web: <http://www.cdc.gov/hiv/library/reports/surveillance/>

Confidential information, referrals, and educational material on HIV infection

CDC-INFO

1-800-232-4636 (in English, en Español)

1-888-232-6348 (TTY)

<http://www.cdc.gov/cdc-info/requestform.html>

Acknowledgments

Publication of this report would not have been possible without the contributions of the state and territorial health departments and the HIV surveillance programs that provided surveillance data to CDC.

This report was prepared by the following CDC staff and contractors: Anna Satcher Johnson, Cheryl Williams, Xiaohong Hu, Jianmin Li, Baohua Wu, Xiuchan Guo, Hong Zhu, Marie Morgan, Michael Friend, Mi Chen, Sridevi Wilmore, John Gerstle, Shelley Ren, Jason Price, Irene Hall, Timothy Green, Kimberly Crenshaw, the HIV Incidence and Case Surveillance Branch, the Quantitative Sciences and Data Management Branch, and the Prevention Communication Branch.

Contents

Commentary	5
Technical Notes	11
Tables	
Section 1 Diagnoses of HIV Infection and Diagnoses of Infection Classified as Stage 3 (AIDS)	
1a Diagnoses of HIV infection, by year of diagnosis and selected characteristics, 2009–2013—United States	17
1b Diagnoses of HIV infection, by year of diagnosis and selected characteristics, 2009–2013—United States and 6 dependent areas	19
2a Stage 3 (AIDS), by year of diagnosis and selected characteristics, 2009–2013 and cumulative—United States	21
2b Stage 3 (AIDS), by year of diagnosis and selected characteristics, 2009–2013 and cumulative—United States and 6 dependent areas	23
3a Diagnoses of HIV infection, by race/ethnicity and selected characteristics, 2013—United States	25
3b Diagnoses of HIV infection, by race/ethnicity and selected characteristics, 2013—United States and 6 dependent areas	27
4a Stage 3 (AIDS), by race/ethnicity and selected characteristics, 2013—United States	29
4b Stage 3 (AIDS), by race/ethnicity and selected characteristics, 2013—United States and 6 dependent areas	31
5a Diagnoses of HIV infection among children aged <13 years, by race/ethnicity, 2009–2013—United States	33
5b Diagnoses of HIV infection among children aged <13 years, by race/ethnicity, 2009–2013—United States and 6 dependent areas	33
6a Stage 3 (AIDS) among children aged <13 years, by race/ethnicity, 2009–2013 and cumulative—United States	34
6b Stage 3 (AIDS) among children aged <13 years, by race/ethnicity, 2009–2013 and cumulative—United States and 6 dependent areas	34
7 Stage 3 (AIDS) among children aged <13 years, by year of diagnosis, 1992–2013—United States and 6 dependent areas	35
8 Diagnoses of HIV infection among adult and adolescent Hispanics/Latinos, by transmission category and place of birth, 2013—United States and 6 dependent areas	36
9 Stage 3 (AIDS) among adult and adolescent Hispanics/Latinos, by transmission category and place of birth, 2013—United States and 6 dependent areas	37
Section 2 Deaths and Survival after a Diagnosis of HIV Infection or Stage 3 (AIDS) Classification	
10a Deaths of persons with diagnosed HIV infection, by year of death and selected characteristics, 2009–2012—United States	38
10b Deaths of persons with diagnosed HIV infection, by year of death and selected characteristics, 2009–2012—United States and 6 dependent areas	40
11a Deaths of persons with diagnosed HIV infection ever classified as stage 3 (AIDS), by year of death and selected characteristics, 2009–2012 and cumulative—United States	42
11b Deaths of persons with diagnosed HIV infection ever classified as stage 3 (AIDS), by year of death and selected characteristics, 2009–2012 and cumulative—United States and 6 dependent areas	44

12a	Survival for more than 12, 24, and 36 months after a diagnosis of HIV infection during 2004–2009, by selected characteristics—United States	46
12b	Survival for more than 12, 24, and 36 months after a diagnosis of HIV infection during 2004–2009, by selected characteristics—United States and 6 dependent areas	47
13a	Survival for more than 12, 24, and 36 months after a stage 3 (AIDS) classification during 2004–2009, by selected characteristics—United States	48
13b	Survival for more than 12, 24, and 36 months after a stage 3 (AIDS) classification during 2004–2009, by selected characteristics—United States and 6 dependent areas	49

Section 3 Persons Living with Diagnosed HIV Infection or with Infection Ever Classified as Stage 3 (AIDS) (Prevalence)

14a	Persons living with diagnosed HIV infection, by year and selected characteristics, 2009–2012—United States	50
14b	Persons living with diagnosed HIV infection, by year and selected characteristics, 2009–2012—United States and 6 dependent areas	52
15a	Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by year and selected characteristics, 2009–2012—United States	54
15b	Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by year and selected characteristics, 2009–2012—United States and 6 dependent areas	56
16a	Persons living with diagnosed HIV infection, by race/ethnicity and selected characteristics, year-end 2012—United States	58
16b	Persons living with diagnosed HIV infection, by race/ethnicity and selected characteristics, year-end 2012—United States and 6 dependent areas	60
17a	Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and selected characteristics, year-end 2012—United States	62
17b	Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and selected characteristics, year-end 2012—United States and 6 dependent areas	64

Section 4 Data for States and Metropolitan Statistical Areas

18	Diagnoses of HIV infection, by area of residence, 2013—United States and 6 dependent areas	66
19	Stage 3 (AIDS), by area of residence, 2013 and cumulative—United States and 6 dependent areas	67
20	Adults and adolescents living with diagnosed HIV infection, by race/ethnicity and area of residence, year-end 2012—United States and 6 dependent areas	69
21	Adults and adolescents living with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and area of residence, year-end 2012—United States and 6 dependent areas	71
22	Diagnoses of HIV infection, 2013, and persons living with diagnosed HIV infection (prevalence), year-end 2012, by metropolitan statistical area of residence—United States and Puerto Rico	73
23	Stage 3 (AIDS), 2013 and cumulative, and persons living with diagnosed HIV infection ever classified as stage 3 (AIDS) (prevalence), year-end 2012, by metropolitan statistical area of residence—United States and Puerto Rico	77

Web Addresses for Reports of State and Local HIV Surveillance	82
--	-----------

The Centers for Disease Control and Prevention (CDC) collects, analyzes, and disseminates surveillance data on HIV infection; these data are one of the nation's primary sources of information on HIV in the United States. The annual surveillance report, published by the Division of HIV/AIDS Prevention (DHAP), summarizes information about diagnosed HIV infection in the United States and dependent areas. HIV surveillance data are used by CDC's public health partners in other federal agencies, health departments, nonprofit organizations, academic institutions, and the general public to help focus prevention efforts, plan services, allocate resources, develop policy, and monitor trends in HIV infection.

Estimated numbers and rates of diagnoses of HIV infection that were made during 2009–2013 are included from all 50 states, the District of Columbia, and 6 U.S. dependent areas (American Samoa, Guam, the Northern Mariana Islands, Puerto Rico, the Republic of Palau, and the U.S. Virgin Islands).

During 2009–2013, the annual estimated number and rate of diagnoses of HIV infection in the United States remained stable. Estimated numbers and rates of diagnoses of HIV infection increased in some subgroups and decreased in others. Variations in trends among groups are expected and may be due to targeted HIV testing initiatives, differences in testing behaviors, or changes in the numbers of new HIV infections. HIV diagnosis data can be used to monitor primary prevention efforts, testing initiatives, and awareness of serostatus among persons living with HIV. HIV incidence surveillance data provide the best indication of changes in the numbers of new HIV infections.

DEFINITIONS AND DATA SPECIFICATIONS

In this report, the term *diagnosis of HIV infection* is defined as a diagnosis of HIV infection regardless of the stage of disease (stage 1, 2, 3 [AIDS], or unknown) and refers to all persons with a diagnosis of HIV infection. The term *HIV infection, stage 3 (AIDS)*, and its condensed version—*stage 3 (AIDS)*—refer specifically to persons with diagnosed HIV whose infection was classified as stage 3 (AIDS) during a given year (for diagnoses) or whose infection

has ever been classified as stage 3 (AIDS) (for prevalence and deaths). In this report, the 2008 HIV case definition was used to classify HIV infection among adults and adolescents and among children. For more information on the stages of HIV infection, please review the Technical Notes.

The data on diagnoses of HIV infection reflect the date of diagnosis (diagnosed by December 31, 2013; reported to CDC as of June 30, 2014), not the date of report to CDC. Similarly, the data specific to stage 3 (AIDS) reflect the first date the criteria for stage 3 (AIDS) were met. HIV diagnoses do not necessarily represent new infections (incidence): some persons with newly diagnosed infections were infected recently, and others were infected at some time in the past. Because of reporting delays, the actual number of cases diagnosed in a given year may be higher than the numbers of diagnoses of HIV infection (unadjusted) presented for recent years; however, reporting delays are typically resolved in 2 to 3 years. Only a few cases are reported more than 3 years after diagnosis. Where indicated, counts of diagnoses and deaths have been statistically adjusted (estimated) to correct for delays in reporting.

HIGHLIGHTS OF ANALYSES

The statements in this section, unless otherwise indicated, are based on estimated data. All rates are per 100,000 population. The standard used for reporting trends in numbers and rates is based on an increase or a decrease of 5% or more during the specified time-frame (e.g., when comparing 2009 and 2013). Exceptions to this standard were made where numbers were small or rates were based on small numbers.

Please read all table titles and footnotes carefully to ensure a complete understanding of the displayed data.

Diagnoses of HIV Infection and Diagnoses of Infection Classified as Stage 3 (AIDS)

Diagnoses of HIV infection

From 2009 through 2013, the annual estimated number of diagnoses of HIV infection in the United States and the estimated rate remained stable (Table 1a). In 2013, the estimated rate was 15.0.

- **Age group:** From 2009 through 2013, the rates for persons aged 13–14, 20–24, 25–29, 60–64, and 65 years and older increased. The rates for persons aged 15–19, 35–39, and 40–44 years decreased. The rates remained stable for children (aged less than 13 years) and persons aged 30–34, 45–49, 50–54, and 55–59 years. In 2013, the highest rate was for persons aged 25–29 years (36.3), followed by the rate for persons aged 20–24 years (35.3).
- **Race/ethnicity:** From 2009 through 2013, the rates for American Indians/Alaska Natives and Asians increased. The rates for Hispanics/Latinos, Native Hawaiians/other Pacific Islanders, and persons of multiple races decreased. The rates for blacks/African Americans and whites remained stable. In 2013, the rates were 55.9 for blacks/African Americans, 18.7 for Hispanics/Latinos, 16.8 for persons of multiple races, 12.7 for Native Hawaiians/other Pacific Islanders, 9.4 for American Indians/Alaska Natives, 6.6 for whites, and 6.0 for Asians. Please use caution when interpreting data for Native Hawaiians/other Pacific Islanders: the numbers are small.
- **Sex:** From 2009 through 2013, the rates for female adults and adolescents decreased; the rates for males remained stable. In 2013, males accounted for 80% of all diagnoses of HIV infection among adults and adolescents. The rate for adult and adolescent males was 29.4, and the rate for females was 6.9.
- **Transmission category:** From 2009 through 2013, among adult and adolescent males, the annual number of diagnosed HIV infections attributed to male-to-male sexual contact increased. The numbers of infections attributed to injection drug use, male-to-male sexual contact *and* injection drug use, or heterosexual contact decreased. Among adult and adolescent females, the numbers of infections attributed to injection drug use or heterosexual contact decreased. In 2013, among adult and adolescent males and females, the diagnosed infections attributed to male-to-male sexual contact (68%, including male-to-male sexual contact *and* injection drug use) and those attributed to heterosexual contact (25%) accounted for approximately 93% of diagnosed HIV infections in the United States.
- **Region:** From 2009 through 2013, the rates of diagnoses of HIV infection in the West decreased,

and the rates in the Northeast, Midwest, and South remained stable. In 2013, rates were 20.5 in the South, 15.9 in the Northeast, 10.8 in the West, and 9.0 in the Midwest.

Stage 3 (AIDS)

From 2009 through 2013, the annual estimated number and the estimated rate of infections classified as stage 3 (AIDS) in the United States decreased (Table 2a). In 2013, the estimated rate of infections classified as stage 3 (AIDS) was 8.4.

- **Age group:** From 2009 through 2013, the rates of stage 3 (AIDS) classifications for persons aged 20–24 and 60–64 years increased. The rates for persons aged 13–14, 30–34, 35–39, 40–44, 45–49, 50–54, 55–59, and 65 years and older decreased. The rates remained stable for children (aged less than 13 years) and persons aged 15–19 and 25–29 years. In 2013, the highest rate was for persons aged 45–49 years (17.8), followed by the rate for persons aged 40–44 years (16.8).
- **Race/ethnicity:** From 2009 through 2013, rates remained stable for American Indians/Alaska Natives and decreased for all other race/ethnicity groups: Asians, blacks/African Americans, Hispanics/Latinos, Native Hawaiians/other Pacific Islanders, whites, and persons of multiple races. In 2013, the rates were 33.7 for blacks/African Americans, 14.0 for persons of multiple races, 9.9 for Hispanics/Latinos, 6.9 for Native Hawaiians/other Pacific Islanders, 4.5 for American Indians/Alaska Natives, 3.4 for whites, and 2.6 for Asians. Please use caution when interpreting data for Native Hawaiians/other Pacific Islanders: the numbers are small.
- **Sex:** From 2009 through 2013, the rates for adult and adolescent males and females decreased. In 2013, adult and adolescent males accounted for 76% of all infections classified as stage 3 (AIDS) among adults and adolescents. The 2013 rate for males was 15.7; the 2013 rate for females was 4.8.
- **Transmission category:** From 2009 through 2013, the annual number of infections classified as stage 3 (AIDS) among adult and adolescent males with HIV infection attributed to male-to-male sexual contact, injection drug use, male-to-male sexual contact *and* injection drug use, or heterosexual contact decreased. The number of infections classified as stage 3 (AIDS) among adult and adolescent females with HIV infection

attributed to injection drug use or heterosexual contact decreased.

- **Region:** From 2009 through 2013, the rates of infections classified as stage 3 (AIDS) decreased in all 4 regions of the United States: Northeast, Midwest, South, and West. In 2013, rates were 12.1 in the South, 8.7 in the Northeast, 5.7 in the West, and 4.8 in the Midwest.

Deaths

Deaths of persons with diagnosed HIV infection

From 2009 through 2012 in the United States, the annual estimated number and rate of deaths of persons with diagnosed HIV infection decreased (Table 10a). In 2012, the estimated rate of deaths of persons with diagnosed HIV infection was 5.5. Deaths of persons with diagnosed HIV infection may be due to any cause (i.e., may or may not be related to HIV infection).

- **Age group:** From 2009 through 2012, the rates of deaths for persons aged 60–64 years increased. The rates of deaths for persons aged 15–19, 20–24, 25–29, 30–34, 35–39, 40–44, 45–49, and 50–54 years decreased. The rates remained stable for children (aged less than 13 years at death) and persons aged 13–14, 55–59, and 65 years and older.
- **Race/ethnicity:** From 2009 through 2012, the rates of deaths for whites remained stable. The rates of deaths for all other race/ethnicity groups—American Indians/Alaska Natives, Asians, blacks/African Americans, Hispanics/Latinos, Native Hawaiians/other Pacific Islanders, and persons of multiple races—decreased. In 2012, the highest rate of deaths was for blacks/African Americans: 21.1. Please use caution when interpreting data for Native Hawaiians/other Pacific Islanders: the numbers are small.
- **Sex and transmission category:** From 2009 through 2012, the rates of deaths for adult and adolescent males and females decreased. The numbers of deaths among males with diagnosed HIV infection attributed to perinatal transmission increased. The numbers of deaths among males with diagnosed HIV infection attributed to male-to-male sexual contact remained stable. The numbers of deaths among males with infection attributed to injection drug use, male-to-male sexual contact *and* injection drug use, or heterosexual contact decreased. The numbers of deaths of adult and adolescent females with infection

attributed to perinatal transmission remained stable. The numbers of deaths of females with infection attributed to injection drug use or heterosexual contact decreased. Please use caution when interpreting trend data for persons with infection attributed to perinatal transmission: the numbers are small.

- **Region:** From 2009 through 2012, the rates of deaths in the South, the Northeast, and the Midwest decreased. The rates in the West remained stable. In 2012, the rate was 7.8 in the Northeast, 7.0 in the South, 3.7 in the West, and 2.8 in the Midwest.

Deaths of persons with infection ever classified as stage 3 (AIDS)

From 2009 through 2012 in the United States, the annual estimated number and rate of deaths of persons with infection ever classified as stage 3 (AIDS) decreased (Table 11a). In 2012, the estimated rate of deaths of persons with stage 3 (AIDS) was 4.4. Deaths of persons with stage 3 (AIDS) may be due to any cause.

- **Age group:** From 2009 through 2012, the rates of deaths among persons aged 60–64 and 65 years and older increased. Rates of deaths among persons aged 20–24, 25–29, 30–34, 35–39, 40–44, 45–49, 50–54, and 55–59 years decreased. Rates remained stable among children (aged less than 13 years) and among persons aged 13–14 and 15–19 years.
- **Race/ethnicity:** From 2009 through 2012, the annual rate of deaths for whites remained stable. The rates of deaths for all other race/ethnicity groups—American Indians/Alaska Natives, Asians, blacks/African Americans, Hispanics/Latinos, Native Hawaiians/other Pacific Islanders, and persons of multiple races—decreased. The highest death rate was for blacks/African Americans: 16.9. Please use caution when interpreting data for Native Hawaiians/other Pacific Islanders: the numbers are small.
- **Sex and transmission category:** From 2009 through 2012, the overall rates for adult and adolescent males and females decreased. The numbers of deaths of males with stage 3 (AIDS) whose HIV infection was attributed to perinatal transmission increased. The numbers of deaths of males with infection attributed to male-to-male sexual contact, injection drug use, male-to-male sexual contact *and* injection drug use, or

heterosexual contact decreased. Among adult and adolescent females, the numbers of deaths of those with infection attributed to perinatal transmission increased. The numbers of deaths of females with infection attributed to injection drug use or to heterosexual contact decreased. Please use caution when interpreting data for persons with infection attributed to perinatal transmission: the numbers are small.

- **Region:** From 2009 through 2012, the rates of deaths decreased in all 4 regions of the United States: Northeast, Midwest, South, and West. In 2012, the rate was 6.2 in the Northeast, 5.6 in the South, 3.0 in the West, and 2.2 in the Midwest.

Survival

Survival after a diagnosis of HIV infection

Unadjusted data on diagnoses of HIV infection made during 2004–2009 in the United States were used to describe the proportions of persons surviving 12, 24, and 36 months after a diagnosis of HIV infection (Table 12a).

- **Year of diagnosis:** Survival increased with the year of diagnosis for diagnoses made during 2004–2009, although year-to-year differences were small.
- **Age group:** Survival decreased as age at diagnosis increased, particularly among persons aged 45 years and older at time of diagnosis.
- **Race/ethnicity:** Survival was greatest among Asians, followed by Hispanics/Latinos, Native Hawaiians/other Pacific Islanders, whites, and persons of multiple races. Survival was lowest among American Indians/Alaska Natives and blacks/African Americans.
- **Transmission category:** Survival was greatest among children regardless of transmission category; survival was slightly lower among children with diagnosed HIV infection attributed to perinatal exposure. Among adults and adolescents, survival was greatest among males with diagnosed HIV infection attributed to male-to-male sexual contact, females with infection attributed to heterosexual contact, and males with infection attributed to male-to-male sexual contact *and* injection drug use. Survival was intermediate among adult and adolescent males with infection attributed to heterosexual contact. Survival was lowest among

adult and adolescent males and females with infection attributed to injection drug use.

- **Region:** Survival was greatest among persons residing in the Midwest and the West, followed by those in the Northeast and the South.

Survival after stage 3 (AIDS) classification

Unadjusted data from diagnosed infections classified as stage 3 (AIDS) during 2004–2009 in the United States were used to describe persons surviving 12, 24, and 36 months after stage 3 (AIDS) classification (Table 13a).

- **Year of diagnosis:** In general, survival increased with the year of diagnosis for diagnoses made during 2004–2009, although year-to-year differences were small.
- **Age group:** In general, survival decreased as age at diagnosis increased, particularly among persons aged 30 years and older. Survival was greatest for persons aged 13–14 and 15–19 years at the time of stage 3 (AIDS) classification.
- **Race/ethnicity:** Survival was greatest among Asians, followed by Hispanics/Latinos, Native Hawaiians/other Pacific Islanders, persons of multiple races, whites, blacks/African Americans, and American Indians/Alaska Natives.
- **Transmission category:** Survival was greatest among children with infection attributed to perinatal transmission, followed by adult and adolescent males with infection attributed to male-to-male sexual contact. Survival was intermediate among adult and adolescent males with infection attributed to male-to-male sexual contact *and* injection drug use and among adult and adolescent males and females with infection attributed to heterosexual contact. Survival was lowest among males and females with infection attributed to injection drug use.
- **Region:** Survival was greatest among persons residing in the West, followed by those in the Northeast and the Midwest. Survival was lowest in the South.

Prevalence: Persons Living with Diagnosed HIV Infection and Persons Living with Infection Ever Classified as Stage 3 (AIDS)

Prevalence of diagnosed HIV infection

From 2009 through 2012, the estimated number of persons in the United States living with diagnosed HIV infection increased (Table 14a). At the end of 2012, an estimated 914,826 persons in the United States were living with diagnosed HIV infection. The estimated prevalence rate of diagnosed HIV infection was 291.5.

- **Age group:** At the end of 2012, the highest prevalence rate (769.4) and the largest percentage (18%) were those among persons aged 45–49 years, followed by those aged 50–54 (682.7 and 17%, respectively) and those aged 40–44 (596.6 and 14%, respectively). From 2009 through 2012, the largest percentage increase in rates (41%) was among persons aged 65 years and older (from 76.1 in 2009 to 107.4 in 2012).
- **Race/ethnicity:** At the end of 2012, the highest rate (1,011.0) and the largest percentage (43%) were those among blacks/African Americans. Among the remaining race/ethnicity groups, the rates were 488.1, persons of multiple races; 347.8, Hispanics/Latinos; 166.4, Native Hawaiians/other Pacific Islanders; 149.2, whites; 124.1, American Indians/Alaska Natives; and 70.5, Asians.
- **Sex and transmission category:** At the end of 2012, the estimated rates were 540.7 for adult and adolescent males, 167.0 for adult and adolescent females, and 4.8 for children (aged less than 13 years at year-end 2012). Among a total of 689,267 males (aged 13 years and older at year-end 2012), 69% of infections were attributed to male-to-male sexual contact, 12% to injection drug use, 11% to heterosexual contact, 7% to male-to-male sexual contact *and* injection drug use, and 1% to perinatal transmission. Among a total of 223,041 females (aged 13 years and older at year-end 2012), 74% of infections were attributed to heterosexual contact, 24% to injection drug use, and 2% to perinatal transmission. Among a total of 2,519 children aged less than 13 years at year-end 2012, 83% of infections were attributed to perinatal transmission.
- **Region:** Prevalence rates were highest in the Northeast (418.6), followed by the South (338.3), the West (239.0), and the Midwest (162.0).

Prevalence of stage 3 (AIDS)

From 2009 through 2012, the estimated number of persons living with infection ever classified as stage 3 (AIDS) increased (Table 15a). At the end of 2012, an estimated 508,845 persons in the United States were living with infection ever classified as stage 3 (AIDS). The estimated prevalence rate of stage 3 (AIDS) was 162.1.

- **Age group:** At the end of 2012, the highest rate (469.3) and the largest percentage (20%) were those among persons aged 45–49 years, followed by persons aged 50–54 (437.7 and 19%, respectively). From 2009 through 2012, the largest percentage increase in rates (41%) was among persons aged 65 years and older (from 51.3 in 2009 to 72.4 in 2012).
- **Race/ethnicity:** At the end of 2012, the highest rate (551.0) and the largest percentage (42%) were those among blacks/African Americans. Among the remaining race/ethnicity groups, the rates were 299.7, persons of multiple races; 206.7, Hispanics/Latinos; 89.1, Native Hawaiians/other Pacific Islanders; 81.0, whites; 66.8, American Indians/Alaska Natives; and 37.4, Asians.
- **Sex and transmission category:** At the end of 2012, the estimated rates, by age designation, were 304.8 for adult and adolescent males, 89.9 for adult and adolescent females, and 0.6 for children (aged less than 13 years at year-end 2012). Among a total of 388,527 males (aged 13 years and older at year-end 2012), 65% of infections were attributed to male-to-male sexual contact, 14% to injection drug use, 11% to heterosexual contact, 8% to male-to-male sexual contact *and* injection drug use, and 1% to perinatal transmission. Among a total of 120,012 females (aged 13 years and older at year-end 2012), 70% of infections were attributed to heterosexual contact, 27% to injection drug use, and 2% to perinatal transmission. Among a total of 305 children aged less than 13 years at year-end 2012, 94% of infections were attributed to perinatal transmission.
- **Region:** Prevalence rates were highest in the Northeast (248.7), followed by the South (179.6), the West (140.5), and the Midwest (83.6).

ADDITIONAL RESOURCES

The following were prepared by using HIV surveillance data:

- *MMWR* articles (selected): <http://www.cdc.gov/hiv/library/reports/mmwr.html>
- Other surveillance reports: <http://www.cdc.gov/hiv/library/reports/surveillance/>
- Public-use slides updated annually: <http://www.cdc.gov/hiv/library/slideSets/>
- NCHHSTP Atlas [interactive tool for accessing HIV/AIDS, STD, TB, and hepatitis data]: <http://www.cdc.gov/nchhstp/atlas/>

SUGGESTED READINGS

CDC. Establishing a holistic framework to reduce inequities in HIV, viral hepatitis, STDs, and tuberculosis in the United States. <http://stacks.cdc.gov/view/cdc/11585>. Published October 2010. Accessed January 6, 2015.

CDC. Estimated HIV incidence in the United States, 2007–2010. *HIV Surveillance Supplemental Report* 2012;17(No. 4). <http://go.usa.gov/p8P4>. Published December 2012. Accessed January 6, 2015.

CDC. HIV surveillance—United States, 1981–2008. *MMWR* 2011;60(21):689–693.

CDC. Monitoring selected national HIV prevention and care objectives by using HIV surveillance data—United States and 6 dependent areas—2012. *HIV Surveillance Supplemental Report* 2013;19(No. 3). <http://go.usa.gov/FUaw>. Published November 2014. Accessed January 6, 2015.

CDC. Revised surveillance case definitions for HIV infection among adults, adolescents, and children aged <18 months and for HIV infection and AIDS among children aged 18 months to <13 years—United States, 2008. *MMWR* 2008;57(RR-10):1–12.

CDC. Terms, definitions, and calculations used in CDC HIV surveillance publications. <http://www.cdc.gov/hiv/statistics/recommendations/terms.html>. Published September 2014. Accessed January 6, 2015.

CDC. Vital Signs: HIV diagnosis, care, and treatment among persons living with HIV—United States, 2011. *MMWR* 2014;63(47):1113–1117.

CDC. Vital Signs: HIV infection, testing, and risk behaviors among youths—United States. *MMWR* 2012;61(47):971–976.

CDC. Vital Signs: HIV prevention through care and treatment—United States. *MMWR* 2011;60(47):1618–1623.

CDC. Vital Signs: HIV testing and diagnosis among adults—United States. *MMWR* 2010;59(47):1550–1555.

Cohen SM, Gray KM, Ocfemia MC, Johnson AS, Hall HI. The status of the National HIV Surveillance System, United States, 2013. *Public Health Rep* 2014;129(4):335–341.

Hall HI, Song R, Rhodes P, et al. Estimation of HIV incidence in the United States. *JAMA* 2008;300(5):520–529.

Johnson AS, Hall HI, Hu X, Lansky A, Holtgrave DR, Mermin J. Trends in diagnoses of HIV infection in the United States, 2002–2011 [Research Letter]. *JAMA* 2014;312(4):432–434. doi:10.1001/jama.2014.8534.

Prejean J, Song R, Hernandez A, et al. Estimated HIV incidence in the United States, 2006–2009. *PLoS One* 2011;6(8):e17502. doi:10.1371/journal.pone.0017502.

SURVEILLANCE OF HIV INFECTION

This report includes case report data from 50 states, the District of Columbia, and 6 U.S. dependent areas (American Samoa, Guam, the Northern Mariana Islands, Puerto Rico, the Republic of Palau, and the U.S. Virgin Islands) in which laws or regulations require confidential reporting to the jurisdiction (not to CDC), by name for adults, adolescents, and children with confirmed diagnoses of HIV infection. After the removal of personal identifying information, data from these reports were submitted to CDC. Although AIDS cases have been reported to CDC since 1981, the implementation of HIV infection reporting has differed from state to state. All states, the District of Columbia, and 6 U.S. dependent areas had fully implemented name-based HIV infection reporting by April 2008.

Please use caution when interpreting data on diagnoses of HIV infection. HIV surveillance reports may not be representative of all persons with HIV because not all infected persons have been (1) tested or (2) tested at a time when the infection could be detected and diagnosed. Also, some states offer anonymous HIV testing; the results of anonymous tests are not reported to the confidential name-based HIV registries of state and local health departments. Therefore, reports of confidential test results may not represent all persons who tested positive for HIV infection. In addition, testing patterns are influenced by many factors, including the extent to which testing is routinely offered to specific groups, or the availability of, and access to, medical care and testing services. The data presented in this report provide a minimum estimate of persons for whom HIV infection has been diagnosed and reported to the surveillance system. Finally, because surveillance practices among jurisdictions differ, the reporting and the updating of a person's clinical and vital status differ among states. The completeness of reporting of HIV infection is estimated at more than 80% [1].

STAGES OF HIV INFECTION

The 2008 revised HIV case definition was used to classify HIV infection among adults and adolescents and among children [2]. The 2008 definition incorpo-

rates an HIV infection classification staging system that includes AIDS (HIV infection, stage 3). The stages of HIV infection are defined as follows:

- **HIV infection, stage 1:** No AIDS-defining condition and either CD4 count of ≥ 500 cells/ μ L or CD4 percentage of total lymphocytes of ≥ 29 .
- **HIV infection, stage 2:** No AIDS-defining condition and either CD4 count of 200–499 cells/ μ L or CD4 percentage of total lymphocytes of 14–28.
- **HIV infection, stage 3 (AIDS):** Documentation of an AIDS-defining condition or either a CD4 count of < 200 cells/ μ L or CD4 percentage of total lymphocytes of < 14 . Documentation of an AIDS-defining condition supersedes a CD4 count or percentage that would not, by itself, be the basis for a stage 3 (AIDS) classification.
- **HIV infection, stage unknown:** No reported information on AIDS-defining conditions and no information available on CD4 count or percentage.

In this report, the term *diagnosis of HIV infection* is defined as a diagnosis of HIV infection regardless of the stage of disease (stage 1, 2, 3 [AIDS], or unknown) and refers to all persons with a diagnosis of HIV infection. The term *HIV infection, stage 3 (AIDS)*, and its condensed version—*stage 3 (AIDS)*—refer specifically to persons with diagnosed HIV whose infection was classified as stage 3 (AIDS) during a given year (for diagnoses) or whose infection has ever been classified as stage 3 (AIDS) (for prevalence and deaths).

TABULATION AND PRESENTATION OF DATA

The data in this report include information received by CDC through June 30, 2014. The data are organized into 4 sections.

- Section 1 (Tables 1a/b–9): numbers (statistically adjusted [estimated] and unadjusted) and rates (adjusted) of diagnoses of HIV infection (any stage) and diagnoses of infection classified as stage 3 (AIDS)
- Section 2 (Tables 10a/b–13a/b): numbers (statistically adjusted and unadjusted) and rates (adjusted) of deaths and proportions (unadjusted) of survival among persons with diagnosed HIV infection (any

stage) and persons with infection ever classified as stage 3 (AIDS)

- Section 3 (Tables 14a/b–17a/b): numbers (statistically adjusted and unadjusted) and rates (adjusted) of prevalence (i.e., data on persons living with diagnosed HIV infection [any stage] and data on persons living with infection ever classified as stage 3 [AIDS])
- Section 4 (Tables 18–23): numbers (statistically adjusted and unadjusted) and rates (adjusted) of diagnoses and prevalence of HIV infection (any stage) and diagnoses and prevalence of infection classified as stage 3 (AIDS), presented by state and metropolitan statistical area (MSA)

Many tables in the 2013 report present data in 2 formats. Tables in the first format—labeled “a”—exclude data from the dependent areas (American Samoa, Guam, the Northern Mariana Islands, Puerto Rico, the Republic of Palau, and the U.S. Virgin Islands). Tables in the second format—labeled “b”—include data from the dependent areas.

Please use caution when interpreting reported numbers less than 12, as well as estimated numbers and accompanying rates and trends based on these numbers: the underlying relative standard errors are greater than 30%, so the numbers are considered unreliable. For the assessment of trends in diagnoses, deaths, and prevalence, it is preferable to use statistically adjusted (estimated) data to eliminate artifacts of reporting in the surveillance system.

Diagnoses of HIV Infection and of Infection Classified as Stage 3 (AIDS)

As of April 2008, 57 areas (50 states, the District of Columbia, and 6 U.S. dependent areas) had implemented confidential name-based HIV infection reporting. All 57 areas were included in tables displaying numbers (unadjusted), estimated numbers, and estimated rates of diagnoses of HIV infection and of diagnoses of infection classified as stage 3 (AIDS), by selected characteristics, area of residence, and MSA (Tables 1a/b–9, 12a/b, 13a/b, 18, 19, 22, and 23).

Deaths

Persons reported to the National HIV Surveillance System are assumed alive unless their deaths have been reported to CDC. Death data include deaths of persons with diagnosed HIV infection (Tables 10a/b) or with infection ever classified as stage 3 (AIDS)

(Tables 11a/b) regardless of the cause of death.

Because of delays in the reporting of deaths, 4 years (2009–2012) of death data are displayed. The exclusion of data from the most recent year allowed at least 18 months for deaths to be reported to CDC. The estimated numbers and rates of deaths resulted from statistical adjustment for delays in reporting. Please use caution when interpreting trend data on the estimated numbers of deaths: the estimates for the most recent year are most subject to uncertainty.

Survival Analyses

The Kaplan-Meier method was used to estimate the probability of survival for 12, 24, and 36 months after a diagnosis of HIV infection (Tables 12a/b) or after a diagnosis of infection classified as stage 3 (AIDS) (Tables 13a/b). To allow at least 3 years from the time of diagnosis to deaths occurring through December 31, 2012 (reported to CDC by June 30, 2014), table data were limited to persons whose diagnosis was made during 2004–2009. Data for each HIV reporting area were included in the survival tables beginning with the first full calendar year after implementation of code-based or name-based HIV infection reporting.

Prevalence: Persons Living with Diagnosed HIV Infection and Persons Living with Infection Ever Classified as Stage 3 (AIDS)

Because of delays in the reporting of deaths, 4 years of prevalence data are displayed. The data reflect the persons living with diagnosed HIV infection (Tables 14a/b, 16a/b, 20, and 22) and persons living with infection ever classified as stage 3 (AIDS) (Tables 15a/b, 17a/b, 21, and 23) at the end of a given year during 2009–2012. The exclusion of data from the most recent year allowed at least 18 months for deaths to be reported and for these deaths to be factored into calculations of prevalence.

Age

The designation “adults and adolescents” refers to persons aged 13 years and older; the designation “children” refers to persons aged less than 13 years. For presentations of data on persons living with diagnosed HIV infection (Tables 14a/b, 16a/b, and 20) and presentations of data on persons living with infection classified as stage 3 (AIDS) (Tables 15a/b, 17a/b, and 21), the age-group assignment (for example, 20–24 years) or the age designation (for example, “adults and

adolescents”) is based on the person’s age as of December 31 of the specified year. For presentations of data on deaths of persons with diagnosed HIV infection (Tables 10a/b) and deaths of persons with infection classified as stage 3 (AIDS) (Tables 11a/b), age-group assignment is determined by the person’s age at the time of death. For all other tables, the age designation or the specific age-group assignment is based on the person’s age at the time of diagnosis (HIV infection or infection classified as stage 3 [AIDS]).

Age-group designations for transmission categories in tables displaying data on prevalence and on deaths are based on a person’s age at the end of the specified year or age at death, respectively. That is, the transmission categories among “adults and adolescents” in these tables include persons who were children at the time of diagnosis but who were aged 13 years and older during the specified year or at death.

Sex

Sex designations in this report are based on a person’s sex at birth. *Transgender* is an umbrella term that is used to identify persons whose sex assigned at birth does not match current gender identity or expression. Data for transgender persons are not presented in this report because information on gender identity (a person’s understanding of his or her gender or the gender with which a person identifies) is not consistently collected or documented in the data sources used by HIV reporting jurisdictions. HIV surveillance personnel voluntarily collect this information, when available, from sources such as case report forms submitted by health care or HIV testing providers and medical records, or by matching with other databases (e.g., Ryan White program data) within the health department. In May 2012, CDC issued guidance to state and local programs on methods for collecting data on transgender persons and working with transgender-specific data. However, fully characterizing the HIV burden among transgender persons may require that HIV surveillance data be supplemented with data from special studies.

Race and Ethnicity

In the *Federal Register* [3] for October 30, 1997, the Office of Management and Budget (OMB) announced the Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity. Implementation by January 1, 2003, was mandated.

At a minimum, data on the following race categories should be collected:

- American Indian or Alaska Native
- Asian
- black or African American
- Native Hawaiian or other Pacific Islander
- white

Additionally, systems must be able to retain information when multiple race categories are reported. In addition to data on race, data on 2 categories of ethnicity should be collected:

- Hispanic or Latino
- not Hispanic or Latino

The Asian or Pacific Islander category displayed in annual surveillance reports published prior to the 2007 surveillance report was split into 2 categories: (1) Asian and (2) Native Hawaiian or other Pacific Islander. The Asian category (in tables where footnoted) includes the cases in Asians/Pacific Islanders (referred to as legacy cases) that were reported before the implementation of the new race categories in 2003 (e.g., cases of HIV infection that were diagnosed and reported to CDC before 2003 but that progressed to AIDS after 2003) and a small percentage of cases that were reported after 2003 but that were reported according to the old race category (Asian/Pacific Islander). In tables of diagnoses of HIV infection during 2009–2013, the Asian category does not include Asian/Pacific Islander cases because these cases were diagnosed after 2003 and were reported to CDC in accordance with OMB’s Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity [3].

This report also presents data for persons for whom multiple race categories are reported. Persons whose race was unknown are included in the total numbers in Tables 10a/b–17a/b, 20, and 21, and in the cumulative totals of Tables 2a/b and 6a/b. In this report, persons categorized by race were not Hispanic or Latino. The number of persons reported in each race category may, however, include persons whose ethnicity was not reported.

Geographic Designations

Geographic designations reflect the area of residence at the time of diagnosis of HIV infection or at the time of stage 3 (AIDS) classification. The 4 regions

of residence used in this report are defined by the U.S. Census Bureau as follows:

Northeast: Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, and Vermont

Midwest: Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin

South: Alabama, Arkansas, Delaware, District of Columbia, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, and West Virginia

West: Alaska, Arizona, California, Colorado, Hawaii, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, and Wyoming

U.S. dependent areas: American Samoa, Guam, the Northern Mariana Islands, Puerto Rico, the Republic of Palau, and the U.S. Virgin Islands

Metropolitan Statistical Areas

In the *Federal Register* for June 28, 2010, the OMB published revised standards for defining MSAs in federal statistical activities [4]. These standards, which provided for the identification of MSAs in the United States and Puerto Rico, replaced the 2000 standards. The adoption of the new standards was effective as of June 28, 2010. On February 28, 2013, the OMB announced new MSA delineations based on the new standards and Census 2010 data [5]. Tables 22 (data on diagnosed HIV infection) and 23 (data on stage 3 [AIDS]) present numbers and rates of diagnoses and of prevalence, by MSA, for areas with populations of 500,000 or more. The MSAs listed in these tables were defined according to the OMB's most recent update (February 2013) of statistical areas [5].

Transmission Categories

Transmission category is the term for the classification of cases that summarizes a person's possible HIV risk factors; the summary classification results from selecting, from the presumed hierarchical order of probability, the single risk factor most likely to have been responsible for transmission. For surveillance purposes, a diagnosis of HIV infection is counted only once in the hierarchy of transmission categories. Persons with more than 1 reported risk factor for HIV infection are classified in the transmission category listed first in the hierarchy. The exception is men who

had sexual contact with other men *and* injected drugs; this group makes up a separate transmission category.

Persons whose transmission category is classified as male-to-male sexual contact include men who have ever had sexual contact with other men (i.e., homosexual contact) and men who have ever had sexual contact with both men and women (i.e., bisexual contact). Persons whose transmission category is classified as heterosexual contact are persons who have ever had heterosexual contact with a person known to have, or to be at high risk for, HIV infection (e.g., an injection drug user).

Tables of prevalence data and tables of death data include a separate transmission category of adults and adolescents with HIV infection attributed to perinatal transmission. The category includes persons infected through perinatal transmission but aged 13 years and older during the specified year or at death.

Cases of HIV infection reported without a risk factor listed in the hierarchy of transmission categories are classified as "no identified risk (NIR)" [6]. Cases classified as NIR include cases that are being followed up by local health department staff; cases in persons whose risk-factor information is missing because they died, declined to be interviewed, or were lost to follow-up; and cases in persons who were interviewed or for whom other follow-up information was available but for whom no risk factor was identified.

Because a substantial proportion of cases of HIV infection are reported to CDC without an identified risk factor, multiple imputation is used to assign a transmission category [7]. Multiple imputation is a statistical approach in which each missing transmission category is replaced with a set of plausible values that represent the uncertainty about the true, but missing, value [8]. The plausible values are analyzed by using standard procedures, and the results from these analyses are then combined to produce the final results. In tables displaying transmission categories, multiple imputation was used for adults and adolescents, but not for children (because the number of cases in children is small, missing transmission categories were not imputed).

Reporting Delays

Reporting delays (time between diagnosis or death and the reporting of diagnosis or death to CDC) may differ among demographic and geographic categories; for some, delays in reporting have been as long as several

years. The statistical adjustment of the data on diagnoses and deaths is based on estimates of reporting-delay distributions, which are calculated by using a modified semiparametric life-table statistical procedure. This procedure takes into account differences in reporting delays due to sex, race/ethnicity, HIV transmission categories, geographic area (i.e., reporting city, state, or dependent area; region of residence), the size of the MSA, and the type of facility where the diagnosis was made or death occurred [9]. The statistical adjustment of data on diagnoses takes into consideration the information reported from the state or dependent area where the diagnosis was made; the statistical adjustment of data on deaths takes into consideration the information from the state where the death occurred.

Rates

Rates per 100,000 population were calculated for (1) the numbers of diagnoses of HIV infection and the numbers of infections classified as stage 3 (AIDS), (2) the numbers of deaths of persons with diagnosed HIV infection and deaths of persons with infection ever classified as stage 3 (AIDS), and (3) the numbers of persons living with diagnosed HIV infection and persons living with infection ever classified as stage 3 (AIDS).

The population denominators used to compute the rates for the 50 states, the District of Columbia, and Puerto Rico were based on the Vintage 2009 postcensal estimates file (for the year 2009) and the Vintage 2013 file (for years 2010–2013) from the U.S. Census Bureau [10]. The population denominators for American Samoa, Guam, the Northern Mariana Islands, the Republic of Palau, and the U.S. Virgin Islands were based on estimates and projections from the U.S. Census Bureau's International Data Base [11]. Each rate was calculated by dividing the estimated total number of diagnoses (or deaths or prevalence) for the calendar year by the population for that calendar year and then multiplying the result by 100,000. The denominators used for calculating the rates specific to age, sex, and race/ethnicity were computed by applying the appropriate vintage estimates for age, sex, and race/ethnicity for the 50 states and the District of Columbia [10]. The same method was used to calculate the denominators for Puerto Rico, with the exception of race/ethnicity estimates; these data are not available for Puerto Rico (see next paragraph). For the other 5 U.S. dependent areas, estimates from the U.S. Census

Bureau's International Data Base were used for age- and sex-specific population denominators [11].

CDC currently does not provide subpopulation rates by race/ethnicity for the 6 U.S. dependent areas because the U.S. Census Bureau does not collect information from all dependent areas. Rates for transmission categories are not provided in this report because of the absence of denominator data from the U.S. Census Bureau, the source of data used for calculating all rates in this report.

REFERENCES

1. Hall HI, Song R, Gerstle JE III, Lee LM; on behalf of the HIV/AIDS Reporting System Evaluation Group. Assessing the completeness of reporting of human immunodeficiency virus diagnoses in 2002–2003: capture-recapture methods. *Am J Epidemiol* 2006;164(4):391–397.
2. CDC. Revised surveillance case definitions for HIV infection among adults, adolescents, and children aged <18 months and for HIV infection and AIDS among children aged 18 months to <13 years—United States, 2008. *MMWR* 2008;57(RR-10):1–12.
3. Office of Management and Budget. Revisions to the standards for the classification of federal data on race and ethnicity. *Federal Register* 1997;62:58781–58790. <http://go.usa.gov/vSdR>. Accessed January 6, 2015.
4. Office of Management and Budget. 2010 Standards for defining metropolitan and micropolitan statistical areas. *Federal Register* 2010;75(123):37246–39052. <http://go.usa.gov/vSpG>. Accessed January 6, 2015.
5. Office of Management and Budget. Revised definitions of metropolitan statistical areas, new definitions of micropolitan statistical areas and combined statistical areas, and guidance on uses of the delineations of these areas. OMB Bulletin 13-01. <http://go.usa.gov/vSyB>. Published February 28, 2013. Accessed January 6, 2015.
6. CDC. Current Trends: heterosexually acquired AIDS—United States, 1993. *MMWR* 1994;43(9):158, Fig 3.
7. Harrison KM, Kajese T, Hall HI, Song R. Risk factor redistribution of the national HIV/AIDS surveillance data: an alternative approach. *Public Health Rep* 2008;123(5):618–627.
8. Rubin, DB. *Multiple Imputation for Nonresponse in Surveys*. New York: John Wiley & Sons Inc; 1987.
9. Song R, Green TA. An improved approach to accounting for reporting delay in case surveillance systems. *JP J Biostat* 2012;7(1):1–14.

10. U.S. Census Bureau. Population estimates [entire data set]. <http://www.census.gov/popest/data/>. Updated July 1, 2014. Accessed January 6, 2015.
11. U.S. Census Bureau. International Data Base. <http://go.usa.gov/vSUj>. Updated December 2013. Accessed January 6, 2015.

Table 1a. Diagnoses of HIV infection, by year of diagnosis and selected characteristics, 2009–2013—United States

	2009			2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	No.		Rate	No.		No.	Rate		No.	No.
Age at diagnosis (yr)															
<13	218	223	0.4	225	232	0.4	187	196	0.4	232	250	0.5	164	187	0.4
13–14	30	31	0.4	40	41	0.5	44	47	0.6	49	53	0.6	39	45	0.5
15–19	2,187	2,228	10.3	2,069	2,123	9.7	1,988	2,074	9.6	1,846	1,989	9.3	1,652	1,863	8.8
20–24	6,741	6,875	31.9	7,064	7,265	33.5	7,066	7,386	33.3	7,196	7,752	34.3	7,059	8,053	35.3
25–29	6,510	6,641	30.6	6,342	6,533	30.9	6,366	6,658	31.3	6,556	7,083	33.1	6,844	7,825	36.3
30–34	5,714	5,835	29.3	5,481	5,652	28.2	5,262	5,508	26.9	5,589	6,037	28.9	5,404	6,165	29.0
35–39	5,659	5,779	28.1	5,028	5,184	25.8	4,456	4,673	23.8	4,264	4,611	23.7	4,246	4,858	24.8
40–44	5,991	6,120	29.2	5,233	5,402	25.8	4,800	5,032	23.9	4,554	4,923	23.4	4,217	4,820	23.1
45–49	5,297	5,413	23.7	4,860	5,014	22.1	4,628	4,859	21.9	4,449	4,822	22.2	4,311	4,961	23.4
50–54	3,659	3,742	17.2	3,512	3,629	16.2	3,364	3,532	15.6	3,274	3,554	15.7	3,254	3,747	16.6
55–59	2,163	2,212	11.7	2,065	2,137	10.8	1,992	2,095	10.3	1,975	2,146	10.3	2,151	2,467	11.6
60–64	1,007	1,031	6.5	1,073	1,109	6.5	1,069	1,123	6.3	1,066	1,164	6.5	1,150	1,316	7.3
≥65	823	842	2.1	789	815	2.0	812	858	2.1	825	901	2.1	896	1,045	2.3
Race/ethnicity															
American Indian/Alaska Native	167	169	7.2	180	183	8.1	172	177	7.8	194	202	8.8	197	218	9.4
Asian	718	734	5.4	717	742	5.0	787	827	5.4	833	901	5.8	862	973	6.0
Black/African American	21,661	22,136	58.7	20,510	21,148	55.6	19,453	20,436	53.3	19,079	20,803	53.7	18,803	21,836	55.9
Hispanic/Latino ^b	9,411	9,620	19.9	9,009	9,317	18.4	8,852	9,299	17.9	8,980	9,710	18.3	8,878	10,117	18.7
Native Hawaiian/Other Pacific Islander	64	65	14.5	59	61	12.1	56	59	11.5	71	75	14.4	61	67	12.7
White	12,518	12,750	6.4	11,957	12,290	6.2	11,465	11,923	6.0	11,593	12,372	6.3	11,672	13,101	6.6
Multiple races	1,460	1,497	32.8	1,349	1,395	24.7	1,249	1,318	22.6	1,125	1,223	20.4	914	1,039	16.8
Transmission category															
Male adult or adolescent															
Male-to-male sexual contact	21,811	27,394	—	21,712	27,106	—	21,791	27,357	—	21,962	28,967	—	21,498	30,689	—
Injection drug use	1,507	2,501	—	1,306	2,205	—	1,076	1,879	—	922	1,799	—	887	1,942	—
Male-to-male sexual contact and injection drug use	1,251	1,611	—	1,177	1,507	—	1,024	1,346	—	978	1,316	—	851	1,270	—
Heterosexual contact ^c	3,120	4,501	—	2,871	4,176	—	2,725	3,959	—	2,452	3,776	—	2,199	3,887	—
Other ^d	7,625	56	—	6,935	55	—	6,415	53	—	6,953	79	—	7,745	99	—
Subtotal	35,314	36,062	29.1	34,001	35,049	28.0	33,031	34,595	27.4	33,267	35,937	28.2	33,180	37,887	29.4

Table 1a. Diagnoses of HIV infection, by year of diagnosis and selected characteristics, 2009–2013—United States (cont)

	2009			2010			2011			2012			2013		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
Female adult or adolescent															
Injection drug use	931	1,687	—	777	1,426	—	661	1,288	—	599	1,227	—	510	1,154	—
Heterosexual contact ^c	5,014	8,943	—	4,669	8,382	—	4,200	7,905	—	3,800	7,811	—	3,446	8,031	—
Other ^d	4,522	56	—	4,109	46	—	3,955	57	—	3,977	61	—	4,087	93	—
Subtotal	10,467	10,686	8.3	9,555	9,855	7.5	8,816	9,250	7.0	8,376	9,099	6.8	8,043	9,278	6.9
Child (<13 yrs at diagnosis)															
Perinatal	175	179	—	175	180	—	135	142	—	154	165	—	93	107	—
Other ^e	43	44	—	50	52	—	52	54	—	78	85	—	71	80	—
Subtotal	218	223	0.4	225	232	0.4	187	196	0.4	232	250	0.5	164	187	0.4
Region of residence															
Northeast	9,006	9,209	16.7	8,421	8,711	15.7	7,865	8,307	14.9	7,829	8,629	15.5	7,499	8,908	15.9
Midwest	5,812	5,917	8.9	5,559	5,698	8.5	5,444	5,634	8.4	5,608	5,906	8.8	5,600	6,109	9.0
South	23,032	23,513	20.7	21,895	22,564	19.6	21,244	22,285	19.2	20,804	22,629	19.3	21,066	24,323	20.5
West	8,149	8,332	11.6	7,906	8,163	11.3	7,481	7,814	10.7	7,634	8,122	11.0	7,222	8,013	10.8
Total^f	45,999	46,971	15.3	43,781	45,136	14.6	42,034	44,040	14.1	41,875	45,287	14.4	41,387	47,352	15.0

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b Hispanics/Latinos can be of any race.

^c Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^d Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 1b. Diagnoses of HIV infection, by year of diagnosis and selected characteristics, 2009–2013—United States and 6 dependent areas

	2009			2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Age at diagnosis (yr)															
<13	219	224	0.4	229	236	0.4	190	199	0.4	232	250	0.5	164	187	0.4
13–14	31	32	0.4	40	41	0.5	44	47	0.6	50	54	0.6	40	46	0.5
15–19	2,206	2,248	10.3	2,097	2,154	9.7	2,009	2,098	9.6	1,865	2,011	9.3	1,670	1,885	8.8
20–24	6,824	6,963	31.9	7,138	7,345	33.4	7,141	7,470	33.3	7,302	7,876	34.4	7,132	8,144	35.3
25–29	6,610	6,747	30.7	6,436	6,635	31.0	6,461	6,765	31.4	6,653	7,197	33.2	6,911	7,909	36.2
30–34	5,844	5,973	29.6	5,592	5,772	28.4	5,353	5,610	27.0	5,686	6,151	29.1	5,491	6,274	29.2
35–39	5,770	5,897	28.3	5,127	5,291	26.0	4,540	4,767	24.0	4,330	4,688	23.7	4,308	4,936	24.9
40–44	6,125	6,263	29.4	5,348	5,527	26.1	4,897	5,142	24.1	4,642	5,026	23.6	4,284	4,903	23.2
45–49	5,401	5,524	23.9	4,970	5,134	22.4	4,733	4,976	22.2	4,545	4,935	22.5	4,394	5,066	23.6
50–54	3,739	3,828	17.4	3,581	3,704	16.4	3,449	3,627	15.9	3,342	3,634	15.9	3,335	3,850	16.9
55–59	2,215	2,267	11.8	2,110	2,186	10.9	2,040	2,149	10.5	2,014	2,192	10.4	2,193	2,519	11.7
60–64	1,030	1,056	6.6	1,109	1,148	6.7	1,092	1,148	6.4	1,091	1,194	6.6	1,180	1,354	7.4
≥65	856	877	2.2	818	847	2.1	837	886	2.1	864	946	2.2	916	1,070	2.4
Race/ethnicity															
American Indian/Alaska Native	167	169	—	180	183	—	172	177	—	194	202	—	197	218	—
Asian	718	734	—	719	744	—	787	827	—	833	901	—	862	973	—
Black/African American	21,680	22,156	—	20,524	21,163	—	19,470	20,455	—	19,084	20,809	—	18,816	21,853	—
Hispanic/Latino ^b	10,253	10,515	—	9,802	10,179	—	9,584	10,120	—	9,710	10,565	—	9,493	10,888	—
Native Hawaiian/Other Pacific Islander	66	67	—	62	64	—	58	61	—	72	76	—	61	67	—
White	12,523	12,755	—	11,959	12,292	—	11,466	11,925	—	11,597	12,377	—	11,675	13,105	—
Multiple races	1,463	1,500	—	1,349	1,395	—	1,249	1,318	—	1,126	1,224	—	914	1,039	—
Transmission category															
Male adult or adolescent															
Male-to-male sexual contact	22,058	27,668	—	21,985	27,422	—	22,042	27,658	—	22,239	29,310	—	21,746	31,023	—
Injection drug use	1,664	2,687	—	1,420	2,351	—	1,172	2,008	—	999	1,904	—	958	2,051	—
Male-to-male sexual contact and injection drug use	1,279	1,643	—	1,199	1,533	—	1,038	1,364	—	993	1,336	—	861	1,284	—
Heterosexual contact ^c	3,264	4,667	—	3,008	4,337	—	2,855	4,120	—	2,587	3,946	—	2,297	4,021	—
Other ^d	7,669	56	—	6,986	55	—	6,467	54	—	6,993	79	—	7,789	99	—
Subtotal	35,934	36,721	29.2	34,598	35,698	28.2	33,574	35,204	27.5	33,811	36,575	28.3	33,651	38,479	29.5

Table 1b. Diagnoses of HIV infection, by year of diagnosis and selected characteristics, 2009–2013—United States and 6 dependent areas (*cont*)

	2009			2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Female adult or adolescent															
Injection drug use	964	1,725	—	796	1,449	—	692	1,325	—	618	1,253	—	533	1,188	—
Heterosexual contact ^c	5,210	9,170	—	4,842	8,591	—	4,361	8,099	—	3,944	8,015	—	3,546	8,195	—
Other ^d	4,543	56	—	4,130	46	—	3,969	57	—	4,011	61	—	4,124	95	—
Subtotal	10,717	10,952	8.3	9,768	10,086	7.6	9,022	9,481	7.1	8,573	9,329	6.9	8,203	9,479	6.9
Child (<13 yrs at diagnosis)															
Perinatal	176	180	—	177	182	—	137	144	—	154	165	—	93	107	—
Other ^e	43	44	—	52	54	—	53	55	—	78	85	—	71	80	—
Subtotal	219	224	0.4	229	236	0.4	190	199	0.4	232	250	0.5	164	187	0.4
Region of residence															
Northeast	9,006	9,209	16.7	8,421	8,711	15.7	7,865	8,307	14.9	7,829	8,629	15.5	7,499	8,908	15.9
Midwest	5,812	5,917	8.9	5,559	5,698	8.5	5,444	5,634	8.4	5,608	5,906	8.8	5,600	6,109	9.0
South	23,032	23,513	20.7	21,895	22,564	19.6	21,244	22,285	19.2	20,804	22,629	19.3	21,066	24,323	20.5
West	8,149	8,332	11.6	7,906	8,163	11.3	7,481	7,814	10.7	7,634	8,122	11.0	7,222	8,013	10.8
U.S. dependent areas	871	926	21.1	814	885	21.5	752	843	20.7	741	868	21.5	631	792	19.8
Total^f	46,870	47,897	15.4	44,595	46,021	14.7	42,786	44,883	14.2	42,616	46,154	14.5	42,018	48,145	15.0

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited. Rates are not calculated by transmission category because of the lack of denominator data.

^b Hispanics/Latinos can be of any race.

^c Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^d Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 2a. Stage 3 (AIDS), by year of diagnosis and selected characteristics, 2009–2013 and cumulative—United States

	2009			2010			2011			2012			2013			Cumulative ^b	
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Est. No. ^a
		No.	Rate		No.	No.		Rate	No.		No.	Rate		No.	No.		
Age at diagnosis (yr)																	
<13	14	14	0.0	23	24	0.0	15	16	0.0	10	10	0.0	7	8	0.0	9,399	9,421
13–14	45	46	0.6	47	49	0.6	37	39	0.5	28	29	0.4	25	30	0.4	1,442	1,461
15–19	439	450	2.1	454	470	2.1	428	447	2.1	340	361	1.7	381	435	2.1	8,636	8,793
20–24	1,947	1,996	9.3	1,987	2,062	9.5	1,971	2,064	9.3	1,863	1,982	8.8	1,996	2,239	9.8	50,040	50,787
25–29	3,140	3,220	14.9	2,884	2,993	14.2	2,763	2,896	13.6	2,701	2,877	13.4	2,787	3,123	14.5	138,724	139,957
30–34	3,651	3,743	18.8	3,302	3,423	17.1	3,129	3,276	16.0	3,218	3,424	16.4	2,927	3,268	15.4	222,750	224,308
35–39	4,335	4,442	21.6	3,708	3,844	19.1	3,179	3,329	17.0	2,891	3,074	15.8	2,869	3,200	16.3	241,443	243,169
40–44	4,989	5,116	24.4	4,304	4,463	21.3	3,753	3,930	18.7	3,424	3,640	17.3	3,121	3,496	16.8	202,176	204,006
45–49	4,813	4,935	21.6	4,273	4,431	19.6	3,953	4,138	18.7	3,614	3,843	17.7	3,386	3,781	17.8	137,405	139,062
50–54	3,492	3,578	16.4	3,180	3,298	14.8	2,900	3,043	13.5	2,863	3,045	13.5	2,803	3,135	13.9	81,518	82,718
55–59	1,939	1,987	10.5	1,759	1,826	9.2	1,793	1,874	9.2	1,756	1,872	9.0	1,789	1,998	9.4	44,927	45,628
60–64	917	939	5.9	933	967	5.7	910	950	5.3	984	1,049	5.9	1,019	1,144	6.3	23,899	24,274
≥65	777	796	2.0	741	767	1.9	724	754	1.8	721	766	1.8	740	831	1.9	20,169	20,455
Race/ethnicity																	
American Indian/Alaska Native	104	106	4.5	119	122	5.4	114	117	5.1	101	105	4.5	96	104	4.5	3,486	3,514
Asian ^c	374	384	2.8	365	380	2.6	367	384	2.5	358	382	2.4	374	415	2.6	9,560	9,712
Black/African American	14,326	14,687	39.0	13,440	13,951	36.7	12,353	12,946	33.7	11,949	12,735	32.9	11,678	13,172	33.7	491,715	497,267
Hispanic/Latino ^d	6,454	6,629	13.7	5,599	5,817	11.5	5,215	5,469	10.5	4,981	5,307	10.0	4,814	5,336	9.9	213,246	215,685
Native Hawaiian/Other Pacific Islander	37	38	8.4	39	40	8.1	32	33	6.5	30	32	6.1	34	37	6.9	845	855
White	7,917	8,097	4.1	6,979	7,208	3.7	6,505	6,786	3.4	6,079	6,429	3.3	6,090	6,759	3.4	433,681	436,557
Multiple races	1,286	1,322	29.0	1,054	1,097	19.4	969	1,020	17.5	915	984	16.4	764	867	14.0	29,995	30,448
Transmission category																	
Male adult or adolescent																	
Male-to-male sexual contact	12,574	15,530	—	11,635	14,575	—	11,157	13,958	—	10,700	13,821	—	10,542	14,611	—	520,784	577,403
Injection drug use	1,741	2,432	—	1,482	2,152	—	1,242	1,825	—	1,105	1,705	—	932	1,610	—	164,750	187,218
Male-to-male sexual contact and injection drug use	1,322	1,595	—	1,130	1,389	—	977	1,232	—	860	1,115	—	716	1,026	—	77,304	83,828
Heterosexual contact ^e	2,626	3,549	—	2,239	3,155	—	2,093	2,918	—	1,891	2,790	—	1,774	2,865	—	64,842	82,447
Other ^f	4,398	125	—	4,138	116	—	3,696	122	—	3,821	115	—	4,164	144	—	106,136	11,545
Subtotal	22,661	23,232	18.7	20,624	21,387	17.1	19,165	20,055	15.9	18,377	19,546	15.3	18,128	20,256	15.7	933,816	942,440

Table 2a. Stage 3 (AIDS), by year of diagnosis and selected characteristics, 2009–2013 and cumulative—United States (cont)

	2009			2010			2011			2012			2013			Cumulative ^b	
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a			Cumulative ^b	
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	Est. No. ^a
Female adult or adolescent																	
Injection drug use	1,160	1,734	—	959	1,472	—	839	1,313	—	755	1,225	—	645	1,143	—	74,970	89,790
Heterosexual contact ^e	4,108	6,156	—	3,584	5,599	—	3,265	5,240	—	3,003	5,077	—	2,696	5,109	—	111,561	146,521
Other ^f	2,555	125	—	2,405	136	—	2,271	132	—	2,268	114	—	2,374	172	—	52,782	5,868
Subtotal	7,823	8,016	6.2	6,948	7,206	5.5	6,375	6,684	5.0	6,026	6,417	4.8	5,715	6,424	4.8	239,313	242,178
Child (<13 yrs at diagnosis)																	
Perinatal	13	13	—	17	18	—	12	13	—	9	9	—	6	7	—	8,532	8,553
Other ^g	1	1	—	6	6	—	3	3	—	1	1	—	1	1	—	867	869
Subtotal	14	14	0.0	23	24	0.0	15	16	0.0	10	10	0.0	7	8	0.0	9,399	9,421
Region of residence																	
Northeast	6,516	6,750	12.2	5,639	5,923	10.7	5,119	5,449	9.8	4,771	5,173	9.3	4,301	4,872	8.7	348,674	352,167
Midwest	3,652	3,714	5.6	3,369	3,456	5.2	3,249	3,356	5.0	3,130	3,283	4.9	2,944	3,221	4.8	125,360	126,372
South	14,718	15,036	13.3	13,715	14,183	12.3	12,908	13,466	11.6	12,382	13,109	11.2	12,769	14,345	12.1	473,007	477,964
West	5,612	5,761	8.0	4,872	5,054	7.0	4,279	4,484	6.2	4,130	4,408	6.0	3,836	4,251	5.7	235,487	237,536
Total^h	30,498	31,262	10.2	27,595	28,616	9.3	25,555	26,755	8.6	24,413	25,973	8.3	23,850	26,688	8.4	1,182,528	1,194,039

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b From the beginning of the epidemic through 2013.

^c Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^d Hispanics/Latinos can be of any race.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^g Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^h Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 2b. Stage 3 (AIDS), by year of diagnosis and selected characteristics, 2009–2013 and cumulative—United States and 6 dependent areas

	2009			2010			2011			2012			2013			Cumulative ^b	
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Est. No. ^a
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		No.	Rate		
Age at diagnosis (yr)																	
<13	14	14	0.0	23	24	0.0	16	17	0.0	10	10	0.0	7	8	0.0	9,820	9,843
13–14	45	46	0.6	47	49	0.6	38	40	0.5	29	31	0.4	26	31	0.4	1,500	1,520
15–19	444	455	2.1	458	474	2.1	436	456	2.1	341	362	1.7	384	439	2.0	8,868	9,027
20–24	1,964	2,013	9.2	2,005	2,080	9.5	1,986	2,080	9.3	1,882	2,003	8.8	2,016	2,266	9.8	51,441	52,204
25–29	3,196	3,277	14.9	2,916	3,026	14.1	2,797	2,932	13.6	2,733	2,912	13.5	2,810	3,152	14.4	142,921	144,180
30–34	3,720	3,813	18.9	3,350	3,473	17.1	3,174	3,324	16.0	3,257	3,467	16.4	2,960	3,311	15.4	229,616	231,215
35–39	4,416	4,525	21.7	3,782	3,921	19.3	3,250	3,404	17.1	2,943	3,132	15.9	2,903	3,244	16.3	248,612	250,387
40–44	5,103	5,232	24.6	4,393	4,555	21.5	3,824	4,005	18.8	3,488	3,710	17.4	3,164	3,551	16.8	207,924	209,804
45–49	4,924	5,048	21.8	4,379	4,540	19.8	4,047	4,238	18.9	3,696	3,934	17.9	3,440	3,853	17.9	141,291	143,001
50–54	3,577	3,665	16.6	3,252	3,373	14.9	2,980	3,128	13.7	2,920	3,107	13.6	2,862	3,213	14.1	83,820	85,064
55–59	1,987	2,036	10.6	1,800	1,868	9.3	1,825	1,908	9.3	1,791	1,910	9.1	1,824	2,045	9.5	46,287	47,014
60–64	943	966	6.0	963	998	5.8	926	967	5.4	1,003	1,070	5.9	1,036	1,167	6.4	24,665	25,054
≥65	804	823	2.1	761	788	1.9	755	787	1.9	748	796	1.8	759	854	1.9	21,098	21,398
Race/ethnicity																	
American Indian/Alaska Native	104	106	—	119	122	—	114	117	—	101	105	—	96	104	—	3,487	3,515
Asian ^c	374	384	—	365	380	—	368	385	—	358	382	—	374	415	—	9,617	9,770
Black/African American	14,341	14,702	—	13,449	13,961	—	12,360	12,953	—	11,960	12,747	—	11,683	13,179	—	492,184	497,742
Hispanic/Latino ^d	7,073	7,262	—	6,120	6,355	—	5,703	5,986	—	5,395	5,763	—	5,148	5,773	—	247,904	250,671
Native Hawaiian/Other Pacific Islander	38	39	—	41	42	—	34	35	—	31	33	—	34	37	—	861	872
White	7,921	8,101	—	6,981	7,210	—	6,505	6,786	—	6,081	6,431	—	6,092	6,762	—	433,803	436,681
Multiple races	1,286	1,322	—	1,054	1,097	—	970	1,021	—	915	984	—	764	867	—	30,007	30,460
Transmission category																	
Male adult or adolescent																	
Male-to-male sexual contact	12,708	15,670	—	11,748	14,694	—	11,274	14,084	—	10,817	13,953	—	10,636	14,739	—	526,753	583,510
Injection drug use	1,900	2,599	—	1,610	2,287	—	1,340	1,934	—	1,185	1,796	—	992	1,694	—	178,022	200,761
Male-to-male sexual contact and injection drug use	1,350	1,624	—	1,157	1,417	—	996	1,253	—	877	1,133	—	725	1,036	—	79,800	86,354
Heterosexual contact ^e	2,730	3,662	—	2,335	3,258	—	2,196	3,030	—	1,964	2,874	—	1,853	2,974	—	68,789	86,531
Other ^f	4,415	128	—	4,150	120	—	3,709	125	—	3,832	117	—	4,177	146	—	106,670	11,745
Subtotal	23,103	23,684	18.8	21,000	21,775	17.2	19,515	20,426	16.0	18,675	19,873	15.4	18,383	20,590	15.8	960,034	968,901

Table 2b. Stage 3 (AIDS), by year of diagnosis and selected characteristics, 2009–2013 and cumulative—United States and 6 dependent areas (cont)

	2009			2010			2011			2012			2013			Cumulative ^b	
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Est. No. ^a
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		No.	Rate		
Female adult or adolescent																	
Injection drug use	1,204	1,780	—	989	1,503	—	866	1,342	—	777	1,250	—	661	1,165	—	77,959	92,845
Heterosexual contact ^e	4,255	6,310	—	3,706	5,729	—	3,378	5,362	—	3,100	5,193	—	2,758	5,198	—	116,964	152,110
Other ^f	2,561	127	—	2,411	137	—	2,279	136	—	2,279	117	—	2,382	174	—	53,086	6,011
Subtotal	8,020	8,217	6.3	7,106	7,369	5.5	6,523	6,841	5.1	6,156	6,560	4.8	5,801	6,537	4.8	248,009	250,967
Child (<13 yrs at diagnosis)																	
Perinatal	13	13	—	17	18	—	13	14	—	9	9	—	6	7	—	8,927	8,948
Other ^g	1	1	—	6	6	—	3	3	—	1	1	—	1	1	—	893	895
Subtotal	14	14	0.0	23	24	0.0	16	17	0.0	10	10	0.0	7	8	0.0	9,820	9,843
Region of residence																	
Northeast	6,516	6,750	12.2	5,639	5,923	10.7	5,119	5,449	9.8	4,771	5,173	9.3	4,301	4,872	8.7	348,674	352,167
Midwest	3,652	3,714	5.6	3,369	3,456	5.2	3,249	3,356	5.0	3,130	3,283	4.9	2,944	3,221	4.8	125,360	126,372
South	14,718	15,036	13.3	13,715	14,183	12.3	12,908	13,466	11.6	12,382	13,109	11.2	12,769	14,345	12.1	473,007	477,964
West	5,612	5,761	8.0	4,872	5,054	7.0	4,279	4,484	6.2	4,130	4,408	6.0	3,836	4,251	5.7	235,487	237,536
U.S. dependent areas	639	653	14.9	534	552	13.4	499	529	13.0	428	471	11.6	341	447	11.2	35,335	35,672
Total^h	31,137	31,915	10.2	28,129	29,168	9.3	26,054	27,283	8.6	24,841	26,444	8.3	24,191	27,135	8.5	1,217,863	1,229,711

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited. Rates are not calculated by transmission category because of the lack of denominator data.

^b From the beginning of the epidemic through 2013.

^c Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^d Hispanics/Latinos can be of any race.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^g Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^h Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 3a. Diagnoses of HIV infection, by race/ethnicity and selected characteristics, 2013—United States

	American Indian/ Alaska Native			Asian			Black/African American			Hispanic/Latino ^a			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total				
	Estimated ^b			Estimated ^b			Estimated ^b			Estimated ^b			Estimated ^b			Estimated ^b			Estimated ^b			Estimated ^b				
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^c
Age at diagnosis (yr)																										
<13	0	0	0.0	14	15	0.6	108	122	1.7	11	13	0.1	0	0	0.0	23	26	0.1	8	10	0.5	164	187	0.4		
13–14	1	1	1.4	1	1	0.3	26	30	2.6	4	4	0.2	0	0	0.0	6	7	0.2	1	1	0.4	39	45	0.5		
15–19	3	4	2.2	9	10	1.1	1,107	1,257	41.1	305	340	7.4	1	1	2.6	184	202	1.7	43	49	7.6	1,652	1,863	8.8		
20–24	34	37	18.4	126	145	12.2	4,004	4,611	135.3	1,384	1,575	33.7	7	8	16.2	1,327	1,481	11.6	177	196	36.1	7,059	8,053	35.3		
25–29	33	36	20.8	152	174	13.0	3,220	3,741	131.4	1,659	1,887	42.9	12	13	26.7	1,606	1,793	14.5	162	181	44.1	6,844	7,825	36.3		
30–34	35	39	24.5	144	161	11.7	2,151	2,489	91.4	1,435	1,635	37.4	14	15	32.9	1,502	1,682	13.8	123	144	39.7	5,404	6,165	29		
35–39	17	18	12.6	124	140	10.6	1,681	1,955	79.0	1,112	1,268	31.3	5	5	13.6	1,208	1,360	12.1	99	112	37.7	4,246	4,858	24.8		
40–44	21	22	15.2	125	137	10.3	1,598	1,868	71.6	958	1,090	29.1	5	7	18.4	1,435	1,609	12.7	75	86	31.6	4,217	4,820	23.1		
45–49	20	22	14.2	81	90	7.8	1,723	2,027	76.5	826	946	28.8	8	9	26.5	1,564	1,765	12.9	89	102	42.7	4,311	4,961	23.4		
50–54	19	23	14.3	34	39	3.7	1,370	1,609	59.3	569	652	23.3	7	7	22.4	1,198	1,351	8.7	57	66	27.7	3,254	3,747	16.6		
55–59	5	6	3.9	29	35	3.6	936	1,088	44.8	340	388	17.7	1	1	4.2	799	903	5.9	41	46	22.4	2,151	2,467	11.6		
60–64	4	4	3.6	12	13	1.6	494	580	30.2	147	168	10.4	1	1	4.9	470	523	3.9	22	26	16.6	1,150	1,316	7.3		
≥65	5	6	2.6	11	12	0.7	385	458	11.9	128	150	4.5	0	0	0.0	350	398	1.1	17	21	6.8	896	1,045	2.3		
Transmission category																										
Male adult or adolescent																										
Male-to-male sexual contact	91	120	—	487	703	—	7,902	12,069	—	5,273	7,199	—	33	47	—	7,212	9,853	—	500	699	—	21,498	30,689	—		
Injection drug use	19	22	—	17	26	—	265	903	—	230	480	—	0	0	—	337	473	—	19	37	—	887	1,942	—		
Male-to-male sexual contact and injection drug use	6	10	—	8	12	—	139	331	—	188	271	—	2	3	—	488	611	—	20	32	—	851	1,270	—		
Heterosexual contact ^d	14	17	—	38	56	—	1,254	2,493	—	483	718	—	1	2	—	360	530	—	49	71	—	2,199	3,887	—		
Other ^e	24	0	—	157	2	—	4,092	52	—	1,461	16	—	11	0	—	1,851	28	—	149	0	—	7,745	99	—		
Subtotal	154	169	18.4	707	799	12.5	13,652	15,847	105.7	7,635	8,686	41.8	47	52	24.1	10,248	11,495	13.8	737	839	44.1	33,180	37,887	29.4		

Table 3a. Diagnoses of HIV infection, by race/ethnicity and selected characteristics, 2013—United States (cont)

	American Indian/ Alaska Native			Asian			Black/African American			Hispanic/Latino ^a			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^b			Estimated ^b			Estimated ^b			Estimated ^b			Estimated ^b			Estimated ^b			Estimated ^b					
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^c	Rate
Female adult or adolescent																								
Injection drug use	7	14	—	1	7	—	151	532	—	75	175	—	2	3	—	259	393	—	15	31	—	510	1,154	—
Heterosexual contact ^d	12	34	—	67	150	—	2,182	5,268	—	559	1,232	—	5	13	—	551	1,174	—	70	159	—	3,446	8,031	—
Other ^e	24	0	—	73	2	—	2,710	67	—	598	11	—	7	.	—	591	13	—	84	0	—	4,087	93	—
Subtotal	43	49	5.1	141	159	2.2	5,043	5,867	34.8	1,232	1,419	7.0	14	15	7.3	1,401	1,579	1.8	169	190	9.3	8,043	9,278	6.9
Child (<13 yrs at diagnosis)																								
Perinatal	0	0	—	7	8	—	60	68	—	8	10	—	0	0	—	14	16	—	4	5	—	93	107	—
Other ^f	0	0	—	7	7	—	48	54	—	3	4	—	0	0	—	9	10	—	4	5	—	71	80	—
Subtotal	0	0	0.0	14	15	0.6	108	122	1.7	11	13	0.1	0	0	0.0	23	26	0.1	8	10	0.5	164	187	0.4
Region of residence																								
Northeast	11	11	9.1	165	192	5.7	3,127	3,704	59.2	2,012	2,402	31.8	2	2	10.3	1,897	2,267	6.0	285	329	38.4	7,499	8,908	15.9
Midwest	25	27	6.8	84	92	4.7	2,764	3,016	43.3	614	698	14.0	2	2	7.2	1,968	2,121	4.1	143	153	12.9	5,600	6,109	9
South	56	69	8.9	231	263	7.2	11,616	13,669	60.8	3,634	4,080	20.6	12	15	18.2	5,162	5,814	8.4	355	412	20.5	21,066	24,323	20.5
West	105	111	10.7	382	426	6.0	1,296	1,447	43.0	2,618	2,937	13.5	45	49	12.1	2,645	2,898	7.5	131	145	6.8	7,222	8,013	10.8
Total^g	197	218	9.4	862	973	6.0	18,803	21,836	55.9	8,878	10,117	18.7	61	67	12.7	11,672	13,101	6.6	914	1,039	16.8	41,387	47,352	15

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Hispanics/Latinos can be of any race.

^b Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^c Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 3b. Diagnoses of HIV infection, by race/ethnicity and selected characteristics, 2013—United States and 6 dependent areas

	American Indian/ Alaska Native		Asian		Black/African American		Hispanic/Latino ^a		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^{b,c}
Age at diagnosis (yr)																
<13	0	0	14	15	108	122	11	13	0	0	23	26	8	10	164	187
13–14	1	1	1	1	26	30	5	6	0	0	6	7	1	1	40	46
15–19	3	4	9	10	1,107	1,257	323	362	1	1	184	202	43	49	1,670	1,885
20–24	34	37	126	145	4,006	4,613	1,455	1,664	7	8	1,327	1,481	177	196	7,132	8,144
25–29	33	36	152	174	3,222	3,743	1,724	1,969	12	13	1,606	1,793	162	181	6,911	7,909
30–34	35	39	144	161	2,151	2,489	1,521	1,742	14	15	1,503	1,683	123	144	5,491	6,274
35–39	17	18	124	140	1,682	1,957	1,173	1,344	5	5	1,208	1,360	99	112	4,308	4,936
40–44	21	22	125	137	1,599	1,869	1,024	1,173	5	7	1,435	1,609	75	86	4,284	4,903
45–49	20	22	81	90	1,726	2,032	905	1,046	8	9	1,565	1,767	89	102	4,394	5,066
50–54	19	23	34	39	1,372	1,611	647	750	7	7	1,199	1,353	57	66	3,335	3,850
55–59	5	6	29	35	936	1,088	382	441	1	1	799	903	41	46	2,193	2,519
60–64	4	4	12	13	496	583	175	203	1	1	470	523	22	26	1,180	1,354
≥65	5	6	11	12	385	458	148	176	0	0	350	398	17	21	916	1,070
Transmission category																
Male adult or adolescent																
Male-to-male sexual contact	91	120	487	703	7,904	12,072	5,517	7,527	33	47	7,214	9,857	500	699	21,746	31,023
Injection drug use	19	22	17	26	265	904	301	588	0	0	337	473	19	37	958	2,051
Male-to-male sexual contact and injection drug use	6	10	8	12	139	331	198	285	2	3	488	611	20	32	861	1,284
Heterosexual contact ^d	14	17	38	56	1,255	2,496	580	849	1	2	360	531	49	71	2,297	4,021
Other ^e	24	0	157	2	4,095	52	1,501	16	11	0	1,852	28	149	0	7,789	99
Subtotal	154	169	707	799	13,658	15,855	8,097	9,266	47	52	10,251	11,500	737	839	33,651	38,479

Table 3b. Diagnoses of HIV infection, by race/ethnicity and selected characteristics, 2013—United States and 6 dependent areas (cont)

	American Indian/ Alaska Native		Asian		Black/African American		Hispanic/Latino ^a		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^{b,c}
Female adult or adolescent																
Injection drug use	7	14	1	7	151	533	98	208	2	3	259	393	15	31	533	1,188
Heterosexual contact ^d	12	34	67	150	2,185	5,276	656	1,389	5	13	551	1,174	70	159	3,546	8,195
Other ^e	24	0	73	2	2,714	67	631	13	7	0	591	13	84	0	4,124	95
Subtotal	43	49	141	159	5,050	5,876	1,385	1,610	14	15	1,401	1,579	169	190	8,203	9,479
Child (<13 yrs at diagnosis)																
Perinatal	0	0	7	8	60	68	8	10	0	0	14	16	4	5	93	107
Other ^f	0	0	7	7	48	54	3	4	0	0	9	10	4	5	71	80
Subtotal	0	0	14	15	108	122	11	13	0	0	23	26	8	10	164	187
Region of residence																
Northeast	11	11	165	192	3,127	3,704	2012	2,402	2	2	1,897	2,267	285	329	7,499	8,908
Midwest	25	27	84	92	2,764	3,016	614	698	2	2	1,968	2,121	143	153	5,600	6,109
South	56	69	231	263	11,616	13,669	3634	4,080	12	15	5,162	5,814	355	412	21,066	24,323
West	105	111	382	426	1,296	1,447	2618	2,937	45	49	2,645	2,898	131	145	7,222	8,013
U.S. dependent areas	0	0	0	0	13	17	615	771	0	0	3	4	0	0	631	792
Total^g	197	218	862	973	18,816	21,853	9,493	10,888	61	67	11,675	13,105	914	1,039	42,018	48,145

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Hispanics/Latinos can be of any race.

^b Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^c Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 4a. Stage 3 (AIDS), by race/ethnicity and selected characteristics, 2013—United States

	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total					
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c					
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d	Rate
Age at diagnosis (yr)																											
<13	0	0	0.0	0	0	0.0	3	3	0.0	2	2	0.0	0	0	0.0	0	0	0.0	2	2	0.1	2	2	0.1	7	8	0.0
13–14	0	0	0.0	0	0	0.0	21	26	2.2	1	1	0.1	0	0	0.0	3	4	0.1	0	0	0.0	25	30	0.4			
15–19	0	0	0.0	3	4	0.4	288	332	10.9	62	69	1.5	0	0	0.0	20	22	0.2	8	9	1.4	381	435	2.1			
20–24	5	5	2.7	24	27	2.3	1,307	1,476	43.3	349	386	8.3	2	2	4.4	244	270	2.1	65	72	13.3	1,996	2,239	9.8			
25–29	10	11	6.3	42	46	3.4	1,541	1,740	61.1	632	699	15.9	6	6	13.2	466	518	4.2	90	103	24.9	2,787	3,123	14.5			
30–34	19	20	12.9	57	63	4.5	1,385	1,557	57.2	740	817	18.7	2	2	4.7	633	705	5.8	91	104	28.6	2,927	3,268	15.4			
35–39	8	8	5.9	62	68	5.1	1,280	1,437	58.1	730	808	20.0	4	4	11.1	687	763	6.8	98	110	37.0	2,869	3,200	16.3			
40–44	13	14	9.6	73	82	6.1	1,367	1,541	59.1	673	748	20.0	4	5	12.7	887	988	7.8	104	118	43.3	3,121	3,496	16.8			
45–49	14	15	9.9	53	59	5.2	1,540	1,735	65.5	659	730	22.2	9	10	28.2	1,021	1,129	8.2	90	103	43.1	3,386	3,781	17.8			
50–54	15	17	10.2	28	31	2.9	1,260	1,422	52.4	461	510	18.2	6	6	19.3	940	1,044	6.7	93	106	44.5	2,803	3,135	13.9			
55–59	4	4	2.9	16	18	1.9	861	969	39.8	246	273	12.4	0	0	0.0	601	665	4.4	61	69	33.8	1,789	1,998	9.4			
60–64	3	3	2.7	9	10	1.2	490	555	28.9	137	155	9.5	0	0	0.0	349	386	2.9	31	36	23.1	1,019	1,144	6.3			
≥65	5	5	2.4	7	8	0.4	335	379	9.9	122	138	4.1	1	1	2.7	239	265	0.8	31	35	11.5	740	831	1.9			
Transmission category																											
Male adult or adolescent																											
Male-to-male sexual contact	40	53	—	200	300	—	3,990	5,804	—	2,439	3,310	—	15	25	—	3,526	4,664	—	332	455	—	10,542	14,611	—			
Injection drug use	6	7	—	5	11	—	401	816	—	224	386	—	0	0	—	265	340	—	31	49	—	932	1,610	—			
Male-to-male sexual contact and injection drug use	5	8	—	4	6	—	189	338	—	138	194	—	1	2	—	338	420	—	41	58	—	716	1,026	—			
Heterosexual contact ^e	9	11	—	24	36	—	1,085	1,892	—	372	529	—	1	2	—	235	330	—	48	66	—	1,774	2,865	—			
Other ^f	13	0	—	87	1	—	2,255	81	—	839	24	—	10	0	—	854	34	—	106	4	—	4,164	144	—			
Subtotal	73	79	8.6	320	354	5.5	7,920	8,930	59.6	4,012	4,443	21.4	27	29	13.6	5,218	5,788	6.9	558	633	33.3	18,128	20,256	15.7			

Table 4a. Stage 3 (AIDS), by race/ethnicity and selected characteristics, 2013—United States (cont)

	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c					
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d	Rate
Female adult or adolescent																								
Injection drug use	5	8	—	1	4	—	281	597	—	95	165	—	0	0	—	230	314	—	33	55	—	645	1,143	—
Heterosexual contact ^e	9	17	—	28	54	—	1,800	3,514	—	388	707	—	4	7	—	372	641	—	95	169	—	2,696	5,109	—
Other ^f	9	0	—	25	2	—	1,674	128	—	317	19	—	3	0	—	270	16	—	76	7	—	2,374	172	—
Subtotal	23	25	2.6	54	60	0.8	3,755	4,238	25.1	800	891	4.4	7	7	3.5	872	971	1.1	204	231	11.3	5,715	6,424	4.8
Child (<13 yrs at diagnosis)																								
Perinatal	0	0	—	0	0	—	3	3	—	2	2	—	0	0	—	—	0	—	1	1	—	6	7	—
Other ^g	0	0	—	0	0	—	—	0	—	—	0	—	0	0	—	—	0	—	1	1	—	1	1	—
Subtotal	0	0	0.0	0	0	0.0	3	3	0.0	2	2	0.0	0	0	0.0	—	0	0.0	2	2	0.1	7	8	0.0
Region of residence																								
Northeast	4	4	3.4	73	82	2.4	1,980	2,251	36.0	1,151	1,304	17.3	0	0	0	895	1,008	2.7	198	223	26.1	4,301	4,872	8.7
Midwest	14	15	3.8	40	43	2.2	1,463	1,606	23.1	320	351	7.0	0	0	0	1,009	1,098	2.1	98	108	9.1	2,944	3,221	4.8
South	20	22	2.9	83	93	2.5	7,555	8,557	38.0	1,980	2,178	11.0	8	9	10.9	2,757	3,068	4.4	366	418	20.8	12,769	14,345	12.1
West	58	62	6.0	178	197	2.8	680	758	22.5	1,363	1,503	6.9	26	28	6.9	1,429	1,585	4.1	102	117	5.5	3,836	4,251	5.7
Total^h	96	104	4.5	374	415	2.6	11,678	13,172	33.7	4,814	5,336	9.9	34	37	6.9	6,090	6,759	3.4	764	867	14.0	23,850	26,688	8.4

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^g Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^h Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 4b. Stage 3 (AIDS), by race/ethnicity and selected characteristics, 2013—United States and 6 dependent areas

	American Indian/ Alaska Native		Asian ^a		Black/African American		Hispanic/Latino ^b		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Age at diagnosis (yr)																
<13	0	0	0	0	3	3	2	2	0	0	0	0	2	2	7	8
13–14	0	0	0	0	21	26	2	2	0	0	3	4	0	0	26	31
15–19	0	0	3	4	288	332	65	73	0	0	20	22	8	9	384	439
20–24	5	5	24	27	1,307	1,476	369	413	2	2	244	270	65	72	2,016	2,266
25–29	10	11	42	46	1,541	1,740	655	728	6	6	466	518	90	103	2,810	3,152
30–34	19	20	57	63	1,385	1,557	773	860	2	2	633	705	91	104	2,960	3,311
35–39	8	8	62	68	1,280	1,437	764	853	4	4	687	763	98	110	2,903	3,244
40–44	13	14	73	82	1,368	1,543	715	802	4	5	887	988	104	118	3,164	3,551
45–49	14	15	53	59	1,541	1,736	711	799	9	10	1,022	1,131	90	103	3,440	3,853
50–54	15	17	28	31	1,261	1,423	518	585	6	6	941	1,045	93	106	2,862	3,213
55–59	4	4	16	18	862	970	280	319	0	0	601	665	61	69	1,824	2,045
60–64	3	3	9	10	491	556	153	176	0	0	349	386	31	36	1,036	1,167
≥65	5	5	7	8	335	379	141	161	1	1	239	265	31	35	759	854
Transmission category																
Male adult or adolescent																
Male-to-male sexual contact	40	53	200	300	3,990	5,804	2,532	3,435	15	25	3,527	4,666	332	455	10,636	14,739
Injection drug use	6	7	5	11	401	816	284	469	0	0	265	341	31	49	992	1,694
Male-to-male sexual contact and injection drug use	5	8	4	6	189	338	147	205	1	2	338	420	41	58	725	1,036
Heterosexual contact ^e	9	11	24	36	1,086	1,894	450	636	1	2	235	330	48	66	1,853	2,974
Other ^f	13	0	87	1	2,256	81	850	26	10	0	855	34	106	4	4,177	146
Subtotal	73	79	320	354	7,922	8,933	4,263	4,771	27	29	5,220	5,791	558	633	18,383	20,590

Table 4b. Stage 3 (AIDS), by race/ethnicity and selected characteristics, 2013—United States and 6 dependent areas (cont)

	American Indian/ Alaska Native		Asian ^a		Black/African American		Hispanic/Latino ^b		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Female adult or adolescent																
Injection drug use	5	8	1	4	281	597	111	187	0	0	230	314	33	55	661	1,165
Heterosexual contact ^e	9	17	28	54	1,802	3,518	448	792	4	7	372	641	95	169	2,758	5,198
Other ^f	9	.	25	2	1,675	128	324	22	3	0	270	16	76	7	2,382	174
Subtotal	23	25	54	60	3,758	4,242	883	1,001	7	7	872	971	204	231	5,801	6,537
Child (<13 yrs at diagnosis)																
Perinatal	0	0	0	0	3	3	2	2	0	0	0	0	1	1	6	7
Other ^g	0	0	0	0	0	0	0	0	0	0	0	0	1	1	1	1
Subtotal	0	0	0	0	3	3	2	2	0	0	0	0	2	2	7	8
Region of residence																
Northeast	4	4	73	82	1,980	2,251	1,151	1,304	0	0	895	1,008	198	223	4,301	4,872
Midwest	14	15	40	43	1,463	1,606	320	351	0	0	1,009	1,098	98	108	2,944	3,221
South	20	22	83	93	7,555	8,557	1,980	2,178	8	9	2,757	3,068	366	418	12,769	14,345
West	58	62	178	197	680	758	1,363	1,503	26	28	1,429	1,585	102	117	3,836	4,251
U.S. dependent areas	0	0	0	0	5	7	334	437	0	0	2	3	0	0	341	447
Total^h	96	104	374	415	11,683	13,179	5,148	5,773	34	37	6,092	6,762	764	867	24,191	27,135

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^g Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^h Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 5a. Diagnoses of HIV infection among children aged <13 years, by race/ethnicity, 2009–2013—United States

Race/ethnicity	2009			2010			2011			2012			2013		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		No.	Rate
American Indian/Alaska Native	0	0	0.0	0	0	0.0	1	1	0.2	0	0	0.0	0	0	0.0
Asian	8	8	0.3	2	2	0.1	6	6	0.3	12	13	0.5	14	15	0.6
Black/African American	152	156	2.1	148	152	2.1	118	124	1.7	164	177	2.4	108	122	1.7
Hispanic/Latino ^b	32	33	0.3	33	34	0.3	23	24	0.2	15	17	0.1	11	13	0.1
Native Hawaiian/Other Pacific Islander	0	0	0.0	0	0	0.0	0	0	0.0	1	1	1.0	0	0	0.0
White	17	17	0.1	31	32	0.1	30	31	0.1	31	33	0.1	23	26	0.1
Multiple races	9	9	0.6	11	11	0.5	9	9	0.4	9	10	0.4	8	10	0.5
Total^c	218	223	0.4	225	232	0.4	187	196	0.4	232	250	0.5	164	187	0.4

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^b Hispanics/Latinos can be of any race.

^c Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 5b. Diagnoses of HIV infection among children aged <13 years, by race/ethnicity, 2009–2013—United States and 6 dependent areas

Race/ethnicity	2009		2010		2011		2012		2013	
	No.	Est. No. ^a	No.	Est. No. ^a	No.	Est. No. ^a	No.	Est. No. ^a	No.	Est. No. ^a
American Indian/Alaska Native	0	0	0	0	1	1	0	0	0	0
Asian	8	8	2	2	6	6	12	13	14	15
Black/African American	152	156	148	152	118	124	164	177	108	122
Hispanic/Latino ^b	33	34	37	39	25	26	15	17	11	13
Native Hawaiian/Other Pacific Islander	0	0	0	0	1	1	1	1	0	0
White	17	17	31	32	30	31	31	33	23	26
Multiple races	9	9	11	11	9	9	9	10	8	10
Total^c	219	224	229	236	190	199	232	250	164	187

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting.

^b Hispanics/Latinos can be of any race.

^c Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 6a. Stage 3 (AIDS) among children aged <13 years, by race/ethnicity, 2009–2013 and cumulative—United States

Race/ethnicity	2009			2010			2011			2012			2013			Cumulative ^b	
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Est. No. ^a
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		No.	Rate		
American Indian/Alaska Native	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0	26	26
Asian ^c	1	1	0.0	1	1	0.0	0	0	0.0	0	0	0.0	0	0	0.0	46	46
Black/African American	8	8	0.1	15	15	0.2	11	12	0.2	8	8	0.1	3	3	0.0	5,601	5,615
Hispanic/Latino ^d	4	4	0.0	4	4	0.0	2	2	0.0	1	1	0.0	2	2	0.0	1,955	1,960
Native Hawaiian/Other Pacific Islander	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0	6	6
White	1	1	0.0	3	3	0.0	2	2	0.0	1	1	0.0	0	0	0.0	1,540	1,542
Multiple races	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0	2	2	0.1	225	226
Total^e	14	14	0.0	23	24	0.0	15	16	0.0	10	10	0.0	7	8	0.0	9,399	9,421

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^b From the beginning of the epidemic through 2013.

^c Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^d Hispanics/Latinos can be of any race.

^e Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 6b. Stage 3 (AIDS) among children aged <13 years, by race/ethnicity, 2009–2013 and cumulative—United States and 6 dependent areas

Race/ethnicity	2009		2010		2011		2012		2013		Cumulative ^a	
	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b	No.	Est. No. ^b
American Indian/Alaska Native	0	0	0	0	0	0	0	0	0	0	26	26
Asian ^c	1	1	1	1	0	0	0	0	0	0	47	47
Black/African American	8	8	15	15	11	12	8	8	3	3	5,611	5,625
Hispanic/Latino ^d	4	4	4	4	2	2	1	1	2	2	2,364	2,370
Native Hawaiian/Other Pacific Islander	0	0	0	0	1	1	0	0	0	0	7	7
White	1	1	3	3	2	2	1	1	0	0	1,540	1,542
Multiple races	0	0	0	0	0	0	0	0	2	2	225	226
Total^e	14	14	23	24	16	17	10	10	7	8	9,820	9,843

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a From the beginning of the epidemic through 2013.

^b Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting.

^c Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^d Hispanics/Latinos can be of any race.

^e Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 7. Stage 3 (AIDS) among children aged <13 years, by year of diagnosis, 1992–2013—United States and 6 dependent areas

Year of diagnosis	United States		United States and dependent areas	
	No.	Est. No. ^a	No.	Est. No. ^a
1992	902	903	950	951
1993	884	886	922	924
1994	811	813	836	838
1995	680	682	703	705
1996	521	522	535	536
1997	342	343	354	355
1998	247	248	254	255
1999	198	199	203	204
2000	133	134	135	136
2001	130	131	131	132
2002	108	109	115	116
2003	75	75	77	77
2004	59	59	60	60
2005	53	53	53	53
2006	40	41	43	44
2007	34	34	34	34
2008	39	39	40	40
2009	14	14	14	14
2010	23	24	23	24
2011	15	16	16	17
2012	10	10	10	10
2013	7	8	7	8

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting.

Table 8. Diagnoses of HIV infection among adult and adolescent Hispanics/Latinos, by transmission category and place of birth, 2013—United States and 6 dependent areas

Transmission category	Central America			Cuba			Mexico			Puerto Rico			South America			United States			Total		
	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No. ^b	Est. No. ^{a,b}	%
Male adult or adolescent																					
Male-to-male sexual contact	273	410	75.9	277	304	86.9	814	1,119	82.7	360	476	59.9	301	365	88.0	2,515	3,221	84.1	5,517	7,527	81.2
Injection drug use	5	20	3.6	2	3	0.9	29	52	3.8	103	148	18.6	3	10	2.3	118	216	5.6	301	588	6.3
Male-to-male sexual contact and injection drug use	9	15	2.7	3	3	1.0	21	35	2.6	17	23	2.9	4	6	1.4	112	148	3.9	198	285	3.1
Heterosexual contact ^c	64	96	17.8	33	39	11.2	81	144	10.6	104	147	18.5	23	34	8.3	183	233	6.1	580	849	9.2
Other ^d	125	0	0.0	18	0	0.0	259	3	0.2	80	1	0.1	40	0	0.0	427	10	0.3	1,501	16	0.2
Subtotal	476	541	100	333	350	100	1,204	1,353	100	664	795	100	371	414	100	3,355	3,828	100	8,097	9,266	100
Female adult or adolescent																					
Injection drug use	0	5	3.9	1	1	4.5	3	8	5.8	29	41	15.8	2	4	7.3	48	98	17.7	98	208	12.9
Heterosexual contact ^c	52	124	95.9	20	27	91.2	63	124	92.9	128	220	83.9	30	52	90.3	244	454	81.5	656	1,389	86.3
Other ^d	60	0	0.2	7	1	4.2	51	2	1.3	62	1	0.3	20	1	2.4	190	5	0.8	631	13	0.8
Subtotal	112	129	100	28	29	100	117	133	100	219	262	100	52	58	100	482	557	100	1,385	1,610	100
Total^e	588	670	100	361	379	100	1,321	1,486	100	883	1,058	100	423	472	100	3,837	4,385	100	9,482	10,875	100

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^b Entries include persons whose place of birth is not among those listed and persons whose place of birth is unknown.

^c Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^d Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^e Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 9. Stage 3 (AIDS) among adult and adolescent Hispanics/Latinos, by transmission category and place of birth, 2013—United States and 6 dependent areas

Transmission category	Central America			Cuba			Mexico			Puerto Rico			South America			United States			Total		
	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No.	Est. No. ^a	%	No. ^b	Est. No. ^{a,b}	%
Male adult or adolescent																					
Male-to-male sexual contact	186	264	72.3	101	114	73.6	471	648	78.0	149	196	40.7	137	164	82.0	1,061	1,360	74.1	2,532	3,435	72.0
Injection drug use	5	16	4.3	4	6	3.8	20	37	4.5	104	138	28.6	4	7	3.7	116	179	9.8	284	469	9.8
Male-to-male sexual contact and injection drug use	6	10	2.7	3	3	2.2	17	28	3.3	18	22	4.7	3	5	2.4	84	108	5.9	147	205	4.3
Heterosexual contact ^c	52	76	20.7	26	31	20.3	69	116	14.0	90	122	25.3	16	23	11.3	136	171	9.3	450	636	13.3
Other ^d	83	0	0.1	11	0	0.1	182	2	0.2	32	4	0.8	23	1	0.6	251	18	1.0	850	26	0.5
Subtotal	332	366	100	145	155	100	759	831	100	393	483	100	183	201	100	1,648	1,835	100	4,263	4,771	100
Female adult or adolescent																					
Injection drug use	3	7	8.0	1	1	15.0	5	9	11.0	33	43	23.6	0	1	2.8	59	98	23.3	111	187	18.7
Heterosexual contact ^c	36	79	90.6	5	7	85.0	42	71	87.3	89	137	75.0	16	28	96.4	194	310	73.6	448	792	79.2
Other ^d	39	1	1.4	2	0	0.0	28	1	1.7	27	3	1.4	10	0	0.7	126	13	3.1	324	22	2.2
Subtotal	78	88	100	8	9	100	75	81	100	149	183	100	26	29	100	379	421	100	883	1,001	100
Total^e	410	453	100	153	164	100	834	913	100	542	665	100	209	229	100	2,027	2,256	100	5,146	5,771	100

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^b Entries include persons whose place of birth is not among those listed and persons whose place of birth is unknown.

^c Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^d Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^e Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 10a. Deaths of persons with diagnosed HIV infection, by year of death and selected characteristics, 2009–2012—United States

	2009			2010			2011			2012		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
Age at death (yr)												
<13	6	6	0.0	5	6	0.0	0	0	0.0	5	6	0.0
13–14	3	3	0.0	3	3	0.0	1	1	0.0	2	2	0.0
15–19	34	35	0.2	38	40	0.2	25	27	0.1	21	25	0.1
20–24	189	197	0.9	190	202	0.9	194	211	1.0	156	188	0.8
25–29	502	523	2.4	414	440	2.1	409	446	2.1	386	460	2.2
30–34	850	885	4.5	746	789	3.9	674	728	3.5	640	767	3.7
35–39	1,503	1,561	7.6	1,184	1,250	6.2	1,029	1,105	5.6	854	1,005	5.2
40–44	2,572	2,676	12.7	2,087	2,205	10.6	1,800	1,940	9.2	1,438	1,691	8.0
45–49	3,439	3,576	15.7	3,186	3,375	14.9	2,899	3,122	14.1	2,384	2,799	12.9
50–54	3,326	3,462	15.9	3,161	3,342	15.0	3,098	3,340	14.8	2,832	3,325	14.7
55–59	2,588	2,695	14.2	2,551	2,700	13.6	2,610	2,811	13.9	2,401	2,814	13.5
60–64	1,566	1,625	10.3	1,679	1,776	10.5	1,693	1,826	10.2	1,689	1,998	11.2
≥65	1,741	1,814	4.6	1,671	1,767	4.4	1,798	1,939	4.7	1,781	2,092	4.8
Race/ethnicity												
American Indian/Alaska Native	66	68	2.9	75	79	3.5	56	59	2.6	53	63	2.7
Asian ^b	78	81	0.6	72	75	0.5	87	93	0.6	64	73	0.5
Black/African American	9,203	9,599	25.5	8,182	8,686	22.9	7,798	8,444	22.0	6,880	8,171	21.1
Hispanic/Latino ^c	2,832	2,916	6.0	2,695	2,812	5.5	2,634	2,799	5.4	2,259	2,586	4.9
Native Hawaiian/Other Pacific Islander	7	7	1.6	7	7	1.4	10	11	2.1	3	4	0.7
White	5,326	5,545	2.8	5,090	5,395	2.7	4,920	5,307	2.7	4,604	5,426	2.7
Multiple races	807	842	18.5	794	842	14.9	725	783	13.4	726	849	14.1
Transmission category												
Male adult or adolescent												
Male-to-male sexual contact	5,597	7,110	—	5,360	6,854	—	5,235	6,857	—	4,788	6,777	—
Injection drug use	2,816	3,417	—	2,546	3,133	—	2,265	2,864	—	1,991	2,723	—
Male-to-male sexual contact and injection drug use	1,335	1,514	—	1,176	1,355	—	1,147	1,343	—	1,030	1,303	—
Heterosexual contact ^d	1,310	1,871	—	1,272	1,787	—	1,235	1,781	—	1,114	1,719	—
Perinatal	19	20	—	28	30	—	34	37	—	38	44	—
Other ^e	2,407	100	—	2,148	92	—	2,119	94	—	1,802	79	—
Subtotal	13,484	14,032	11.3	12,530	13,250	10.6	12,035	12,977	10.3	10,763	12,645	9.9

Table 10a. Deaths of persons with diagnosed HIV infection, by year of death and selected characteristics, 2009–2012—United States (cont)

	2009			2010			2011			2012		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
Female adult or adolescent												
Injection drug use	1,523	1,914	—	1,315	1,684	—	1,304	1,669	—	1,133	1,611	—
Heterosexual contact ^d	2,092	3,030	—	1,956	2,895	—	1,823	2,767	—	1,705	2,832	—
Perinatal	44	45	—	30	32	—	44	47	—	39	45	—
Other ^e	1,170	31	—	1,079	29	—	1,024	36	—	944	34	—
Subtotal	4,829	5,020	3.9	4,380	4,640	3.5	4,195	4,519	3.4	3,821	4,521	3.4
Child (<13 yrs at death)												
Perinatal	5	5	—	4	4	—	—	—	—	4	5	—
Other ^e	1	1	—	1	1	—	—	—	—	1	1	—
Subtotal	6	6	0.0	5	6	0.0	0	0	0.0	5	6	0.0
Region of residence												
Northeast	4,800	4,994	9.0	4,603	4,874	8.8	4,291	4,616	8.3	3,752	4,362	7.8
Midwest	1,972	2,042	3.1	1,917	2,016	3.0	1,845	1,979	2.9	1,620	1,907	2.8
South	8,908	9,307	8.2	7,901	8,401	7.3	7,616	8,263	7.1	6,886	8,218	7.0
West	2,639	2,715	3.8	2,494	2,605	3.6	2,478	2,638	3.6	2,331	2,686	3.7
Total^f	18,319	19,058	6.2	16,915	17,896	5.8	16,230	17,496	5.6	14,589	17,173	5.5

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Deaths of persons with diagnosed HIV infection may be due to any cause.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 10b. Deaths of persons with diagnosed HIV infection, by year of death and selected characteristics, 2009–2012—United States and 6 dependent areas

	2009			2010			2011			2012		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Age at death (yr)												
<13	6	6	0.0	5	6	0.0	0	0	0.0	5	6	0.0
13–14	3	3	0.0	3	3	0.0	1	1	0.0	2	2	0.0
15–19	38	39	0.2	40	42	0.2	26	28	0.1	22	27	0.1
20–24	199	207	0.9	193	205	0.9	198	215	1.0	159	192	0.8
25–29	523	546	2.5	424	452	2.1	423	463	2.1	395	472	2.2
30–34	882	920	4.6	785	833	4.1	696	753	3.6	666	800	3.8
35–39	1,568	1,633	7.8	1,236	1,309	6.4	1,072	1,155	5.8	891	1,053	5.3
40–44	2,680	2,795	13.1	2,149	2,276	10.7	1,870	2,021	9.5	1,479	1,743	8.2
45–49	3,554	3,703	16.0	3,294	3,498	15.3	2,997	3,237	14.4	2,460	2,896	13.2
50–54	3,435	3,582	16.3	3,244	3,436	15.2	3,196	3,455	15.1	2,892	3,402	14.9
55–59	2,662	2,777	14.4	2,613	2,771	13.8	2,674	2,886	14.1	2,448	2,875	13.7
60–64	1,608	1,672	10.4	1,726	1,830	10.6	1,738	1,878	10.4	1,721	2,040	11.3
≥65	1,805	1,885	4.7	1,729	1,833	4.5	1,857	2,008	4.8	1,828	2,154	4.9
Race/ethnicity												
American Indian/Alaska Native	66	68	—	75	79	—	56	59	—	53	63	—
Asian ^b	80	83	—	72	75	—	89	95	—	65	74	—
Black/African American	9,217	9,614	—	8,188	8,692	—	7,805	8,452	—	6,888	8,181	—
Hispanic/Latino ^c	3,456	3,604	—	3,211	3,398	—	3,140	3,389	—	2,624	3,057	—
Native Hawaiian/Other Pacific Islander	7	7	—	7	7	—	11	12	—	3	4	—
White	5,329	5,549	—	5,093	5,399	—	4,921	5,308	—	4,606	5,428	—
Multiple races	808	843	—	795	843	—	726	784	—	729	853	—
Transmission category												
Male adult or adolescent												
Male-to-male sexual contact	5,678	7,205	—	5,424	6,930	—	5,316	6,957	—	4,833	6,839	—
Injection drug use	3,047	3,683	—	2,730	3,351	—	2,444	3,081	—	2,116	2,890	—
Male-to-male sexual contact and injection drug use	1,375	1,559	—	1,212	1,397	—	1,177	1,379	—	1,053	1,333	—
Heterosexual contact ^d	1,395	1,971	—	1,351	1,882	—	1,301	1,863	—	1,181	1,812	—
Perinatal	23	24	—	28	30	—	36	39	—	39	46	—
Other ^e	2,429	100	—	2,167	93	—	2,136	94	—	1,816	80	—
Subtotal	13,947	14,542	11.6	12,912	13,683	10.8	12,410	13,414	10.5	11,038	13,000	10.1

Table 10b. Deaths of persons with diagnosed HIV infection, by year of death and selected characteristics, 2009–2012—United States and 6 dependent areas (cont)

	2009			2010			2011			2012		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Female adult or adolescent												
Injection drug use	1,565	1,962	—	1,361	1,738	—	1,352	1,726	—	1,160	1,647	—
Heterosexual contact ^d	2,220	3,176	—	2,048	3,003	—	1,910	2,872	—	1,775	2,928	—
Perinatal	48	50	—	32	34	—	47	51	—	40	46	—
Other ^e	1,177	32	—	1,083	29	—	1,029	37	—	950	34	—
Subtotal	5,010	5,219	4.0	4,524	4,804	3.6	4,338	4,686	3.5	3,925	4,655	3.4
Child (<13 yrs at death)												
Perinatal	5	5	—	4	4	—	0	0	—	4	5	—
Other ^e	1	1	—	1	1	—	0	0	—	1	1	—
Subtotal	6	6	0.0	5	6	0.0	0	0	0.0	5	6	0.0
Region of residence												
Northeast	4,800	4,994	9.0	4,603	4,874	8.8	4,291	4,616	8.3	3,752	4,362	7.8
Midwest	1,972	2,042	3.1	1,917	2,016	3.0	1,845	1,979	2.9	1,620	1,907	2.8
South	8,908	9,307	8.2	7,901	8,401	7.3	7,616	8,263	7.1	6,886	8,218	7.0
West	2,639	2,715	3.8	2,494	2,605	3.6	2,478	2,638	3.6	2,331	2,686	3.7
U.S. dependent areas	644	710	16.2	526	598	14.5	518	604	14.8	379	488	12.1
Total^f	18,963	19,768	6.3	17,441	18,493	5.9	16,748	18,100	5.7	14,968	17,661	5.6

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Deaths of persons with diagnosed HIV infection may be due to any cause.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 11a. Deaths of persons with diagnosed HIV infection ever classified as stage 3 (AIDS), by year of death and selected characteristics, 2009–2012 and cumulative—United States

	2009			2010			2011			2012			Cumulative ^b		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Est. No. ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Age at death (yr)															
<13	3	3	0.0	0	0	0.0	0	0	0.0	1	1	0.0	4,943	4,951	
13–14	2	2	0.0	2	2	0.0	1	1	0.0	1	1	0.0	293	295	
15–19	28	28	0.1	27	28	0.1	20	21	0.1	14	16	0.1	1,293	1,302	
20–24	142	145	0.7	145	151	0.7	145	154	0.7	121	139	0.6	9,732	9,796	
25–29	404	416	1.9	311	325	1.5	307	329	1.5	298	340	1.6	46,742	46,950	
30–34	707	730	3.7	607	634	3.2	547	581	2.8	514	594	2.8	101,463	101,897	
35–39	1,298	1,339	6.5	991	1,034	5.1	873	925	4.7	706	804	4.1	127,398	128,057	
40–44	2,168	2,238	10.7	1,770	1,851	8.9	1,506	1,600	7.6	1,212	1,382	6.6	120,682	121,596	
45–49	2,925	3,019	13.2	2,671	2,799	12.4	2,448	2,602	11.7	2,048	2,342	10.8	92,972	94,089	
50–54	2,748	2,835	13.0	2,599	2,713	12.1	2,589	2,753	12.2	2,362	2,688	11.9	62,731	63,752	
55–59	2,113	2,180	11.5	2,057	2,148	10.8	2,132	2,260	11.2	2,003	2,272	10.9	38,733	39,477	
60–64	1,246	1,278	8.1	1,367	1,427	8.4	1,372	1,454	8.2	1,358	1,548	8.7	22,554	23,020	
≥65	1,314	1,351	3.4	1,304	1,356	3.4	1,385	1,463	3.5	1,404	1,585	3.7	22,873	23,323	
Race/ethnicity															
American Indian/Alaska Native	51	52	2.2	59	61	2.7	49	51	2.2	39	43	1.9	1,852	1,867	
Asian ^c	62	64	0.5	57	59	0.4	71	75	0.5	55	60	0.4	3,420	3,441	
Black/African American	7,536	7,791	20.7	6,702	7,030	18.5	6,362	6,787	17.7	5,676	6,540	16.9	267,545	270,726	
Hispanic/Latino ^d	2,453	2,505	5.2	2,301	2,370	4.7	2,232	2,333	4.5	1,942	2,155	4.1	100,169	100,888	
Native Hawaiian/Other Pacific Islander	6	6	1.3	7	7	1.4	9	9	1.9	2	2	0.4	354	356	
White	4,281	4,412	2.2	4,044	4,225	2.1	3,964	4,205	2.1	3,702	4,199	2.1	267,750	269,653	
Multiple races	709	735	16.1	681	715	12.7	638	683	11.7	626	713	11.9	11,319	11,577	
Transmission category															
Male adult or adolescent															
Male-to-male sexual contact	4,743	5,804	—	4,450	5,503	—	4,414	5,509	—	4,021	5,380	—	287,753	311,087	
Injection drug use	2,362	2,802	—	2,106	2,531	—	1,877	2,306	—	1,698	2,215	—	117,934	130,406	
Male-to-male sexual contact and injection drug use	1,181	1,317	—	1,030	1,164	—	988	1,131	—	891	1,088	—	46,795	50,001	
Heterosexual contact ^e	1,080	1,493	—	1,070	1,453	—	1,019	1,410	—	929	1,359	—	27,498	35,230	
Perinatal	19	20	—	25	26	—	31	33	—	36	42	—	363	376	
Other ^f	1,777	78	—	1,618	78	—	1,533	80	—	1,350	66	—	50,816	8,658	
Subtotal	11,162	11,514	9.3	10,299	10,754	8.6	9,862	10,469	8.3	8,925	10,150	8.0	531,159	535,758	

Table 11a. Deaths of persons with diagnosed HIV infection ever classified as stage 3 (AIDS), by year of death and selected characteristics, 2009–2012 and cumulative—United States (cont)

	2009			2010			2011			2012			Cumulative ^b	
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Est. No. ^a
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		
Female adult or adolescent														
Injection drug use	1,286	1,578	—	1,100	1,371	—	1,106	1,375	—	954	1,299	—	49,370	56,322
Heterosexual contact ^e	1,751	2,402	—	1,630	2,288	—	1,544	2,225	—	1,416	2,191	—	45,702	57,383
Perinatal	40	41	—	28	30	—	42	45	—	39	45	—	474	490
Other ^f	856	27	—	794	25	—	771	28	—	707	25	—	20,761	3,603
Subtotal	3,933	4,048	3.1	3,552	3,713	2.8	3,463	3,674	2.8	3,116	3,561	2.7	116,307	117,797
Child (<13 yrs at death)														
Perinatal	3	3	—	0	0	—	0	0	—	1	1	—	4,483	4,491
Other ^f	0	0	—	0	0	—	0	0	—	0	0	—	460	461
Subtotal	3	3	0.0	0	0	0.0	0	0	0.0	1	1	0.0	4,943	4,951
Region of residence														
Northeast	4,013	4,141	7.5	3,755	3,927	7.1	3,526	3,737	6.7	3,106	3,482	6.2	206,850	208,570
Midwest	1,594	1,631	2.4	1,533	1,585	2.4	1,460	1,535	2.3	1,300	1,474	2.2	66,318	66,856
South	7,235	7,494	6.6	6,480	6,810	5.9	6,252	6,688	5.8	5,656	6,550	5.6	249,947	253,080
West	2,256	2,299	3.2	2,083	2,145	3.0	2,087	2,184	3.0	1,980	2,206	3.0	129,294	130,001
Total^g	15,098	15,565	5.1	13,851	14,467	4.7	13,325	14,143	4.5	12,042	13,712	4.4	652,409	658,507

Note. Deaths of persons with diagnosed HIV infection may be due to any cause.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b From the beginning of the epidemic through 2012.

^c Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^d Hispanics/Latinos can be of any race.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 11b. Deaths of persons with diagnosed HIV infection ever classified as stage 3 (AIDS), by year of death and selected characteristics, 2009–2012 and cumulative—United States and 6 dependent areas

	2009			2010			2011			2012			Cumulative ^b		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Est. No. ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Age at death (yr)															
<13	3	3	0.0	0	0	0.0	0	0	0.0	1	1	0.0	5,191	5,202	
13–14	2	2	0.0	2	2	0.0	1	1	0.0	1	1	0.0	302	304	
15–19	30	31	0.1	29	30	0.1	21	22	0.1	15	17	0.1	1,358	1,370	
20–24	150	154	0.7	146	152	0.7	147	156	0.7	124	143	0.6	10,225	10,296	
25–29	420	434	2.0	318	333	1.6	319	343	1.6	303	347	1.6	48,854	49,088	
30–34	731	756	3.7	632	662	3.3	559	595	2.9	531	615	2.9	105,423	105,911	
35–39	1,352	1,399	6.7	1,028	1,076	5.3	901	958	4.8	729	833	4.2	132,279	133,024	
40–44	2,247	2,326	10.9	1,823	1,911	9.0	1,560	1,663	7.8	1,245	1,424	6.7	124,986	126,005	
45–49	3,013	3,116	13.5	2,771	2,914	12.7	2,537	2,707	12.1	2,104	2,412	11.0	96,079	97,308	
50–54	2,835	2,932	13.3	2,667	2,791	12.3	2,665	2,842	12.4	2,410	2,749	12.0	64,723	65,833	
55–59	2,175	2,248	11.7	2,104	2,201	11.0	2,183	2,320	11.3	2,036	2,313	11.0	39,838	40,634	
60–64	1,279	1,315	8.2	1,406	1,472	8.5	1,405	1,493	8.3	1,382	1,579	8.7	23,285	23,787	
≥65	1,365	1,407	3.5	1,347	1,405	3.4	1,431	1,517	3.6	1,436	1,625	3.7	23,818	24,314	
Race/ethnicity															
American Indian/Alaska Native	51	52	—	59	61	—	49	51	—	39	43	—	1,852	1,867	
Asian ^c	63	65	—	57	59	—	73	77	—	56	61	—	3,455	3,477	
Black/African American	7,546	7,802	—	6,706	7,035	—	6,368	6,794	—	5,683	6,549	—	267,815	271,004	
Hispanic/Latino ^d	2,943	3,048	—	2,719	2,847	—	2,626	2,796	—	2,205	2,486	—	123,727	125,051	
Native Hawaiian/Other Pacific Islander	6	6	—	7	7	—	10	11	—	2	2	—	359	361	
White	4,284	4,416	—	4,044	4,225	—	3,964	4,205	—	3,703	4,200	—	267,828	269,732	
Multiple races	709	735	—	681	715	—	639	684	—	629	717	—	11,325	11,583	
Transmission category															
Male adult or adolescent															
Male-to-male sexual contact	4,811	5,881	—	4,503	5,564	—	4,481	5,591	—	4,054	5,424	—	291,451	314,907	
Injection drug use	2,539	3,001	—	2,260	2,709	—	2,013	2,470	—	1,789	2,332	—	128,099	140,940	
Male-to-male sexual contact and injection drug use	1,216	1,357	—	1,061	1,200	—	1,016	1,165	—	912	1,115	—	48,618	51,881	
Heterosexual contact ^e	1,144	1,568	—	1,130	1,524	—	1,072	1,475	—	976	1,422	—	29,851	37,707	
Perinatal	22	23	—	25	26	—	33	36	—	37	43	—	389	404	
Other ^f	1,786	79	—	1,625	79	—	1,542	80	—	1,357	67	—	51,160	8,785	
Subtotal	11,518	11,909	9.5	10,604	11,102	8.8	10,157	10,816	8.5	9,125	10,403	8.1	549,568	554,623	

Table 11b. Deaths of persons with diagnosed HIV infection ever classified as stage 3 (AIDS), by year of death and selected characteristics, 2009–2012 and cumulative—United States and 6 dependent areas (cont)

	2009			2010			2011			2012			Cumulative ^b		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Est. No. ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Female adult or adolescent															
Injection drug use	1,315	1,611	—	1,137	1,414	—	1,145	1,422	—	976	1,328	—	51,447	58,483	
Heterosexual contact ^e	1,862	2,528	—	1,706	2,377	—	1,608	2,303	—	1,465	2,255	—	48,711	60,544	
Perinatal	44	45	—	30	32	—	44	47	—	40	46	—	505	523	
Other ^f	860	28	—	796	25	—	775	29	—	710	25	—	20,939	3,700	
Subtotal	4,081	4,212	3.2	3,669	3,847	2.9	3,572	3,802	2.8	3,191	3,655	2.7	121,602	123,251	
Child (<13 yrs at death)															
Perinatal	3	3	—	0	0	—	0	0	—	1	1	—	4,714	4,724	
Other ^f	0	0	—	0	0	—	0	0	—	0	0	—	477	478	
Subtotal	3	3	0.0	0	0	0.0	0	0	0.0	1	1	0.0	5,191	5,202	
Region of residence															
Northeast	4,013	4,141	7.5	3,755	3,927	7.1	3,526	3,737	6.7	3,106	3,482	6.2	206,850	208,570	
Midwest	1,594	1,631	2.4	1,533	1,585	2.4	1,460	1,535	2.3	1,300	1,474	2.2	66,318	66,856	
South	7,235	7,494	6.6	6,480	6,810	5.9	6,252	6,688	5.8	5,656	6,550	5.6	249,947	253,080	
West	2,256	2,299	3.2	2,083	2,145	3.0	2,087	2,184	3.0	1,980	2,206	3.0	129,294	130,001	
U.S. dependent areas	504	559	12.7	422	482	11.7	404	475	11.6	275	346	8.6	23,952	24,569	
Total^g	15,602	16,124	5.2	14,273	14,950	4.8	13,729	14,618	4.6	12,317	14,059	4.4	676,361	683,076	

Note. Deaths of persons with diagnosed HIV infection may be due to any cause.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited. Rates are not calculated by transmission category because of the lack of denominator data.

^b From the beginning of the epidemic through 2012.

^c Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^d Hispanics/Latinos can be of any race.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 12a. Survival for more than 12, 24, and 36 months after a diagnosis of HIV infection during 2004–2009, by selected characteristics—United States

	No. of persons	Proportion survived (in months)		
		>12	>24	>36
Age at diagnosis (yr)				
<13	1,577	0.98	0.98	0.98
13–14	271	0.98	0.98	0.98
15–19	11,013	0.99	0.99	0.98
20–24	33,984	0.99	0.98	0.98
25–29	38,137	0.98	0.97	0.96
30–34	37,674	0.97	0.96	0.95
35–39	42,776	0.96	0.94	0.93
40–44	43,990	0.94	0.93	0.91
45–49	34,490	0.92	0.89	0.88
50–54	22,290	0.89	0.86	0.84
55–59	12,724	0.86	0.83	0.80
60–64	6,039	0.82	0.78	0.75
≥65	5,157	0.73	0.67	0.62
Race/ethnicity				
American Indian/Alaska Native	1,010	0.91	0.90	0.88
Asian	3,878	0.95	0.95	0.94
Black/African American	135,131	0.94	0.92	0.90
Hispanic/Latino ^a	57,987	0.95	0.94	0.93
Native Hawaiian/Other Pacific Islander	365	0.96	0.94	0.93
White	82,192	0.95	0.93	0.92
Multiple races	9,559	0.95	0.93	0.92
Transmission category				
Male adult or adolescent				
Male-to-male sexual contact	127,941	0.97	0.96	0.95
Injection drug use	14,360	0.90	0.87	0.84
Male-to-male sexual contact and injection drug use	9,927	0.96	0.95	0.93
Heterosexual contact ^b	21,013	0.94	0.91	0.90
Other ^c	43,715	0.88	0.86	0.84
Subtotal	216,956	0.94	0.93	0.91
Female adult or adolescent				
Injection drug use	8,465	0.93	0.90	0.88
Heterosexual contact ^b	36,159	0.96	0.95	0.94
Other ^c	26,965	0.92	0.90	0.89
Subtotal	71,589	0.94	0.93	0.91
Child (<13 yrs at diagnosis)				
Perinatal	1,288	0.98	0.98	0.98
Other ^d	289	0.99	0.99	0.99
Subtotal	1,577	0.98	0.98	0.98
Region of residence				
Northeast	59,549	0.94	0.93	0.92
Midwest	35,380	0.95	0.94	0.93
South	144,548	0.94	0.92	0.90
West	50,645	0.95	0.94	0.93
Year of diagnosis				
2004	49,135	0.93	0.91	0.90
2005	48,455	0.94	0.92	0.90
2006	48,752	0.94	0.92	0.91
2007	49,749	0.95	0.93	0.92
2008	48,323	0.95	0.94	0.92
2009	45,708	0.95	0.94	0.93
Total	290,122	0.94	0.93	0.91

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data exclude persons whose month of diagnosis or month of death is unknown.

See Technical Notes for method for calculating proportion of persons surviving.

^a Hispanics/Latinos can be of any race.

^b Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^c Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^d Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

Table 12b. Survival for more than 12, 24, and 36 months after a diagnosis of HIV infection during 2004–2009, by selected characteristics—United States and 6 dependent areas

	No. of persons	Proportion survived (in months)		
		>12	>24	>36
Age at diagnosis (yr)				
<13	1,609	0.98	0.98	0.98
13–14	281	0.98	0.98	0.98
15–19	11,156	0.99	0.99	0.98
20–24	34,482	0.99	0.98	0.98
25–29	38,896	0.98	0.97	0.96
30–34	38,519	0.97	0.96	0.95
35–39	43,717	0.95	0.94	0.93
40–44	44,962	0.94	0.92	0.91
45–49	35,272	0.92	0.89	0.87
50–54	22,870	0.89	0.86	0.83
55–59	13,074	0.86	0.82	0.80
60–64	6,232	0.82	0.78	0.74
≥65	5,398	0.73	0.67	0.62
Race/ethnicity				
American Indian/Alaska Native	1,010	0.91	0.90	0.88
Asian	3,879	0.95	0.95	0.94
Black/African American	135,251	0.94	0.92	0.90
Hispanic/Latino ^a	64,179	0.94	0.93	0.92
Native Hawaiian/Other Pacific Islander	374	0.95	0.94	0.93
White	82,208	0.95	0.93	0.92
Multiple races	9,567	0.95	0.93	0.92
Transmission category				
Male adult or adolescent				
Male-to-male sexual contact	129,286	0.97	0.96	0.95
Injection drug use	15,845	0.89	0.86	0.83
Male-to-male sexual contact and injection drug use	10,146	0.96	0.95	0.93
Heterosexual contact ^b	22,065	0.93	0.91	0.89
Other ^c	44,021	0.88	0.86	0.84
Subtotal	221,363	0.94	0.92	0.91
Female adult or adolescent				
Injection drug use	8,820	0.93	0.90	0.87
Heterosexual contact ^b	37,542	0.96	0.95	0.93
Other ^c	27,134	0.92	0.90	0.89
Subtotal	73,496	0.94	0.93	0.91
Child (<13 yrs at diagnosis)				
Perinatal	1,315	0.98	0.98	0.98
Other ^d	294	0.99	0.99	0.99
Subtotal	1,609	0.98	0.98	0.98
Region of residence				
Northeast	59,549	0.94	0.93	0.92
Midwest	35,380	0.95	0.94	0.93
South	144,548	0.94	0.92	0.90
West	50,645	0.95	0.94	0.93
U.S. dependent areas	6,346	0.86	0.84	0.82
Year of diagnosis				
2004	50,302	0.93	0.91	0.89
2005	49,711	0.94	0.92	0.90
2006	49,841	0.94	0.92	0.91
2007	50,738	0.95	0.93	0.92
2008	49,301	0.95	0.93	0.92
2009	46,575	0.95	0.94	0.93
Total	296,468	0.94	0.93	0.91

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Data exclude persons whose month of diagnosis or month of death is unknown.

See Technical Notes for method for calculating proportion of persons surviving.

^a Hispanics/Latinos can be of any race.

^b Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^c Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^d Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

Table 13a. Survival for more than 12, 24, and 36 months after a stage 3 (AIDS) classification during 2004–2009, by selected characteristics—United States

	No. of persons	Proportion survived (in months)		
		>12	>24	>36
Age at diagnosis (yr)				
<13	235	0.93	0.93	0.91
13–14	395	0.98	0.97	0.96
15–19	2,407	0.98	0.96	0.95
20–24	10,263	0.96	0.94	0.92
25–29	18,626	0.94	0.92	0.90
30–34	24,845	0.93	0.91	0.89
35–39	33,421	0.92	0.89	0.87
40–44	37,929	0.90	0.87	0.85
45–49	31,098	0.87	0.84	0.81
50–54	20,313	0.84	0.80	0.77
55–59	11,313	0.81	0.76	0.72
60–64	5,445	0.76	0.71	0.67
≥65	4,514	0.67	0.60	0.55
Race/ethnicity				
American Indian/Alaska Native	711	0.87	0.83	0.79
Asian ^a	2,227	0.91	0.90	0.89
Black/African American	94,381	0.88	0.84	0.81
Hispanic/Latino ^b	41,015	0.91	0.88	0.87
Native Hawaiian/Other Pacific Islander	213	0.92	0.88	0.87
White	54,093	0.89	0.87	0.84
Multiple races	8,164	0.91	0.88	0.85
Transmission category				
Male adult or adolescent				
Male-to-male sexual contact	78,517	0.92	0.90	0.88
Injection drug use	15,329	0.85	0.80	0.76
Male-to-male sexual contact and injection drug use	10,129	0.92	0.88	0.86
Heterosexual contact ^c	16,830	0.89	0.85	0.83
Other ^d	26,494	0.82	0.78	0.76
Subtotal	147,299	0.89	0.86	0.84
Female adult or adolescent				
Injection drug use	9,713	0.88	0.82	0.77
Heterosexual contact ^c	27,830	0.91	0.88	0.86
Other ^d	15,727	0.86	0.83	0.80
Subtotal	53,270	0.89	0.85	0.83
Child (<13 yrs at diagnosis)				
Perinatal	220	0.94	0.93	0.91
Other ^e	15	0.87	0.87	0.87
Subtotal	235	0.93	0.93	0.91
Region of residence				
Northeast	48,023	0.90	0.88	0.85
Midwest	23,018	0.90	0.87	0.85
South	95,321	0.88	0.84	0.81
West	34,442	0.91	0.88	0.87
Year of diagnosis				
2004	37,399	0.88	0.85	0.81
2005	35,331	0.89	0.85	0.83
2006	33,497	0.89	0.85	0.83
2007	32,422	0.89	0.86	0.84
2008	31,718	0.90	0.87	0.85
2009	30,437	0.90	0.88	0.86
Total	200,804	0.89	0.86	0.84

Note. Data exclude persons whose month of diagnosis or month of death is unknown. See Technical Notes for method for calculating proportion of persons surviving.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^d Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

Table 13b. Survival for more than 12, 24, and 36 months after a stage 3 (AIDS) classification during 2004–2009, by selected characteristics—United States and 6 dependent areas

	No. of persons	Proportion survived (in months)		
		>12	>24	>36
Age at diagnosis (yr)				
<13	240	0.92	0.92	0.90
13–14	406	0.98	0.97	0.96
15–19	2,453	0.98	0.96	0.95
20–24	10,397	0.95	0.94	0.92
25–29	18,969	0.94	0.92	0.90
30–34	25,407	0.93	0.91	0.89
35–39	34,153	0.91	0.89	0.87
40–44	38,831	0.90	0.87	0.85
45–49	31,886	0.87	0.83	0.81
50–54	20,865	0.84	0.80	0.77
55–59	11,623	0.80	0.76	0.72
60–64	5,615	0.76	0.70	0.66
≥65	4,705	0.66	0.60	0.55
Race/ethnicity				
American Indian/Alaska Native	711	0.87	0.83	0.79
Asian ^a	2,228	0.91	0.90	0.89
Black/African American	94,465	0.88	0.84	0.81
Hispanic/Latino ^b	45,656	0.89	0.86	0.85
Native Hawaiian/Other Pacific Islander	220	0.91	0.87	0.86
White	54,104	0.89	0.87	0.84
Multiple races	8,166	0.91	0.88	0.85
Transmission category				
Male adult or adolescent				
Male-to-male sexual contact	79,373	0.92	0.90	0.88
Injection drug use	16,729	0.84	0.79	0.75
Male-to-male sexual contact and injection drug use	10,352	0.91	0.88	0.85
Heterosexual contact ^c	17,575	0.88	0.85	0.82
Other ^d	26,596	0.82	0.78	0.76
Subtotal	150,625	0.89	0.86	0.84
Female adult or adolescent				
Injection drug use	10,069	0.87	0.82	0.77
Heterosexual contact ^c	28,835	0.91	0.88	0.85
Other ^d	15,781	0.86	0.82	0.80
Subtotal	54,685	0.89	0.85	0.82
Child (<13 yrs at diagnosis)				
Perinatal	225	0.92	0.92	0.90
Other ^e	15	0.87	0.87	0.87
Subtotal	240	0.92	0.92	0.90
Region of residence				
Northeast	48,023	0.90	0.88	0.85
Midwest	23,018	0.90	0.87	0.85
South	95,321	0.88	0.84	0.81
West	34,442	0.91	0.88	0.87
U.S. dependent areas	4,746	0.74	0.71	0.68
Year of diagnosis				
2004	38,314	0.88	0.84	0.81
2005	36,266	0.88	0.85	0.82
2006	34,303	0.88	0.85	0.83
2007	33,154	0.89	0.86	0.84
2008	32,437	0.89	0.87	0.84
2009	31,076	0.90	0.88	0.86
Total	205,550	0.89	0.86	0.83

Note. Data exclude persons whose month of diagnosis or month of death is unknown.

See Technical Notes for method for calculating proportion of persons surviving.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^d Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

Table 14a. Persons living with diagnosed HIV infection, by year and selected characteristics, 2009–2012—United States

	2009			2010			2011			2012		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
Age at end of year												
<13	3,197	3,221	6.0	2,885	2,912	5.5	2,614	2,648	5.0	2,470	2,519	4.8
13–14	1,301	1,306	16.1	1,161	1,166	14.1	1,028	1,034	12.5	859	867	10.5
15–19	7,410	7,470	34.7	7,083	7,160	32.6	6,626	6,737	31.1	6,119	6,290	29.5
20–24	26,729	27,033	125.5	28,976	29,412	135.5	30,545	31,208	140.9	31,771	32,839	145.4
25–29	47,791	48,230	222.5	49,269	49,859	235.8	51,482	52,319	245.8	54,004	55,297	258.5
30–34	65,716	66,172	332.7	68,032	68,628	342.0	69,973	70,780	345.1	71,796	72,942	348.9
35–39	95,578	96,051	467.7	89,414	89,962	448.1	85,746	86,433	441.0	84,421	85,332	437.8
40–44	141,659	142,124	677.1	136,695	137,237	656.5	130,466	131,118	623.2	124,660	125,457	596.6
45–49	165,890	166,186	727.9	169,300	169,611	749.3	169,715	170,107	767.5	166,420	166,862	769.4
50–54	125,820	125,877	578.4	135,840	135,879	607.9	144,886	144,905	642.1	154,249	154,130	682.7
55–59	79,328	79,258	417.7	87,483	87,347	441.3	96,794	96,581	476.7	106,574	106,150	510.9
60–64	39,939	39,877	252.2	46,620	46,492	273.6	53,361	53,134	298.2	60,279	59,815	335.8
≥65	30,227	30,123	76.1	34,693	34,503	85.2	40,019	39,708	96.0	46,919	46,327	107.4
Race/ethnicity												
American Indian/Alaska Native	2,501	2,504	106.1	2,606	2,608	115.0	2,722	2,727	119.1	2,863	2,866	124.1
Asian ^b	8,726	8,794	64.3	9,371	9,460	64.1	10,071	10,194	67.1	10,840	11,022	70.5
Black/African American	353,649	354,439	940.6	365,977	366,902	965.3	377,632	378,895	987.7	389,831	391,526	1,011.0
Hispanic/Latino ^c	163,240	164,107	338.9	169,554	170,612	336.2	175,772	177,113	341.5	182,493	184,237	347.8
Native Hawaiian/Other Pacific Islander	688	692	154.4	740	746	149.3	786	794	155.7	854	865	166.4
White	274,016	274,534	137.4	280,883	281,428	142.6	287,428	288,044	145.8	294,417	294,991	149.2
Multiple races	27,765	27,857	611.0	28,320	28,411	503.0	28,844	28,946	496.7	29,243	29,319	488.1
Transmission category												
Male adult or adolescent												
Male-to-male sexual contact	362,134	414,232	—	378,486	434,484	—	395,042	454,985	—	412,216	477,175	—
Injection drug use	69,310	85,033	—	68,070	84,105	—	66,881	83,120	—	65,812	82,195	—
Male-to-male sexual contact and injection drug use	44,699	49,929	—	44,700	50,081	—	44,577	50,084	—	44,525	50,098	—
Heterosexual contact ^d	52,570	65,857	—	54,169	68,246	—	55,659	70,424	—	56,997	72,482	—
Perinatal	3,555	3,560	—	3,758	3,762	—	3,925	3,927	—	4,052	4,049	—
Other ^e	87,845	3,300	—	92,651	3,282	—	96,968	3,262	—	102,125	3,268	—
Subtotal	620,113	621,911	501.7	641,834	643,961	515.0	663,052	665,802	527.5	685,727	689,267	540.7

Table 14a. Persons living with diagnosed HIV infection, by year and selected characteristics, 2009–2012—United States (cont)

	2009			2010			2011			2012		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Female adult or adolescent												
Injection drug use	40,109	53,717	—	39,571	53,459	—	38,928	53,078	—	38,394	52,693	—
Heterosexual contact ^d	107,327	148,349	—	110,040	153,837	—	112,417	158,976	—	114,512	163,955	—
Perinatal	3,843	3,848	—	4,077	4,081	—	4,251	4,254	—	4,392	4,391	—
Other ^e	55,996	1,882	—	59,044	1,917	—	61,993	1,955	—	65,046	2,002	—
Subtotal	207,275	207,795	160.7	212,732	213,294	162.4	217,589	218,262	164.8	222,344	223,041	167.0
Child (<13 yrs at end of year)												
Perinatal	2,856	2,875	—	2,534	2,555	—	2,254	2,280	—	2,067	2,101	—
Other ^e	341	346	—	351	358	—	360	368	—	403	418	—
Subtotal	3,197	3,221	6.0	2,885	2,912	5.5	2,614	2,648	5.0	2,470	2,519	4.8
Region of residence												
Northeast	221,239	221,651	400.9	225,057	225,488	407.2	228,631	229,180	412.2	232,708	233,447	418.6
Midwest	97,346	97,711	146.2	100,988	101,393	151.4	104,587	105,049	156.4	108,575	109,048	162.0
South	353,311	354,042	312.4	367,305	368,205	320.6	380,933	382,226	329.4	394,851	396,638	338.3
West	158,689	159,523	222.9	164,101	165,081	228.9	169,104	170,257	233.9	174,407	175,693	239.0
Total^f	830,585	832,927	271.3	857,451	860,167	278.1	883,255	886,712	284.6	910,541	914,826	291.5

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 14b. Persons living with diagnosed HIV infection, by year and selected characteristics, 2009–2012—United States and 6 dependent areas

	2009			2010			2011			2012		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Age at end of year												
<13	3,244	3,269	6.0	2,929	2,958	5.5	2,654	2,690	5.0	2,498	2,548	4.8
13–14	1,330	1,335	16.2	1,179	1,185	14.1	1,042	1,049	12.5	879	887	10.6
15–19	7,582	7,645	35.0	7,247	7,327	32.9	6,768	6,882	31.3	6,237	6,413	29.6
20–24	27,155	27,469	125.7	29,404	29,855	135.8	30,975	31,658	141.0	32,233	33,334	145.7
25–29	48,652	49,113	223.4	50,084	50,699	236.8	52,287	53,156	246.7	54,800	56,137	259.3
30–34	67,208	67,694	335.3	69,503	70,135	344.8	71,410	72,261	347.7	73,173	74,370	351.3
35–39	97,693	98,191	471.3	91,410	91,985	452.1	87,652	88,371	445.0	86,255	87,203	441.6
40–44	144,766	145,242	682.4	139,648	140,206	662.1	133,224	133,897	628.5	127,251	128,077	601.6
45–49	169,486	169,774	734.2	172,947	173,253	756.1	173,336	173,721	774.2	169,977	170,413	776.1
50–54	128,562	128,604	583.6	138,773	138,789	613.6	147,974	147,967	648.1	157,566	157,417	689.2
55–59	81,073	80,988	421.2	89,369	89,213	445.2	98,896	98,655	481.1	108,857	108,398	515.6
60–64	40,869	40,800	254.2	47,718	47,577	276.2	54,576	54,331	301	61,639	61,150	338.9
≥65	31,186	31,068	77.4	35,749	35,538	86.6	41,204	40,864	97.4	48,281	47,648	108.9
Race/ethnicity												
American Indian/Alaska Native	2,502	2,505	—	2,607	2,609	—	2,723	2,728	—	2,864	2,867	—
Asian ^b	8,780	8,848	—	9,427	9,516	—	10,125	10,248	—	10,893	11,075	—
Black/African American	353,989	354,783	—	366,325	367,254	—	377,990	379,258	—	390,186	391,885	—
Hispanic/Latino ^c	180,965	181,870	—	187,556	188,651	—	194,000	195,382	—	201,086	202,890	—
Native Hawaiian/Other Pacific Islander	700	705	—	755	762	—	802	811	—	871	883	—
White	274,088	274,607	—	280,954	281,500	—	287,499	288,116	—	294,490	295,064	—
Multiple races	27,782	27,874	—	28,336	28,427	—	28,859	28,961	—	29,256	29,332	—
Transmission category												
Male adult or adolescent												
Male-to-male sexual contact	365,489	417,721	—	382,050	438,213	—	398,776	458,913	—	416,182	481,385	—
Injection drug use	74,318	90,208	—	73,008	89,208	—	71,736	88,135	—	70,619	87,149	—
Male-to-male sexual contact and injection drug use	45,671	50,912	—	45,658	51,048	—	45,519	51,032	—	45,459	51,035	—
Heterosexual contact ^d	54,944	68,343	—	56,601	70,798	—	58,155	73,055	—	59,561	75,189	—
Perinatal	3,689	3,693	—	3,896	3,899	—	4,063	4,064	—	4,193	4,189	—
Other ^e	88,309	3,355	—	93,147	3,336	—	97,500	3,318	—	102,686	3,325	—
Subtotal	632,420	634,232	504.6	654,360	656,502	518.3	675,749	678,518	530.8	698,700	702,273	544.1

Table 14b. Persons living with diagnosed HIV infection, by year and selected characteristics, 2009–2012—United States and 6 dependent areas (cont)

	2009			2010			2011			2012		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate		No.	Rate
Female adult or adolescent												
Injection drug use	41,547	55,203	—	40,982	54,914	—	40,322	54,513	—	39,780	54,119	—
Heterosexual contact ^d	111,337	152,602	—	114,131	158,190	—	116,582	163,417	—	118,751	168,504	—
Perinatal	3,967	3,971	—	4,201	4,204	—	4,374	4,376	—	4,517	4,514	—
Other ^e	56,291	1,915	—	59,357	1,951	—	62,317	1,989	—	65,400	2,039	—
Subtotal	213,142	213,690	162.9	218,671	219,259	164.7	223,595	224,295	167.1	228,448	229,176	169.4
Child (<13 yrs at end of year)												
Perinatal	2,896	2,915	—	2,570	2,592	—	2,288	2,315	—	2,094	2,129	—
Other ^e	348	353	—	359	366	—	366	375	—	404	419	—
Subtotal	3,244	3,269	6.0	2,929	2,958	5.5	2,654	2,690	5.0	2,498	2,548	4.8
Region of residence												
Northeast	221,239	221,651	400.9	225,057	225,488	407.2	228,631	229,180	412.2	232,708	233,447	418.6
Midwest	97,346	97,711	146.2	100,988	101,393	151.4	104,587	105,049	156.4	108,575	109,048	162.0
South	353,311	354,042	312.4	367,305	368,205	320.6	380,933	382,226	329.4	394,851	396,638	338.3
West	158,689	159,523	222.9	164,101	165,081	228.9	169,104	170,257	233.9	174,407	175,693	239.0
U.S. dependent areas	18,221	18,264	415.7	18,509	18,552	450.6	18,743	18,790	460.5	19,105	19,170	474.0
Total^f	848,806	851,191	273.3	875,960	878,719	280.3	901,998	905,503	286.9	929,646	933,996	293.8

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 15a. Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by year and selected characteristics, 2009–2012—United States

	2009			2010			2011			2012		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
Age at end of year												
<13	532	536	1.0	443	447	0.8	361	365	0.7	301	305	0.6
13–14	458	461	5.7	318	321	3.9	249	251	3.0	188	190	2.3
15–19	2,638	2,659	12.3	2,498	2,525	11.5	2,241	2,274	10.5	1,926	1,963	9.2
20–24	6,872	6,964	32.3	7,468	7,605	35.0	7,974	8,165	36.9	8,168	8,414	37.3
25–29	15,424	15,616	72.0	15,779	16,045	75.9	16,111	16,461	77.3	16,462	16,916	79.1
30–34	27,366	27,632	138.9	27,685	28,038	139.7	27,953	28,409	138.5	28,239	28,806	137.8
35–39	48,561	48,901	238.1	44,732	45,139	224.8	42,060	42,549	217.1	40,509	41,081	210.8
40–44	81,793	82,198	391.6	77,706	78,191	374.0	73,059	73,617	349.9	68,812	69,418	330.1
45–49	103,252	103,592	453.7	105,043	105,442	465.8	104,377	104,848	473.1	101,313	101,762	469.3
50–54	81,366	81,533	374.7	87,675	87,883	393.2	93,224	93,453	414.1	98,684	98,822	437.7
55–59	52,614	52,670	277.6	57,858	57,906	292.5	63,920	63,951	315.6	70,216	70,125	337.5
60–64	26,747	26,769	169.3	31,114	31,119	183.2	35,437	35,395	198.7	39,971	39,793	223.4
≥65	20,301	20,292	51.3	23,349	23,312	57.6	26,932	26,845	64.9	31,480	31,249	72.4
Race/ethnicity												
American Indian/Alaska Native	1,351	1,355	57.4	1,411	1,416	62.4	1,476	1,482	64.7	1,538	1,543	66.8
Asian ^b	4,859	4,904	35.8	5,167	5,225	35.4	5,463	5,535	36.4	5,766	5,857	37.4
Black/African American	193,490	194,096	515.1	200,228	201,017	528.9	206,219	207,175	540.1	212,492	213,369	551.0
Hispanic/Latino ^c	98,943	99,725	206.0	102,241	103,172	203.3	105,224	106,309	205.0	108,263	109,461	206.7
Native Hawaiian/Other Pacific Islander	374	377	84.0	406	410	82.1	429	434	85.1	457	463	89.1
White	151,988	152,352	76.2	154,923	155,335	78.7	157,464	157,916	79.9	159,841	160,146	81.0
Multiple races	16,919	17,015	373.2	17,292	17,397	308.0	17,623	17,734	304.3	17,912	18,005	299.7
Transmission category												
Male adult or adolescent												
Male-to-male sexual contact	201,882	225,743	—	209,067	234,816	—	215,810	243,264	—	222,489	251,705	—
Injection drug use	47,736	56,571	—	47,112	56,192	—	46,477	55,712	—	45,884	55,201	—
Male-to-male sexual contact and injection drug use	29,735	32,450	—	29,835	32,675	—	29,824	32,776	—	29,793	32,802	—
Heterosexual contact ^d	32,365	39,710	—	33,534	41,412	—	34,608	42,920	—	35,570	44,351	—
Perinatal	2,058	2,067	—	2,140	2,150	—	2,210	2,221	—	2,255	2,265	—
Other ^e	43,645	2,248	—	46,109	2,227	—	48,216	2,211	—	50,635	2,203	—
Subtotal	357,421	358,788	289.4	367,797	369,472	295.5	377,145	379,103	300.3	386,626	388,527	304.8

Table 15a. Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by year and selected characteristics, 2009–2012—United States (cont)

	2009			2010			2011			2012		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
Female adult or adolescent												
Injection drug use	25,562	32,361	—	25,421	32,462	—	25,154	32,399	—	24,955	32,325	—
Heterosexual contact ^d	57,901	74,817	—	59,855	78,127	—	61,576	81,142	—	63,163	84,028	—
Perinatal	2,142	2,152	—	2,264	2,277	—	2,359	2,375	—	2,426	2,442	—
Other ^e	24,366	1,170	—	25,888	1,187	—	27,303	1,200	—	28,798	1,217	—
Subtotal	109,971	110,499	85.5	113,428	114,054	86.8	116,392	117,116	88.4	119,342	120,012	89.9
Child (<13 yrs at end of year)												
Perinatal	512	516	—	424	428	—	340	344	—	284	288	—
Other ^e	20	20	—	19	19	—	21	21	—	17	17	—
Subtotal	532	536	1.0	443	447	0.8	361	365	0.7	301	305	0.6
Region of residence												
Northeast	132,381	133,326	241.2	134,265	135,322	244.4	135,858	137,035	246.5	137,523	138,725	248.7
Midwest	50,643	50,793	76.0	52,479	52,664	78.6	54,268	54,486	81.1	56,098	56,295	83.6
South	189,674	189,829	167.5	196,909	197,202	171.7	203,565	203,980	175.8	210,291	210,539	179.6
West	95,226	95,875	134.0	98,015	98,784	137.0	100,207	101,083	138.8	102,357	103,285	140.5
Total^f	467,924	469,823	153.0	481,668	483,972	156.5	493,898	496,584	159.4	506,269	508,845	162.1

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 15b. Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by year and selected characteristics, 2009–2012—United States and 6 dependent areas

	2009			2010			2011			2012		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
Age at end of year												
<13	539	543	1.0	448	452	0.8	366	370	0.7	305	309	0.6
13–14	472	475	5.7	325	328	3.9	253	255	3.1	192	194	2.3
15–19	2,700	2,721	12.4	2,555	2,582	11.6	2,297	2,330	10.6	1,972	2,009	9.3
20–24	7,023	7,114	32.6	7,616	7,753	35.3	8,106	8,297	37.0	8,292	8,539	37.3
25–29	15,664	15,856	72.1	16,008	16,273	76.0	16,344	16,693	77.5	16,683	17,138	79.2
30–34	27,900	28,165	139.5	28,164	28,515	140.2	28,402	28,857	138.9	28,661	29,227	138.1
35–39	49,572	49,901	239.5	45,668	46,064	226.4	42,917	43,397	218.5	41,315	41,881	212.1
40–44	83,602	83,972	394.5	79,380	79,831	377.0	74,569	75,096	352.5	70,161	70,739	332.2
45–49	105,662	105,943	458.1	107,429	107,765	470.3	106,738	107,139	477.5	103,599	103,979	473.5
50–54	83,229	83,337	378.2	89,688	89,826	397.1	95,328	95,479	418.2	100,925	100,976	442.1
55–59	53,868	53,878	280.2	59,189	59,183	295.3	65,398	65,362	318.7	71,829	71,662	340.9
60–64	27,404	27,400	170.7	31,910	31,880	185.1	36,310	36,225	200.7	40,936	40,707	225.6
≥65	20,971	20,926	52.1	24,082	23,999	58.5	27,759	27,614	65.8	32,441	32,138	73.4
Race/ethnicity												
American Indian/Alaska Native	1,352	1,356	—	1,412	1,417	—	1,477	1,483	—	1,539	1,544	—
Asian ^b	4,883	4,927	—	5,191	5,249	—	5,486	5,557	—	5,788	5,878	—
Black/African American	193,674	194,277	—	200,417	201,203	—	206,409	207,362	—	212,686	213,559	—
Hispanic/Latino ^c	109,361	109,872	—	112,762	113,380	—	115,839	116,570	—	119,029	119,846	—
Native Hawaiian/Other Pacific Islander	381	384	—	415	419	—	439	444	—	468	474	—
White	152,027	152,391	—	154,964	155,376	—	157,505	157,956	—	159,883	160,187	—
Multiple races	16,928	17,024	—	17,301	17,406	—	17,632	17,743	—	17,918	18,010	—
Transmission category												
Male adult or adolescent												
Male-to-male sexual contact	203,865	227,712	—	211,110	236,842	—	217,903	245,335	—	224,666	253,865	—
Injection drug use	50,858	59,621	—	50,208	59,199	—	49,535	58,663	—	48,931	58,127	—
Male-to-male sexual contact and injection drug use	30,416	33,113	—	30,512	33,330	—	30,492	33,418	—	30,457	33,436	—
Heterosexual contact ^d	33,768	41,109	—	34,973	42,843	—	36,097	44,398	—	37,085	45,849	—
Perinatal	2,144	2,151	—	2,230	2,238	—	2,300	2,310	—	2,346	2,355	—
Other ^e	43,780	2,283	—	46,246	2,262	—	48,355	2,246	—	50,777	2,238	—
Subtotal	364,831	365,989	291.2	375,279	376,714	297.4	384,682	386,369	302.2	394,262	395,870	306.7

Table 15b. Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by year and selected characteristics, 2009–2012—United States and 6 dependent areas (cont)

	2009			2010			2011			2012		
	Estimated ^a			Estimated ^a			Estimated ^a			Estimated ^a		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate
Female adult or adolescent												
Injection drug use	26,477	33,265	—	26,329	33,355	—	26,050	33,275	—	25,851	33,197	—
Heterosexual contact ^d	60,090	77,018	—	62,090	80,370	—	63,860	83,429	—	65,495	86,367	—
Perinatal	2,227	2,236	—	2,348	2,360	—	2,445	2,459	—	2,512	2,526	—
Other ^e	24,442	1,180	—	25,968	1,199	—	27,384	1,212	—	28,886	1,230	—
Subtotal	113,236	113,699	86.7	116,735	117,283	88.1	119,739	120,375	89.7	122,744	123,320	91.1
Child (<13 yrs at end of year)												
Perinatal	518	521	—	428	432	—	344	348	—	288	292	—
Other ^e	21	21	—	20	20	—	22	22	—	17	17	—
Subtotal	539	543	1.0	448	452	0.8	366	370	0.7	305	309	0.6
Region of residence												
Northeast	132,381	133,326	241.2	134,265	135,322	244.4	135,858	137,035	246.5	137,523	138,725	248.7
Midwest	50,643	50,793	76.0	52,479	52,664	78.6	54,268	54,486	81.1	56,098	56,295	83.6
South	189,674	189,829	167.5	196,909	197,202	171.7	203,565	203,980	175.8	210,291	210,539	179.6
West	95,226	95,875	134.0	98,015	98,784	137.0	100,207	101,083	138.8	102,357	103,285	140.5
U.S. dependent areas	10,682	10,408	236.9	10,794	10,477	254.5	10,889	10,531	258.1	11,042	10,655	263.5
Total^f	478,606	480,231	154.2	492,462	494,449	157.7	504,787	507,115	160.7	517,311	519,500	163.4

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited. Rates are not calculated by transmission category because of the lack of denominator data.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 16a. Persons living with diagnosed HIV infection, by race/ethnicity and selected characteristics, year-end 2012—United States

	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c					
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d	Rate
Age at end of year																								
<13	6	6	1.3	55	57	2.3	1,636	1,669	23.0	374	380	2.9	3	3	2.9	294	299	1.1	102	104	4.8	2,470	2,519	4.8
13–14	0	0	0.0	7	7	2.0	569	575	49.1	174	175	9.5	0	0	0.0	75	76	1.7	34	34	12.4	859	867	10.5
15–19	14	14	7.7	54	55	5.8	3,978	4,095	130.6	1,212	1,243	27.2	3	3	8.3	653	666	5.6	205	213	34.2	6,119	6,290	29.5
20–24	101	102	51.9	307	321	27.3	19,582	20,236	609.2	5,782	6,001	131.2	32	34	69.7	4,826	4,964	38.9	1,141	1,181	229.9	31,771	32,839	145.4
25–29	177	179	106.3	726	752	57.8	28,595	29,249	1,051.2	11,739	12,073	276.9	69	71	149.0	10,732	10,965	88.9	1,966	2,009	505.2	54,004	55,297	258.5
30–34	289	292	188.1	1,103	1,132	85.0	33,328	33,797	1,260.2	17,317	17,676	411.3	104	106	242.7	17,116	17,366	144.1	2,539	2,574	737.3	71,796	72,942	348.9
35–39	325	325	228.2	1,624	1,661	125.7	36,808	37,144	1,507.3	21,147	21,461	539.1	120	122	324.2	21,556	21,750	193.2	2,841	2,869	1,000.8	84,421	85,332	437.8
40–44	471	471	316.4	1,999	2,029	158.0	50,776	51,045	1,947.1	28,378	28,661	782.0	118	119	338.3	38,761	38,956	299.5	4,157	4,175	1,566.7	124,660	125,457	596.6
45–49	511	512	322.6	1,769	1,789	158.6	65,127	65,257	2419	33,545	33,743	1052.4	139	140	410.7	59,841	59,930	421.4	5,488	5,491	2,285.3	166,420	166,862	769.4
50–54	453	452	277.6	1,274	1,287	125.4	61,280	61,197	2,263.6	27,581	27,646	1030.6	109	109	338.3	58,726	58,634	372.6	4,826	4,804	2,058.8	154,249	154,130	682.7
55–59	288	286	202.6	895	899	96.7	44,851	44,582	1,891.7	17,655	17,650	844.6	85	86	326.5	39,688	39,569	263.2	3,112	3,079	1,575.7	106,574	106,150	510.9
60–64	129	128	115.1	551	556	71.1	25,119	24,836	1,336.2	9,479	9,446	611.1	41	41	199.7	23,245	23,110	173.1	1,715	1,696	1,143.8	60,279	59,815	335.8
≥65	99	97	47.4	476	477	29.6	18,182	17,845	486.5	8,110	8,082	257.0	31	31	85.9	18,904	18,705	54.7	1,117	1,090	377.4	46,919	46,327	107.4
Transmission category																								
Male adult or adolescent																								
Male-to-male sexual contact	1,229	1,383	—	6,008	7,420	—	116,873	145,707	—	84,682	99,495	—	533	606	—	189,728	207,849	—	13,163	14,714	—	412,216	477,175	—
Injection drug use	192	216	—	255	378	—	31,983	42,255	—	17,421	21,780	—	24	27	—	13,831	15,141	—	2,106	2,398	—	65,812	82,195	—
Male-to-male sexual contact and injection drug use	281	316	—	289	344	—	13,172	16,010	—	8,313	9,548	—	34	37	—	20,075	21,254	—	2,361	2,590	—	44,525	50,098	—
Heterosexual contact ^e	143	167	—	617	777	—	35,003	46,795	—	11,438	13,600	—	35	40	—	8,158	9,257	—	1,603	1,845	—	56,997	72,482	—
Perinatal	4	4	—	22	22	—	2,397	2,394	—	978	978	—	1	1	—	515	516	—	135	134	—	4,052	4,049	—
Other ^f	244	14	—	1,672	72	—	53,443	940	—	21,516	467	—	78	3	—	22,814	1,663	—	2,358	109	—	102,125	3,268	—
Subtotal	2,093	2,099	231.5	8,863	9,014	145.4	252,871	254,101	1,717.6	144,348	145,868	720.1	705	715	340.9	255,121	255,680	307	21,726	21,791	1,189.1	685,727	689,267	540.7

Table 16a. Persons living with diagnosed HIV infection, by race/ethnicity and selected characteristics, year-end 2012—United States (cont)

	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c					
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d	Rate
Female adult or adolescent																								
Injection drug use	188	247	—	87	167	—	19,227	28,105	—	6,815	9,286	—	20	25	—	10,269	12,622	—	1,788	2,241	—	38,394	52,693	—
Heterosexual contact ^e	369	495	—	1,091	1,696	—	69,833	103,953	—	20,073	27,425	—	87	118	—	19,330	25,308	—	3,729	4,960	—	114,512	163,955	—
Perinatal	8	8	—	16	16	—	2,723	2,723	—	963	963	—	2	2	—	517	517	—	163	162	—	4,392	4,391	—
Other ^f	199	11	—	728	73	—	43,541	975	—	9,920	315	—	37	2	—	8,886	565	—	1,735	61	—	65,046	2,002	—
Subtotal	764	761	80.4	1,922	1,952	27.9	135,324	135,756	814.4	37,771	37,989	192.9	146	147	70.9	39,002	39,012	44.8	7,415	7,424	373.8	222,344	223,041	167.0
Child (<13 yrs at end of year)																								
Perinatal	5	5	—	40	41	—	1,361	1,384	—	347	352	—	3	3	—	221	224	—	90	92	—	2,067	2,101	—
Other ^f	1	1	—	15	16	—	275	285	—	27	28	—	0	0	—	73	76	—	12	12	—	403	418	—
Subtotal	6	6	1.3	55	57	2.3	1,636	1,669	23.0	374	380	2.9	3	3	2.9	294	299	1.1	102	104	4.8	2,470	2,519	4.8
Region of residence																								
Northeast	170	171	137.3	2,547	2,591	78.2	95,668	95,880	1,541.5	65,512	65,788	891.4	58	59	304.2	58,181	58,394	154.2	10,572	10,564	1,267.2	232,708	233,447	418.6
Midwest	371	370	93.3	936	948	50.8	47,527	47,784	689.6	11,098	11,272	230.4	40	41	146.6	44,574	44,578	85.6	4,029	4,056	352.9	108,575	109,048	162.0
South	721	728	95.5	1,964	2,007	57.0	219,056	220,029	989.2	54,994	55,610	288.3	144	147	190.8	106,956	107,051	154.2	11,016	11,066	571.2	394,851	396,638	338.3
West	1,601	1,597	155.6	5,393	5,476	78.9	27,580	27,833	835	50,889	51,568	240.9	612	619	156.3	84,706	84,967	221.6	3,626	3,633	174.1	174,407	175,693	239.0
Total^g	2,863	2,866	124.1	10,840	11,022	70.5	389,831	391,526	1,011.0	182,493	184,237	347.8	854	865	166.4	294,417	294,991	149.2	29,243	29,319	488.1	910,541	914,826	291.5

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 16b. Persons living with diagnosed HIV infection, by race/ethnicity and selected characteristics, year-end 2012—United States and 6 dependent areas

	American Indian/ Alaska Native		Asian ^a		Black/African American		Hispanic/Latino ^b		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Age at end of year																
<13	6	6	55	57	1,636	1,669	400	407	4	4	294	299	103	105	2,498	2,548
13–14	0	0	7	7	571	577	192	194	0	0	75	76	34	34	879	887
15–19	14	14	54	55	3,981	4,098	1,327	1,363	3	3	653	666	205	213	6,237	6,413
20–24	101	102	307	321	19,586	20,240	6,240	6,491	32	34	4,826	4,964	1,141	1,181	32,233	33,334
25–29	177	179	727	753	28,605	29,259	12,521	12,899	70	72	10,734	10,967	1,966	2,009	54,800	56,137
30–34	289	292	1,108	1,137	33,361	33,831	18,647	19,055	107	109	17,120	17,370	2,541	2,576	73,173	74,370
35–39	325	325	1,632	1,669	36,842	37,179	22,936	23,285	122	124	21,557	21,751	2,841	2,868	86,255	87,203
40–44	471	471	2,012	2,042	50,837	51,107	30,880	31,191	120	121	38,772	38,967	4,159	4,177	127,251	128,077
45–49	511	512	1,777	1,797	65,193	65,324	37,003	37,194	144	145	59,858	59,947	5,491	5,494	169,977	170,413
50–54	454	453	1,282	1,295	61,324	61,241	30,831	30,866	110	110	58,737	58,645	4,828	4,806	157,566	157,417
55–59	288	286	898	902	44,885	44,615	19,887	19,848	86	87	39,699	39,580	3,114	3,080	108,857	108,398
60–64	129	128	554	559	25,146	24,863	10,799	10,741	42	42	23,253	23,118	1,716	1,697	61,639	61,150
≥65	99	97	480	481	18,219	17,882	9,423	9,355	31	31	18,912	18,713	1,117	1,090	48,281	47,648
Transmission category																
Male adult or adolescent																
Male-to-male sexual contact	1,229	1,383	6,029	7,446	116,939	145,785	88,509	103,545	540	614	189,766	207,890	13,170	14,721	416,182	481,385
Injection drug use	192	216	256	386	32,003	42,303	22,203	26,672	24	28	13,835	15,146	2,106	2,398	70,619	87,149
Male-to-male sexual contact and injection drug use	281	316	290	346	13,174	16,014	9,237	10,472	34	37	20,082	21,261	2,361	2,590	45,459	51,035
Heterosexual contact ^e	144	168	619	782	35,047	46,870	13,950	16,221	37	43	8,160	9,260	1,604	1,846	59,561	75,189
Perinatal	4	4	22	22	2,400	2,397	1,116	1,115	1	1	515	516	135	134	4,193	4,189
Other ^f	244	14	1,688	72	53,515	941	21,984	524	79	3	22,818	1,663	2,358	109	102,686	3,325
Subtotal	2,094	2,100	8,904	9,054	253,078	254,310	156,999	158,549	715	725	255,176	255,735	21,734	21,799	698,700	702,273

Table 16b. Persons living with diagnosed HIV infection, by race/ethnicity and selected characteristics, year-end 2012—United States and 6 dependent areas (cont)

	American Indian/ Alaska Native		Asian ^a		Black/African American		Hispanic/Latino ^b		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Female adult or adolescent																
Injection drug use	188	247	87	169	19,236	28,123	8,188	10,687	20	25	10,273	12,628	1,788	2,241	39,780	54,119
Heterosexual contact ^e	369	495	1,094	1,706	69,903	104,079	24,224	31,817	90	124	19,339	25,320	3,732	4,963	118,751	168,504
Perinatal	8	8	16	16	2,727	2,727	1,084	1,082	2	2	517	517	163	162	4,517	4,514
Other ^f	199	11	737	73	43,606	978	10,191	347	40	2	8,891	565	1,736	62	65,400	2,039
Subtotal	764	761	1,934	1,964	135,472	135,906	43,687	43,933	152	154	39,020	39,030	7,419	7,428	228,448	229,176
Child (<13 yrs at end of year)																
Perinatal	5	5	40	41	1,361	1,384	372	378	4	4	221	224	91	93	2,094	2,129
Other ^f	1	1	15	16	275	285	28	29	0	0	73	76	12	12	404	419
Subtotal	6	6	55	57	1,636	1,669	400	407	4	4	294	299	103	105	2,498	2,548
Region of residence																
Northeast	170	171	2,547	2,591	95,668	95,880	65,512	65,788	58	59	58,181	58,394	10,572	10,564	232,708	233,447
Midwest	371	370	936	948	47,527	47,784	11,098	11,272	40	41	44,574	44,578	4,029	4,056	108,575	109,048
South	721	728	1,964	2,007	219,056	220,029	54,994	55,610	144	147	106,956	107,051	11,016	11,066	394,851	396,638
West	1,601	1,597	5,393	5,476	27,580	27,833	50,889	51,568	612	619	84,706	84,967	3,626	3,633	174,407	175,693
U.S. dependent areas	1	1	53	52	355	359	18,593	18,653	17	18	73	74	13	12	19,105	19,170
Total^g	2,864	2,867	10,893	11,075	390,186	391,885	201,086	202,890	871	883	294,490	295,064	29,256	29,332	929,646	933,996

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 17a. Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and selected characteristics, year-end 2012—United States

	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c					
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d	Rate
Age at end of year																								
<13	2	2	0.4	4	4	0.2	196	199	2.7	44	45	0.3	2	2	2.0	38	38	0.1	15	15	0.7	301	305	0.6
13–14	0	0	0.0	2	2	0.6	129	130	11.1	42	43	2.3	0	0	0.0	9	9	0.2	6	6	2.2	188	190	2.3
15–19	4	4	2.2	11	11	1.2	1,306	1,331	42.4	375	384	8.4	2	2	5.0	168	170	1.4	60	61	9.8	1,926	1,963	9.2
20–24	24	24	12.4	49	51	4.3	5,259	5,412	162.9	1,621	1,678	36.7	8	8	17.1	879	901	7.1	328	340	66.1	8,168	8,414	37.3
25–29	51	52	30.9	202	209	16.1	9,023	9,263	332.9	3,963	4,086	93.7	17	18	37.2	2,476	2,537	20.6	730	751	188.9	16,462	16,916	79.1
30–34	113	114	73.7	394	405	30.4	13,586	13,827	515.6	7,537	7,719	179.6	49	50	114.8	5,443	5,548	46.1	1,117	1,141	327.0	28,239	28,806	137.8
35–39	148	149	104.2	706	723	54.7	18,168	18,385	746.1	11,027	11,224	282.0	58	59	156.9	8,843	8,957	79.6	1,559	1,585	552.8	40,509	41,081	210.8
40–44	262	263	176.7	1,091	1,110	86.4	28,393	28,600	1,091.0	16,900	17,116	467.0	60	61	172.4	19,516	19,654	151.1	2,590	2,614	981.0	68,812	69,418	330.1
45–49	310	311	195.8	1,077	1,092	96.8	39,814	39,951	1,481.0	22,098	22,294	695.3	92	93	273.5	34,171	34,252	240.9	3,751	3,769	1,568.7	101,313	101,762	469.3
50–54	287	288	176.8	845	854	83.2	38,825	38,837	1,436.5	19,188	19,310	719.8	72	73	224.6	36,002	35,997	228.7	3,465	3,464	1,484.5	98,684	98,822	437.7
55–59	185	185	130.7	646	651	70.0	29,386	29,300	1,243.3	12,610	12,668	606.2	48	48	183.8	25,098	25,044	166.6	2,243	2,229	1,141.1	70,216	70,125	337.5
60–64	86	86	77.0	395	398	51.0	16,323	16,189	871.0	6,939	6,962	450.4	29	29	141.8	14,945	14,883	111.5	1,254	1,247	840.6	39,971	39,793	223.4
≥65	66	66	32.0	344	345	21.4	12,084	11,944	325.6	5,919	5,935	188.7	20	20	55.4	12,253	12,156	35.6	794	783	271.0	31,480	31,249	72.4
Transmission category																								
Male adult or adolescent																								
Male-to-male sexual contact	648	711	—	3,070	3,799	—	60,623	72,542	—	47,397	55,384	—	289	323	—	102,923	110,517	—	7,539	8,429	—	222,489	251,705	—
Injection drug use	122	134	—	161	244	—	22,327	27,889	—	12,558	15,315	—	15	16	—	9,098	9,795	—	1,603	1,808	—	45,884	55,201	—
Male-to-male sexual contact and injection drug use	189	206	—	188	222	—	9,267	10,699	—	5,795	6,558	—	20	21	—	12,667	13,268	—	1,667	1,827	—	29,793	32,802	—
Heterosexual contact ^e	83	97	—	426	529	—	21,350	27,846	—	7,746	9,164	—	21	24	—	4,868	5,454	—	1,076	1,238	—	35,570	44,351	—
Perinatal	2	2	—	10	10	—	1,371	1,374	—	530	536	—	1	1	—	261	262	—	80	80	—	2,255	2,265	—
Other ^f	112	11	—	932	57	—	25,441	532	—	12,301	334	—	36	2	—	10,394	1,196	—	1,419	70	—	50,635	2,203	—
Subtotal	1,156	1,161	128.0	4,787	4,862	78.4	140,379	140,883	952.3	86,327	87,291	430.9	382	387	184.8	140,211	140,492	168.7	13,384	13,452	734.0	386,626	388,527	304.8

Table 17a. Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and selected characteristics, year-end 2012—United States (cont)

	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c					
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d	Rate
Female adult or adolescent																								
Injection drug use	96	127	—	53	93	—	12,888	17,442	—	4,698	6,113	—	13	15	—	5,943	6,979	—	1,264	1,555	—	24,955	32,325	—
Heterosexual contact ^e	190	242	—	592	843	—	38,380	52,742	—	11,833	15,273	—	44	56	—	9,912	12,016	—	2,212	2,856	—	63,163	84,028	—
Perinatal	4	4	—	5	5	—	1,530	1,539	—	531	537	—	2	2	—	263	264	—	91	91	—	2,426	2,442	—
Other ^f	90	8	—	325	49	—	19,119	564	—	4,830	202	—	14	1	—	3,474	356	—	946	36	—	28,798	1,217	—
Subtotal	380	381	40.2	975	990	14.2	71,917	72,288	433.7	21,892	22,125	112.3	73	74	35.5	19,592	19,616	22.5	4,513	4,538	228.5	119,342	120,012	89.9
Child (<13 yrs at end of year)																								
Perinatal	1	1	—	4	4	—	184	187	—	43	44	—	2	2	—	35	35	—	15	15	—	284	288	—
Other ^f	1	1	—	0	0	—	12	12	—	1	1	—	0	0	—	3	3	—	0	0	—	17	17	—
Subtotal	2	2	0.4	4	4	0.2	196	199	2.7	44	45	0.3	2	2	2.0	38	38	0.1	15	15	0.7	301	305	0.6
Region of residence																								
Northeast	90	91	72.9	1,317	1,345	40.6	56,763	57,293	921.1	40,282	40,751	552.2	26	26	136.2	32,153	32,301	85.3	6,892	6,919	829.9	137,523	138,725	248.7
Midwest	177	177	44.6	449	454	24.3	24,445	24,556	354.4	6,266	6,332	129.4	19	19	70.4	22,483	22,486	43.2	2,259	2,272	197.7	56,098	56,295	83.6
South	345	346	45.5	915	928	26.4	115,003	115,051	517.2	30,679	30,933	160.3	54	55	71.0	56,747	56,645	81.6	6,548	6,581	339.7	210,291	210,539	179.6
West	926	929	90.5	3,085	3,130	45.1	16,281	16,469	494.1	31,036	31,445	146.9	358	363	91.6	48,458	48,715	127.1	2,213	2,234	107.1	102,357	103,285	140.5
Total^g	1,538	1,543	66.8	5,766	5,857	37.4	212,492	213,369	551.0	108,263	109,461	206.7	457	463	89.1	159,841	160,146	81.0	17,912	18,005	299.7	506,269	508,845	162.1

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 17b. Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and selected characteristics, year-end 2012—United States and 6 dependent areas

	American Indian/ Alaska Native		Asian ^a		Black/African American		Hispanic/Latino ^b		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Age at end of year																
<13	2	2	4	4	196	199	47	47	3	3	38	38	15	15	305	309
13–14	0	0	2	2	129	130	46	47	0	0	9	9	6	6	192	194
15–19	4	4	11	11	1,309	1,334	418	427	2	2	168	170	60	61	1,972	2,009
20–24	24	24	49	51	5,260	5,413	1,744	1,801	8	8	879	901	328	340	8,292	8,539
25–29	51	52	202	209	9,024	9,264	4,183	4,307	17	18	2,476	2,537	730	751	16,683	17,138
30–34	113	114	395	406	13,595	13,836	7,946	8,127	51	52	5,443	5,548	1,118	1,142	28,661	29,227
35–39	148	149	708	725	18,184	18,401	11,815	12,006	58	59	8,843	8,957	1,559	1,584	41,315	41,881
40–44	262	263	1,096	1,114	28,424	28,631	18,206	18,395	62	63	19,521	19,659	2,590	2,614	70,161	70,739
45–49	310	311	1,084	1,099	39,857	39,994	24,318	24,445	97	98	34,180	34,261	3,753	3,771	103,599	103,979
50–54	288	289	847	856	38,852	38,863	21,391	21,427	72	73	36,008	36,003	3,467	3,466	100,925	100,976
55–59	185	185	647	652	29,410	29,323	14,187	14,170	49	49	25,107	25,053	2,244	2,230	71,829	71,662
60–64	86	86	397	400	16,344	16,209	7,876	7,848	29	29	14,950	14,888	1,254	1,247	40,936	40,707
≥65	66	66	346	347	12,102	11,961	6,852	6,798	20	20	12,261	12,163	794	783	32,441	32,138
Transmission category																
Male adult or adolescent																
Male-to-male sexual contact	648	711	3,081	3,812	60,667	72,589	49,487	57,451	293	327	102,947	110,542	7,543	8,433	224,666	253,865
Injection drug use	122	134	162	247	22,342	27,913	15,587	18,210	15	17	9,100	9,797	1,603	1,808	48,931	58,127
Male-to-male sexual contact and injection drug use	189	206	189	224	9,269	10,702	6,452	7,184	20	22	12,671	13,272	1,667	1,827	30,457	33,436
Heterosexual contact ^e	84	98	427	531	21,381	27,889	9,226	10,615	22	25	4,869	5,455	1,076	1,237	37,085	45,849
Perinatal	2	2	10	10	1,372	1,375	620	625	1	1	261	262	80	80	2,346	2,355
Other ^f	112	11	937	57	25,469	532	12,407	370	37	2	10,396	1,196	1,419	70	50,777	2,238
Subtotal	1,157	1,162	4,806	4,880	140,500	141,001	93,779	94,454	388	393	140,244	140,524	13,388	13,455	394,262	395,870

Table 17b. Persons living with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and selected characteristics, year-end 2012—United States and 6 dependent areas (*cont*)

	American Indian/ Alaska Native		Asian ^a		Black/African American		Hispanic/Latino ^b		Native Hawaiian/ Other Pacific Islander		White		Multiple races		Total	
	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^c	No.	Est. No. ^{c,d}
Female adult or adolescent																
Injection drug use	96	127	53	93	12,892	17,450	5,587	6,974	13	15	5,945	6,982	1,265	1,556	25,851	33,197
Heterosexual contact ^e	190	242	594	846	38,418	52,802	14,117	17,539	46	60	9,917	12,022	2,213	2,857	65,495	86,367
Perinatal	4	4	5	5	1,533	1,542	614	618	2	2	263	264	91	91	2,512	2,526
Other ^f	90	8	326	49	19,147	565	4,885	214	16	1	3,476	356	946	36	28,886	1,230
Subtotal	380	381	978	993	71,990	72,359	25,203	25,345	77	78	19,601	19,625	4,515	4,540	122,744	123,320
Child (<13 yrs at end of year)																
Perinatal	1	1	4	4	184	187	46	46	3	3	35	35	15	15	288	292
Other ^f	1	1	0	0	12	12	1	1	0	0	3	3	0	0	17	17
Subtotal	2	2	4	4	196	199	47	47	3	3	38	38	15	15	305	309
Region of residence																
Northeast	90	91	1,317	1,345	56,763	57,293	40,282	40,751	26	26	32,153	32,301	6,892	6,919	137,523	138,725
Midwest	177	177	449	454	24,445	24,556	6,266	6,332	19	19	22,483	22,486	2,259	2,272	56,098	56,295
South	345	346	915	928	115,003	115,051	30,679	30,933	54	55	56,747	56,645	6,548	6,581	210,291	210,539
West	926	929	3,085	3,130	16,281	16,469	31,036	31,445	358	363	48,458	48,715	2,213	2,234	102,357	103,285
U.S. dependent areas	1	1	22	21	194	190	10,766	10,385	11	11	42	41	6	5	11,042	10,655
Total^g	1,539	1,544	5,788	5,878	212,686	213,559	119,029	119,846	468	474	159,883	160,187	17,918	18,010	517,311	519,500

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing transmission category, but not for incomplete reporting.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 18. Diagnoses of HIV infection, by area of residence, 2013—United States and 6 dependent areas

Area of residence	Adults and Adolescents			Children			Total		
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a	
		No.	Rate		No.	Rate		No.	Rate
Alabama	597	622	15.4	0	0	0.0	597	622	12.9
Alaska	23	24	3.9	0	0	0.0	23	24	3.2
Arizona	723	749	13.7	5	5	0.4	728	754	11.4
Arkansas	287	384	15.7	1	1	0.3	288	386	13.0
California	4,620	5,315	16.7	16	18	0.3	4,636	5,334	13.9
Colorado	317	329	7.5	4	4	0.5	321	333	6.3
Connecticut	333	347	11.4	1	1	0.2	334	348	9.7
Delaware	121	127	16.3	0	0	0.0	121	127	13.7
District of Columbia	486	612	109.2	0	0	0.0	486	612	94.6
Florida	5,188	5,364	32.1	12	13	0.4	5,200	5,377	27.5
Georgia	1,872	3,011	36.7	5	9	0.5	1,877	3,020	30.2
Hawaii	85	87	7.4	0	0	0.0	85	87	6.2
Idaho	27	28	2.1	1	2	0.5	28	29	1.8
Illinois	1,706	2,077	19.4	12	15	0.7	1,718	2,091	16.2
Indiana	485	506	9.3	2	2	0.2	487	508	7.7
Iowa	122	127	5.0	0	0	0.0	122	127	4.1
Kansas	148	155	6.5	2	2	0.4	150	157	5.4
Kentucky	388	403	11.0	2	2	0.3	390	405	9.2
Louisiana	1,356	1,398	36.6	1	1	0.1	1,357	1,399	30.3
Maine	30	31	2.7	1	1	0.6	31	32	2.4
Maryland	1,397	2,168	43.7	4	6	0.6	1,401	2,174	36.7
Massachusetts	663	992	17.4	2	3	0.3	665	995	14.9
Michigan	787	814	9.8	1	1	0.1	788	815	8.2
Minnesota	304	316	7.0	6	6	0.7	310	322	5.9
Mississippi	514	536	21.8	3	4	0.7	517	540	18.0
Missouri	466	482	9.6	2	2	0.2	468	484	8.0
Montana	21	22	2.5	0	0	0.0	21	22	2.1
Nebraska	75	77	5.0	0	0	0.0	75	77	4.1
Nevada	443	459	19.9	2	2	0.4	445	461	16.5
New Hampshire	32	33	2.9	0	0	0.0	32	33	2.5
New Jersey	1,387	2,169	29.0	5	8	0.5	1,392	2,177	24.5
New Mexico	144	149	8.7	0	0	0.0	144	149	7.2
New York	3,580	3,800	22.8	3	3	0.1	3,583	3,803	19.4
North Carolina	1,513	1,566	19.1	12	12	0.8	1,525	1,578	16.0
North Dakota	19	19	3.2	1	1	0.8	20	20	2.8
Ohio	1,165	1,200	12.4	8	8	0.4	1,173	1,208	10.4
Oklahoma	340	459	14.5	4	5	0.7	344	464	12.0
Oregon	222	230	6.9	0	0	0.0	222	230	5.8
Pennsylvania	1,365	1,419	13.1	5	6	0.3	1,370	1,425	11.2
Rhode Island	78	80	8.9	1	1	0.7	79	81	7.7
South Carolina	743	776	19.4	4	4	0.5	747	780	16.3
South Dakota	33	34	4.9	0	0	0.0	33	34	4.1
Tennessee	810	839	15.5	4	4	0.4	814	843	13.0
Texas	4,290	4,836	22.7	16	18	0.3	4,306	4,854	18.4
Utah	89	92	4.1	0	0	0.0	89	92	3.2
Vermont	13	14	2.5	0	0	0.0	13	14	2.2
Virginia	1,012	1,054	15.2	8	8	0.6	1,020	1,062	12.9
Washington	459	476	8.2	7	7	0.6	466	483	6.9
West Virginia	76	81	5.1	0	0	0.0	76	81	4.3
Wisconsin	255	263	5.5	1	1	0.1	256	264	4.6
Wyoming	14	14	3.0	0	0	0.0	14	14	2.5
Subtotal	41,223	47,165	17.9	164	187	0.4	41,387	47,352	15.0
U.S. dependent areas									
American Samoa	0	0	0.0	0	0	0.0	0	0	0.0
Guam	1	2	1.3	0	0	0.0	1	2	1.0
Northern Mariana Islands	0	0	0.0	0	0	0.0	0	0	0.0
Puerto Rico	607	761	24.9	0	0	0.0	607	761	21.1
Republic of Palau	0	0	0.0	0	0	0.0	0	0	0.0
U.S. Virgin Islands	23	29	33.3	0	0	0.0	23	29	28.0
Subtotal	631	792	23.5	0	0	0.0	631	792	19.8
Total^b	41,854	47,957	18.0	164	187	0.4	42,018	48,145	15.0

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^b Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 19. Stage 3 (AIDS), by area of residence, 2013 and cumulative—United States and 6 dependent areas

Area of residence	2013									Cumulative ^a		
	Adults and Adolescents			Children			Total			Estimated ^b		
	Estimated ^b			Estimated ^b			Estimated ^b			Adults and adolescents	Children	Total
	No.	No.	Rate ^c	No.	No.	Rate ^c	No.	No.	Rate ^c			
Alabama	439	538	13.3	0	0	0.0	439	538	11.1	11,477	76	11,553
Alaska	16	18	3.0	0	0	0.0	16	18	2.5	807	7	814
Arizona	361	384	7.0	0	0	0.0	361	384	5.8	13,538	47	13,585
Arkansas	169	192	7.8	0	0	0.0	169	192	6.5	5,015	35	5,050
California	2,430	2,724	8.6	1	1	0.0	2,431	2,725	7.1	171,102	692	171,795
Colorado	191	205	4.7	0	0	0.0	191	205	3.9	10,737	33	10,771
Connecticut	218	238	7.8	0	0	0.0	218	238	6.6	16,777	186	16,963
Delaware	96	108	13.8	0	0	0.0	96	108	11.6	4,390	27	4,417
District of Columbia	277	330	59.0	0	0	0.0	277	330	51.1	21,353	190	21,543
Florida	3,055	3,224	19.3	1	1	0.0	3,056	3,225	16.5	130,220	1,553	131,774
Georgia	1,338	1,646	20.1	1	1	0.1	1,339	1,648	16.5	46,159	259	46,417
Hawaii	58	67	5.7	0	0	0.0	58	67	4.7	3,489	17	3,506
Idaho	15	18	1.4	0	0	0.0	15	18	1.1	813	4	817
Illinois	868	983	9.2	0	0	0.0	868	983	7.6	42,111	283	42,394
Indiana	256	276	5.1	0	0	0.0	256	276	4.2	10,663	57	10,720
Iowa	81	87	3.4	0	0	0.0	81	87	2.8	2,304	14	2,318
Kansas	76	81	3.4	0	0	0.0	76	81	2.8	3,518	16	3,534
Kentucky	190	211	5.8	0	0	0.0	190	211	4.8	6,229	35	6,264
Louisiana	744	782	20.5	1	1	0.1	745	783	16.9	23,428	143	23,572
Maine	26	33	2.9	0	0	0.0	26	33	2.5	1,383	9	1,392
Maryland	969	1,286	25.9	0	0	0.0	969	1,286	21.7	39,617	331	39,948
Massachusetts	296	390	6.8	0	0	0.0	296	390	5.8	24,077	224	24,300
Michigan	473	515	6.2	0	0	0.0	473	515	5.2	18,884	118	19,002
Minnesota	163	173	3.9	0	0	0.0	163	173	3.2	6,420	30	6,450
Mississippi	410	438	17.8	0	0	0.0	410	438	14.6	9,069	57	9,126
Missouri	269	290	5.7	0	0	0.0	269	290	4.8	13,790	64	13,855
Montana	14	15	1.8	0	0	0.0	14	15	1.5	531	3	534
Nebraska	57	67	4.4	0	0	0.0	57	67	3.6	1,971	12	1,983
Nevada	250	271	11.7	0	0	0.0	250	271	9.7	7,581	29	7,610
New Hampshire	16	18	1.6	0	0	0.0	16	18	1.3	1,307	10	1,317
New Jersey	800	910	12.2	0	0	0.0	800	910	10.2	57,436	800	58,236
New Mexico	75	83	4.8	0	0	0.0	75	83	4.0	3,356	9	3,365
New York	2,104	2,319	13.9	0	0	0.0	2,104	2,319	11.8	202,274	2,447	204,721
North Carolina	898	958	11.7	1	1	0.1	899	959	9.7	22,728	133	22,862
North Dakota	11	12	2.0	0	0	0.0	11	12	1.7	198	2	200

Table 19. Stage 3 (AIDS), by area of residence, 2013 and cumulative—United States and 6 dependent areas (cont)

Area of residence	2013									Cumulative ^a		
	Adults and Adolescents			Children			Total			Estimated ^b		
	Estimated ^b			Estimated ^b			Estimated ^b			Adults and adolescents	Children	Total
	No.	No.	Rate ^c	No.	No.	Rate ^c	No.	No.	Rate ^c			
Ohio	539	575	5.9	0	0	0.0	539	575	5.0	19,627	148	19,775
Oklahoma	172	189	6.0	0	0	0.0	172	189	4.9	6,179	27	6,206
Oregon	134	146	4.4	0	0	0.0	134	146	3.7	7,308	17	7,325
Pennsylvania	777	892	8.2	1	1	0.1	778	893	7.0	41,157	371	41,529
Rhode Island	59	66	7.3	0	0	0.0	59	66	6.3	3,131	27	3,158
South Carolina	472	504	12.6	0	0	0.0	472	504	10.5	17,755	117	17,872
South Dakota	13	14	2.0	0	0	0.0	13	14	1.6	360	6	366
Tennessee	613	704	13.0	0	0	0.0	613	704	10.8	16,817	61	16,878
Texas	2,448	2,707	12.7	0	0	0.0	2,448	2,707	10.2	90,581	397	90,978
Utah	44	47	2.1	0	0	0.0	44	47	1.6	2,803	21	2,824
Vermont	4	4	0.8	0	0	0.0	4	4	0.7	544	6	550
Virginia	422	463	6.7	0	0	0.0	422	463	5.6	21,356	185	21,541
Washington	233	256	4.4	0	0	0.0	233	256	3.7	14,224	35	14,259
West Virginia	53	60	3.8	0	0	0.0	53	60	3.2	1,952	11	1,963
Wisconsin	137	146	3.0	1	1	0.1	138	147	2.6	5,738	37	5,775
Wyoming	14	16	3.3	0	0	0.0	14	16	2.7	331	2	333
Subtotal	23,843	26,680	10.1	7	8	0.0	23,850	26,688	8.4	1,184,618	9,421	1,194,039
U.S. dependent areas												
American Samoa	0	0	0.0	0	0	0.0	0	0	0.0	1	0	1
Guam	1	1	1.2	0	0	0.0	1	1	0.9	87	2	90
Northern Mariana Islands	0	0	0.0	0	0	0.0	0	0	0.0	2	0	2
Puerto Rico	331	433	14.2	0	0	0.0	331	433	12.0	34,379	403	34,782
Republic of Palau	0	0	0.0	0	0	0.0	0	0	0.0	2	0	2
U.S. Virgin Islands	9	13	14.2	0	0	0.0	9	13	12.0	778	17	795
Subtotal	341	447	13.3	0	0	0.0	341	447	11.2	35,250	422	35,672
Total^d	24,184	27,128	10.2	7	8	0.0	24,191	27,135	8.5	1,219,868	9,843	1,229,711

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a From the beginning of the epidemic through 2013.

^b Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting.

^c Rates are per 100,000 population.

^d Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 20. Adults and adolescents living with diagnosed HIV infection, by race/ethnicity and area of residence, year-end 2012—United States and 6 dependent areas

Area of residence	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d	Rate
Alabama	5	5	22.2	29	29	60.0	7,790	7,691	750.2	325	324	238.1	3	3	178.0	3,521	3,478	126.7	453	448	1116.5	12,126	11,979	298.0
Alaska	145	143	179.8	13	13	38.7	73	73	355.8	64	64	189.8	3	3	50.0	306	303	77.6	24	24	82.2	628	622	105.0
Arizona	474	472	228.3	128	128	81.0	1,293	1,284	620.1	3,719	3,710	254.4	13	13	136.3	6,970	6,927	212.0	225	223	294.5	12,822	12,757	237.0
Arkansas	7	7	41.5	10	10	31.2	2,164	2,201	613.1	278	283	204.8	4	4	92.4	2,320	2,344	126.4	155	157	516.7	4,938	5,006	205.5
California	414	419	299.9	4,097	4,179	96.0	20,318	20,602	1,113.0	38,879	39,587	358.0	272	279	242.9	50,021	50,466	381.1	2,264	2,281	346.4	116,265	117,814	375.0
Colorado	66	65	232.5	98	98	80.8	1,609	1,604	983.9	2,278	2,273	284.0	10	10	196.8	7,358	7,334	236.3	175	174	256.6	11,594	11,558	269.5
Connecticut	8	8	133.8	77	77	64.3	3,338	3,330	1,166.7	3,335	3,326	853.1	4	4	396.4	3,340	3,329	151.1	160	160	457.2	10,262	10,234	336.6
Delaware	3	3	119.1	8	8	31.2	1,883	1,869	1,192.6	232	231	408.3	0	0	0.0	829	825	159.6	74	74	674.8	3,029	3,010	391.0
District of Columbia	13	14	1,062.4	82	85	398.0	11,126	11,512	4,434.9	970	1,006	1,980.2	8	8	3,456.9	2,411	2,489	1,196.8	379	391	3,918.9	14,989	15,505	2,812.4
Florida	66	66	152.6	360	361	86.0	47,938	47,649	1,978.8	20,518	20,445	558.9	35	35	353.9	28,297	28,118	288.5	1,995	1,976	1,065.6	99,209	98,650	599.0
Georgia	16	16	85.6	151	162	57.1	25,799	26,321	1,087.7	3,342	3,464	540.3	12	13	286.6	7,612	7,686	165.2	1,425	1,440	1,521.8	38,357	39,102	481.8
Hawaii	7	7	232.1	378	378	82.0	115	114	512.3	222	222	234.9	233	232	223.8	1,229	1,223	426.0	186	185	95.2	2,370	2,361	202.6
Idaho	13	13	90.4	6	6	36.1	36	36	492.2	113	113	87.6	0	0	0.0	659	655	59.7	19	19	105.1	846	843	65.6
Illinois	39	41	255.8	372	385	73.6	15,942	16,461	1,101.6	5,830	6,018	387.1	23	24	853.6	9,855	10,085	144.2	1,629	1,673	1,538.3	33,690	34,686	324.4
Indiana	6	6	48.0	63	63	67.2	3,307	3,295	698.9	725	726	251.6	0	0	0.0	4,904	4,879	109.1	301	299	507.3	9,306	9,268	171.6
Iowa	3	3	42.8	39	39	80.4	342	341	486.5	179	179	158.9	0	0	0.0	1,260	1,254	54.7	53	53	221.8	1,876	1,868	73.2
Kansas	10	10	51.7	26	26	44.3	730	728	542.3	425	425	192.7	1	1	62.6	1,502	1,494	79.3	123	122	309.9	2,817	2,805	119.0
Kentucky	6	6	77.3	29	29	67.1	1,869	1,851	665.7	323	323	328.0	0	0	0.0	3,237	3,208	100.9	188	187	494.1	5,652	5,604	153.7
Louisiana	22	22	93.0	59	59	92.8	12,631	12,561	1,074.3	779	778	480.8	5	5	357.3	4,721	4,689	200.5	318	315	897.1	18,535	18,430	485.8
Maine	7	7	99.5	4	4	29.8	129	129	1,069.8	57	57	422.1	0	0	0.0	979	980	90.2	14	14	109.7	1,190	1,191	104.1
Maryland	27	29	238.2	165	176	60.7	22,772	23,335	1,650.8	1,537	1,604	415.7	14	14	601.7	4,549	4,658	170.0	1,229	1,250	1,608.4	30,293	31,067	631.1
Massachusetts	18	19	193.3	295	319	99.3	5,235	5,531	1,554.2	4,405	4,636	898.3	2	2	102.5	7,698	8,072	183.9	293	307	456.7	17,946	18,886	333.6
Michigan	33	33	71.0	81	81	38.6	8,556	8,374	743.5	712	703	215.9	2	2	99.6	4,906	4,821	74.6	723	714	600.7	15,013	14,728	177.6
Minnesota	120	119	271.3	108	107	59.4	2,380	2,372	1,110.2	661	659	361.5	7	7	399.6	3,645	3,626	96.1	158	157	260.8	7,079	7,047	158.1
Mississippi	10	10	90.3	13	13	56.5	6,499	6,415	732.6	232	231	367.6	2	2	243.3	1,781	1,759	120.5	324	322	1,878.5	8,861	8,753	357.2
Missouri	14	14	67.1	59	59	68.0	5,115	5,088	912.8	612	610	379.3	2	2	41.3	5,747	5,706	138.5	261	257	384.3	11,810	11,737	234.0
Montana	23	23	50.7	0	0	0.0	13	13	305.7	17	17	74.0	0	0	0.0	325	324	43.1	15	15	99.7	393	392	46.4
Nebraska	21	21	182.8	26	26	90.2	482	481	741.5	244	244	197.7	2	2	235.2	994	991	78.0	40	40	242.8	1,809	1,805	119.0
Nevada	43	43	211.9	196	196	109.3	1,633	1,621	891.1	1,666	1,661	299.1	22	23	172.1	3,666	3,637	284.9	177	176	336.9	7,403	7,357	322.8
New Hampshire	0	0	0.0	14	14	57.2	149	149	1,175.6	145	144	484.2	0	0	0.0	821	818	77.9	35	35	298.4	1,164	1,160	102.5
New Jersey	16	16	155.1	336	353	54.7	17,734	18,167	1,923.0	9,352	9,641	749.8	11	11	520.1	7,690	7,875	176.6	1,998	2,051	2,644.2	37,137	38,115	513.3
New Mexico	204	202	143.6	8	8	32.0	130	129	411.8	1,241	1,236	163.0	0	0	0.0	942	935	127.0	49	49	241.1	2,574	2,558	149.4
New York	85	85	180.0	1,593	1,597	121.6	52,660	52,397	2,206.5	42,618	42,451	1,504.1	16	16	231.9	26,388	26,251	268.4	6,627	6,582	3,310.0	129,987	129,379	782.0
North Carolina	180	179	196.7	108	108	57.2	17,191	17,092	1,005.7	1,736	1,736	294.3	10	10	220.5	6,638	6,604	121.7	711	708	745.6	26,574	26,437	326.5
North Dakota	8	8	29.8	2	2	29.6	38	38	487.6	10	10	81.3	0	0	0.0	135	135	25.7	10	10	145.4	203	203	34.6

Table 20. Adults and adolescents living with diagnosed HIV infection, by race/ethnicity and area of residence, year-end 2012—United States and 6 dependent areas (cont)

Area of residence	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d	Rate
Ohio	15	15	78.6	74	74	43.0	8,225	8,200	726.4	978	976	361.9	1	1	33.5	8,822	8,790	110.6	563	563	464.2	18,678	18,619	192.7
Oklahoma	269	272	111.4	34	35	60.1	1,227	1,248	556.3	477	487	197.4	6	6	176.9	2,979	3,013	135.8	253	257	191.8	5,245	5,319	169.9
Oregon	43	43	118.1	82	82	63.6	358	357	646.8	689	687	204.6	14	14	126.0	3,961	3,940	149.2	103	102	131.8	5,250	5,225	159.1
Pennsylvania	28	28	187.3	183	182	58.2	15,561	15,308	1,391.1	4,909	4,839	839.0	23	23	905.3	9,873	9,676	111.0	1,373	1,343	1,243.6	31,950	31,399	289.9
Rhode Island	8	8	220.2	31	31	111.5	492	491	1,055.3	536	536	514.0	1	1	190.8	972	972	138.2	44	44	348.3	2,084	2,083	231.8
South Carolina	11	11	75.5	30	30	55.1	10,733	10,695	1,013.4	584	585	326.9	2	2	102.5	3,533	3,519	135.4	284	284	726.7	15,177	15,125	383.5
South Dakota	72	71	140.6	3	3	41.2	97	97	924.8	18	18	99.3	0	0	0.0	265	264	45.0	9	9	100.2	464	462	67.7
Tennessee	9	9	60.6	55	54	66.1	9,279	9,069	1,042.1	667	665	306.0	9	9	361.7	5,862	5,735	138.5	325	320	567.5	16,206	15,862	294.6
Texas	55	56	77.0	578	593	68.0	25,848	26,256	1,073.6	21,161	21,615	291.7	25	26	161.0	20,912	21,193	213.1	2,235	2,272	1,091.1	70,814	72,010	343.4
Utah	21	21	96.4	32	32	63.6	210	210	991.2	485	484	183.1	4	4	21.3	1,712	1,707	95.5	49	49	165.4	2,513	2,507	114.3
Vermont	0	0	0.0	3	3	41.0	42	43	866.7	24	24	300.6	1	1	680.3	376	376	73.6	8	8	114.3	454	455	84.2
Virginia	19	19	103.1	234	235	59.3	12,991	12,933	1,007.9	1,644	1,642	317.2	9	9	189.2	6,544	6,523	144.7	562	559	485.6	22,003	21,919	320.3
Washington	137	136	184.4	336	336	76.5	1,621	1,616	802.1	1,400	1,398	245.5	39	39	119.6	7,339	7,298	171.0	317	313	190.2	11,189	11,137	193.7
West Virginia	2	2	61.8	4	4	34.2	436	433	801.4	59	59	314.1	0	0	0.0	1,070	1,067	72.2	60	60	380.7	1,631	1,625	102.7
Wisconsin	28	28	71.0	70	70	65.3	2,042	2,034	736.9	669	667	270.7	1	1	72.8	2,480	2,473	60.8	136	136	289.9	5,426	5,408	113.0
Wyoming	8	8	87.9	3	3	71.0	14	14	208.9	38	38	93.9	0	0	0.0	167	166	40.5	10	10	174.2	240	239	50.1
Subtotal	2,857	2,860	154.3	10,785	10,966	83.1	388,195	389,857	1,239.1	182,119	183,856	460.2	851	862	206.5	294,123	294,691	173.0	29,141	29,215	765.1	908,071	912,308	349.5
U.S. dependent areas^e																								
American Samoa	0	0	0.0	1	1	0.0	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0	1	1	2.4
Guam	0	0	0.0	48	47	0.0	5	5	0.0	4	4	0.0	11	11	0.0	9	9	0.0	7	7	0.0	84	83	67.8
Northern Mariana Islands	0	0	0.0	0	0	0.0	0	0	0.0	0	0	0.0	2	2	0.0	0	0	0.0	0	0	0.0	2	2	5.7
Puerto Rico	1	1	0.0	4	4	0.0	7	7	0.0	18,350	18,405	0.0	1	1	0.0	23	24	0.0	0	0	0.0	18,386	18,442	600.2
Republic of Palau	0	0	0.0	0	0	0.0	1	1	0.0	1	1	0.0	2	2	0.0	0	0	0.0	0	0	0.0	4	4	23.9
U.S. Virgin Islands	0	0	0.0	0	0	0.0	342	346	0.0	212	216	0.0	0	0	0.0	41	41	0.0	5	4	0.0	600	608	688.7
Subtotal	1	1	0.0	53	52	0.0	355	359	0.0	18,567	18,626	0.0	16	17	0.0	73	74	0.0	12	11	0.0	19,077	19,141	565.8
Total^f	2,858	2,861	0.0	10,838	11,018	0.0	388,550	390,217	0.0	200,686	202,483	0.0	867	879	0.0	294,196	294,765	0.0	29,153	29,226	0.0	927,148	931,449	352.3

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 21. Adults and adolescents living with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and area of residence, year-end 2012—United States and 6 dependent areas

Area of residence	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d	Rate
Alabama	1	1	4.5	10	10	21.1	3,152	3,166	308.8	153	154	113.1	2	2	124.2	1,381	1,375	50.1	198	201	501.3	4,897	4,909	122.1
Alaska	90	90	113.6	11	11	33.3	41	41	201.9	46	47	138.7	2	2	34.3	170	170	43.4	14	14	47.8	374	375	63.3
Arizona	269	269	130.0	54	54	34.2	614	615	296.8	1,862	1,864	127.9	9	9	95.6	3,345	3,334	102.0	120	120	158.7	6,273	6,265	116.4
Arkansas	5	5	29.3	5	5	15.5	975	990	275.9	137	138	100.1	3	3	68.9	1,134	1,141	61.5	91	92	304.8	2,350	2,375	97.5
California	243	246	175.9	2,398	2,438	56.0	12,611	12,792	691.0	24,485	24,878	225.0	143	146	127.1	30,722	30,990	234.0	1,418	1,435	217.9	72,020	72,925	232.1
Colorado	32	32	114.1	48	48	39.8	727	727	446.1	1,197	1,199	149.9	4	4	78.7	2,940	2,938	94.7	99	100	147.7	5,047	5,049	117.7
Connecticut	5	5	83.8	37	37	31.0	2,284	2,297	804.6	2,300	2,313	593.3	4	4	406.7	2,195	2,203	100.0	148	149	424.9	6,973	7,008	230.5
Delaware	2	2	79.7	5	5	20.2	1,231	1,232	786.1	141	142	250.2	0	0	0.0	512	514	99.4	46	46	422.9	1,937	1,941	252.1
District of Columbia	5	5	393.0	44	45	207.7	6,883	7,007	2,699.4	584	594	1,169.9	4	4	1,729.2	1,186	1,202	578.2	263	267	2,683.3	8,969	9,125	1,655.2
Florida	23	23	52.9	165	166	39.5	27,672	27,589	1,145.7	11,019	11,005	300.8	11	11	111.3	16,084	16,008	164.3	1,298	1,294	698.0	56,272	56,096	340.6
Georgia	9	8	43.5	66	68	23.8	13,989	13,726	567.2	2,032	2,060	321.4	4	4	91.2	4,216	4,101	88.1	948	945	998.6	21,264	20,912	257.6
Hawaii	3	3	98.5	225	227	49.2	80	81	362.8	143	144	152.8	155	156	150.2	732	733	255.2	110	110	56.9	1,448	1,454	124.7
Idaho	5	5	35.4	3	3	18.1	18	18	244.5	62	62	48.3	0	0	0.0	311	311	28.3	12	12	66.8	411	411	32.0
Illinois	14	14	89.3	189	192	36.8	8,577	8,763	586.5	3,366	3,427	220.5	12	12	448.1	4,988	5,047	72.2	949	965	887.1	18,095	18,421	172.3
Indiana	0	0	0.0	42	42	45.0	1,669	1,674	355.0	417	419	145.5	0	0	0.0	2,639	2,640	59.0	170	171	289.4	4,937	4,946	91.6
Iowa	1	1	14.4	16	16	33.4	234	236	335.8	115	115	102.5	0	0	0.0	761	758	33.1	32	32	134.5	1,159	1,158	45.4
Kansas	6	6	31.2	12	12	20.5	382	382	285.0	275	276	125.4	0	0	0.0	879	876	46.5	74	74	187.9	1,628	1,627	69.0
Kentucky	3	3	38.8	13	13	30.2	910	907	326.1	206	207	210.3	0	0	0.0	1,771	1,765	55.5	113	113	299.4	3,016	3,008	82.5
Louisiana	12	12	51.0	23	23	36.2	6,730	6,737	576.1	415	417	257.4	2	2	142.9	2,583	2,579	110.3	171	170	485.4	9,936	9,940	262.0
Maine	4	4	60.7	3	3	26.4	62	64	526.5	33	33	245.5	0	0	0.0	555	562	51.7	11	11	86.9	668	677	59.2
Maryland	7	8	61.6	72	74	25.6	12,664	12,713	899.3	913	933	241.8	6	6	254.8	2,309	2,307	84.2	791	791	1,017.3	16,762	16,831	341.9
Massachusetts	10	11	106.1	170	180	56.1	3,238	3,405	956.8	2,689	2,820	546.4	1	1	50.9	4,409	4,610	105.1	183	191	284.5	10,700	11,217	198.2
Michigan	13	13	27.8	36	36	17.3	4,465	4,358	386.9	392	388	119.0	2	2	100.0	2,561	2,510	38.8	409	404	340.0	7,878	7,710	93.0
Minnesota	64	64	145.3	53	53	29.4	1,132	1,137	532.2	356	357	195.6	2	2	114.8	1,543	1,540	40.8	75	75	124.4	3,225	3,227	72.4
Mississippi	4	4	35.6	5	5	21.4	2,929	2,893	330.3	116	115	183.3	1	1	122.0	797	787	53.9	191	193	1,122.0	4,043	3,997	163.1
Missouri	9	9	42.9	25	25	29.0	2,722	2,720	488.0	311	311	193.0	1	1	20.7	3,008	2,994	72.7	159	158	235.7	6,235	6,218	123.9
Montana	16	16	35.2	0	0	0.0	8	8	187.3	14	14	61.1	0	0	0.0	207	208	27.7	10	10	68.2	255	256	30.3
Nebraska	10	10	88.1	15	15	52.3	267	268	413.6	160	161	129.9	2	2	236.1	507	507	39.9	29	29	177.7	990	992	65.4
Nevada	23	23	114.0	81	81	45.2	799	797	437.9	850	849	153.0	10	10	76.8	1,720	1,710	133.9	100	101	193.0	3,583	3,571	156.7
New Hampshire	0	0	0.0	6	6	24.6	77	78	612.6	84	84	283.5	0	0	0.0	426	427	40.7	27	27	232.0	620	623	55.0
New Jersey	11	11	105.9	164	168	26.0	9,275	9,430	998.1	4,908	4,996	388.5	5	5	231.3	3,956	4,005	89.8	1,236	1,259	1,623.3	19,555	19,875	267.6
New Mexico	106	106	75.5	5	5	20.4	68	68	217.1	739	739	97.5	0	0	0.0	617	617	83.8	40	40	198.2	1,575	1,575	92.0
New York	45	45	95.4	809	823	62.7	32,222	32,547	1,370.6	26,960	27,234	964.9	4	4	58.8	14,162	14,182	145.0	4,363	4,377	2,200.9	78,565	79,212	478.8
North Carolina	78	78	86.1	32	32	17.1	7,142	7,184	422.7	781	785	133.1	1	1	21.7	2,649	2,660	49.0	321	324	341.6	11,004	11,065	136.7
North Dakota	7	7	26.5	2	2	29.8	13	13	164.9	6	6	50.7	0	0	0.0	57	57	10.8	4	4	58.3	89	89	15.2

Table 21. Adults and adolescents living with diagnosed HIV infection ever classified as stage 3 (AIDS), by race/ethnicity and area of residence, year-end 2012—United States and 6 dependent areas (cont)

Area of residence	American Indian/ Alaska Native			Asian ^a			Black/African American			Hispanic/Latino ^b			Native Hawaiian/ Other Pacific Islander			White			Multiple races			Total		
	Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c			Estimated ^c		
	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No.	Rate	No.	No. ^d	Rate
Ohio	7	7	37.7	29	29	16.9	3,925	3,942	349.3	495	498	184.6	0	0	0.1	4,213	4,227	53.2	283	285	235.2	8,952	8,988	93.0
Oklahoma	147	148	60.5	20	20	34.7	564	571	254.5	225	228	92.3	3	3	88.7	1,459	1,472	66.3	116	118	88.1	2,534	2,559	81.8
Oregon	25	25	68.9	51	51	40.0	219	220	398.1	464	468	139.3	7	7	63.3	2,456	2,463	93.3	68	68	88.3	3,290	3,303	100.6
Pennsylvania	9	9	60.8	101	100	32.0	9,189	9,054	822.8	2,898	2,855	495.1	10	10	389.4	5,586	5,441	62.4	885	865	801.3	18,678	18,335	169.3
Rhode Island	6	6	167.4	24	24	87.2	355	357	767.1	381	385	369.1	1	1	191.6	640	645	91.8	30	30	242.2	1,437	1,449	161.3
South Carolina	7	7	48.6	18	18	33.3	5,936	5,954	564.2	343	345	192.8	2	2	102.5	1,808	1,811	69.7	167	168	430.3	8,281	8,305	210.6
South Dakota	33	33	65.1	1	1	13.8	42	42	403.0	5	5	27.5	0	0	0.0	99	99	16.8	5	5	55.7	185	185	27.1
Tennessee	6	6	38.7	28	27	33.4	4,406	4,283	492.1	361	363	166.9	2	2	81.4	3,057	2,967	71.7	188	186	329.8	8,048	7,834	145.5
Texas	27	27	37.6	286	292	33.5	13,740	13,985	571.9	12,377	12,565	169.5	10	10	63.2	12,023	12,164	122.3	1,326	1,349	647.7	39,789	40,392	192.6
Utah	16	16	73.8	18	18	35.8	108	108	512.4	286	287	108.5	2	2	10.9	974	974	54.5	32	32	109.6	1,436	1,437	65.5
Vermont	0	0	0.0	2	2	28.0	26	26	530.3	15	15	189.3	1	1	683.1	217	218	42.7	7	7	103.0	268	269	49.9
Virginia	9	9	49.6	123	124	31.4	5,769	5,802	452.1	840	847	163.7	3	3	63.1	3,129	3,138	69.6	272	273	237.1	10,145	10,197	149.0
Washington	91	91	123.8	189	191	43.5	965	970	481.5	850	855	150.2	24	24	74.4	4,157	4,161	97.5	178	178	107.9	6,454	6,471	112.6
West Virginia	0	0	0.0	0	0	0.0	201	203	376.1	26	26	138.1	0	0	0.0	632	636	43.0	44	45	288.3	903	910	57.5
Wisconsin	12	12	30.7	28	28	26.1	982	985	357.1	362	363	147.2	0	0	0.0	1,222	1,225	30.1	64	64	137.1	2,670	2,678	56.0
Wyoming	6	6	66.1	0	0	0.0	7	7	106.5	24	24	59.8	0	0	0.0	99	99	24.0	9	9	163.1	145	145	30.5
Subtotal	1,536	1,541	83.2	5,762	5,853	44.3	212,296	213,170	677.5	108,219	109,416	273.9	455	461	110.4	159,803	160,108	94.0	17,897	17,990	471.1	505,968	508,539	194.8
U.S. dependent areas^e																								
American Samoa	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—	0	0	—	1	1	—	1	1	2.4
Guam	0	0	—	22	21	—	1	1	—	1	1	—	7	7	—	1	1	—	2	2	—	34	33	26.6
Northern Mariana Islands	0	0	—	0	0	—	0	0	—	0	0	—	2	2	—	0	0	—	0	0	—	2	2	5.4
Puerto Rico	1	1	—	0	0	—	2	2	—	10,645	10,266	—	0	0	—	14	14	—	0	0	—	10,662	10,283	334.7
Republic of Palau	0	0	—	0	0	—	0	0	—	0	0	—	1	1	—	0	0	—	0	0	—	1	1	5.6
U.S. Virgin Islands	0	0	—	0	0	—	191	187	—	117	115	—	0	0	—	27	27	—	3	2	—	338	332	375.9
Subtotal	1	1	—	22	21	—	194	190	—	10,763	10,383	—	10	10	—	42	41	—	6	5	—	11,038	10,651	314.8
Total^f	1,537	1,542	—	5,784	5,874	—	212,490	213,360	—	118,982	119,799	—	465	471	—	159,845	160,149	—	17,903	17,995	—	517,006	519,191	196.4

Note. Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^b Hispanics/Latinos can be of any race.

^c Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^d Because the estimated totals were calculated independently of the corresponding values for each subpopulation, the subpopulation values may not sum to the totals shown here.

^e Rates by race/ethnicity are not provided because U.S. census information for U.S. dependent areas is limited.

^f Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 22. Diagnoses of HIV infection, 2013, and persons living with diagnosed HIV infection (prevalence), year-end 2012, by metropolitan statistical area of residence—United States and Puerto Rico

MSA of residence	Diagnosis, 2013				Prevalence of diagnosed HIV infection, year-end 2012		
	No.	Estimated ^a		Rank ^b	No.	Estimated ^a	
		No.	Rate			No.	Rate
Akron, OH	48	49	7.0	87	816	814	115.4
Albany–Schenectady–Troy, NY	65	68	7.8	82	2,093	2,061	235.6
Albuquerque, NM	67	70	7.7	83	1,286	1,278	142.0
Allentown–Bethlehem–Easton, PA–NJ	62	65	7.9	80	1,439	1,418	171.5
Atlanta–Sandy Springs–Roswell, GA	1,214	1,919	34.7	5	25,546	26,020	477.0
Augusta–Richmond County, GA–SC	99	153	26.4	11	2,103	2,134	370.2
Austin–Round Rock, TX	287	333	17.7	32	4,750	4,851	264.3
Bakersfield, CA	102	118	13.7	49	1,736	1,769	206.7
Baltimore–Columbia–Towson, MD	650	1,010	36.5	3	17,992	18,288	664.1
Baton Rouge, LA	288	297	36.2	4	4,449	4,425	543.0
Birmingham–Hoover, AL	167	172	15.1	44	3,836	3,794	334.3
Boise City, ID	22	23	3.5	103	452	451	70.7
Boston–Cambridge–Newton, MA–NH ^c	495	735	15.7	40	12,686	13,346	287.5
Boston Division	286	427	22.0	—	7,567	7,959	413.2
Cambridge Division	199	297	12.8	—	4,830	5,100	222.4
Bridgeport–Stamford–Norwalk, CT	91	95	10.1	66	2,687	2,681	287.1
Buffalo–Cheektowaga–Niagara Falls, NY	128	132	11.6	59	2,271	2,229	196.6
Cape Coral–Fort Myers, FL	142	147	22.3	16	1,645	1,635	253.6
Charleston–North Charleston, SC	122	126	17.8	31	2,292	2,283	327.3
Charlotte–Concord–Gastonia, NC–SC	420	434	18.6	26	7,147	7,104	309.6
Chattanooga, TN–GA	47	52	9.7	72	1,150	1,130	210.0
Chicago–Naperville–Elgin, IL–IN–WI	1,548	1,885	19.8	25	29,701	30,604	321.7
Chicago Division	1,387	1,706	23.3	—	27,098	27,976	382.6
Elgin Division	33	38	6.1	—	605	618	98.7
Gary Division	78	84	11.9	—	1,176	1,171	165.8
Lake County Division	50	57	6.5	—	822	839	96.6
Cincinnati, OH–KY–IN	244	251	11.8	58	3,429	3,419	160.6
Cleveland–Elyria, OH	291	299	14.5	46	4,763	4,745	229.8
Colorado Springs, CO	30	31	4.5	99	814	809	121.0
Columbia, SC	170	176	22.2	17	3,935	3,917	499.3
Columbus, OH	302	311	15.8	39	5,031	5,019	258.0
Dallas–Fort Worth–Arlington, TX	1,197	1,389	20.4	21	21,781	22,254	332.0
Dallas Division	928	1,076	23.9	—	17,313	17,690	399.6
Fort Worth Division	269	312	13.5	—	4,468	4,564	200.6
Dayton, OH	87	89	11.1	63	1,422	1,419	176.7
Deltona–Daytona Beach–Ormond Beach, FL	95	98	16.2	37	1,461	1,451	243.8
Denver–Aurora–Lakewood, CO	244	253	9.4	73	9,081	9,054	342.1
Des Moines–West Des Moines, IA	35	36	6.0	93	603	602	102.1

Table 22. Diagnoses of HIV infection, 2013, and persons living with diagnosed HIV infection (prevalence), year-end 2012, by metropolitan statistical area of residence—United States and Puerto Rico (cont)

MSA of residence	Diagnosis, 2013				Prevalence of diagnosed HIV infection, year-end 2012		
	No.	Estimated ^a		Rank ^b	No.	Estimated ^a	
		No.	Rate			No.	Rate
Detroit–Warren–Dearborn, MI	528	544	12.7	54	9,817	9,610	223.9
Detroit Division	349	360	20.3	—	7,177	7,006	390.8
Warren Division	179	185	7.3	—	2,640	2,605	104.2
Durham–Chapel Hill, NC	106	109	20.5	20	2,034	2,020	384.4
El Paso, TX	118	137	16.5	36	1,782	1,825	219.4
Fresno, CA	91	106	11.1	64	1,590	1,623	171.3
Grand Rapids–Wyoming, MI	50	52	5.1	98	981	969	96.3
Greensboro–High Point, NC	132	137	18.5	27	2,359	2,344	318.4
Greenville–Anderson–Mauldin, SC	107	111	13.0	52	1,613	1,607	190.9
Harrisburg–Carlisle, PA	50	52	9.4	74	1,319	1,288	232.2
Hartford–West Hartford–East Hartford, CT	119	123	10.1	67	3,475	3,463	285.2
Honolulu (Urban), HI	57	59	6.0	94	1,642	1,635	167.7
Houston–The Woodlands–Sugar Land, TX	1,469	1,556	24.6	13	24,099	24,270	393.0
Indianapolis–Carmel–Anderson, IN	257	265	13.6	50	4,632	4,615	239.2
Jackson, MS	164	170	29.5	8	3,354	3,307	573.7
Jacksonville, FL	361	373	26.8	9	6,146	6,108	443.3
Kansas City, MO–KS	203	210	10.2	65	4,819	4,794	235.1
Knoxville, TN	60	63	7.4	84	1,177	1,151	135.8
Lakeland–Winter Haven, FL	100	104	16.6	35	1,773	1,762	286.0
Lancaster, PA	33	34	6.4	89	754	738	140.2
Las Vegas–Henderson–Paradise, NV	393	407	20.0	24	6,196	6,160	308.4
Little Rock–North Little Rock–Conway, AR	126	168	23.2	15	1,870	1,898	264.4
Los Angeles–Long Beach–Anaheim, CA	2,048	2,379	18.1	30	51,510	52,354	401.6
Anaheim Division	304	353	11.3	—	6,614	6,720	217.8
Los Angeles Division	1,744	2,026	20.2	—	44,896	45,633	458.5
Louisville/Jefferson County, KY–IN	184	190	15.1	42	2,997	2,965	236.9
Madison, WI	33	34	5.4	95	678	675	108.8
McAllen–Edinburg–Mission, TX	103	119	14.5	45	999	1,026	127.3
Memphis, TN–MS–AR	393	413	30.8	7	7,373	7,233	539.5
Miami–Fort Lauderdale–West Palm Beach, FL	2,502	2,582	44.3	1	51,630	51,349	891.0
Fort Lauderdale Division	836	862	46.9	—	16,717	16,624	916.0
Miami Division	1,290	1,332	50.9	—	27,041	26,904	1,037.7
West Palm Beach Division	376	388	28.3	—	7,872	7,821	576.9
Milwaukee–Waukesha–West Allis, WI	133	137	8.7	76	3,054	3,041	194.2
Minneapolis–St. Paul–Bloomington, MN–WI	267	277	8.0	78	6,289	6,260	182.9
Modesto, CA	20	23	4.4	100	522	530	101.7
Nashville–Davidson–Murfreesboro–Franklin, TN	234	242	13.8	47	5,078	4,974	288.0
New Haven–Milford, CT	94	98	11.4	60	3,110	3,101	359.1

Table 22. Diagnoses of HIV infection, 2013, and persons living with diagnosed HIV infection (prevalence), year-end 2012, by metropolitan statistical area of residence—United States and Puerto Rico (cont)

MSA of residence	Diagnosis, 2013				Prevalence of diagnosed HIV infection, year-end 2012		
	No.	Estimated ^a		Rank ^b	No.	Estimated ^a	
		No.	Rate			No.	Rate
New Orleans–Metairie, LA	521	538	43.4	2	8,153	8,101	659.9
New York–Newark–Jersey City, NY–NJ–PA ^c	4,143	4,941	24.8	12	143,939	144,282	727.3
Nassau County Division	237	248	8.7	—	5,900	5,826	204.7
New York Division	3,351	3,849	27.1	—	121,833	121,892	864.0
Newark Division	525	812	32.5	—	14,917	15,291	614.8
North Port–Sarasota–Bradenton, FL	90	94	12.8	53	1,664	1,656	230.0
Ogden–Clearfield, UT	8	8	1.3	104	289	288	47.0
Oklahoma City, OK	170	229	17.4	33	2,559	2,596	200.1
Omaha–Council Bluffs, NE–IA	55	57	6.3	90	1,280	1,275	144.0
Orlando–Kissimmee–Sanford, FL	584	604	26.6	10	10,145	10,093	453.9
Oxnard–Thousand Oaks–Ventura, CA	52	61	7.2	86	869	883	105.8
Palm Bay–Melbourne–Titusville, FL	65	67	12.2	56	1,212	1,202	219.6
Philadelphia–Camden–Wilmington, PA–NJ–DE–MD	1,098	1,225	20.3	22	26,151	25,920	430.6
Camden Division	143	224	17.9	—	3,075	3,169	252.5
Montgomery County Division	126	131	6.7	—	2,207	2,164	111.5
Philadelphia Division	738	768	36.3	—	18,423	18,151	860.5
Wilmington Division	91	102	14.2	—	2,446	2,436	341.4
Phoenix–Mesa–Scottsdale, AZ	532	550	12.5	55	9,744	9,696	224.0
Pittsburgh, PA	143	148	6.3	91	3,000	2,935	124.3
Portland–South Portland, ME	19	20	3.8	102	651	652	125.8
Portland–Vancouver–Hillsboro, OR–WA	183	189	8.2	77	4,319	4,299	187.8
Providence–Warwick, RI–MA	127	155	9.7	71	3,144	3,198	199.7
Provo–Orem, UT	3	3	0.5	105	149	149	27.0
Raleigh, NC	215	222	18.3	28	3,348	3,340	281.0
Richmond, VA	181	188	15.1	43	4,533	4,510	365.8
Riverside–San Bernardino–Ontario, CA	431	500	11.4	61	7,949	8,085	186.2
Rochester, NY	106	109	10.1	68	2,943	2,896	267.5
Sacramento–Roseville–Arden–Arcade, CA	193	223	10.1	69	3,700	3,763	171.5
St. Louis, MO–IL	312	331	11.8	57	6,793	6,784	242.6
Salt Lake City, UT	67	70	6.1	92	1,898	1,894	168.5
San Antonio–New Braunfels, TX	410	477	20.9	19	5,163	5,289	236.7
San Diego–Carlsbad, CA	474	551	17.2	34	12,242	12,427	391.3
San Francisco–Oakland–Hayward, CA ^c	751	820	18.2	29	24,498	24,556	551.3
Oakland Division	293	340	12.7	—	7,221	7,342	278.9
San Francisco Division	444	464	29.3	—	16,396	16,327	1042.5
San Jose–Sunnyvale–Santa Clara, CA	129	150	7.8	81	3,221	3,273	172.9
San Juan–Carolina–Caguas, PR	448	561	24.5	14	13,112	13,169	570.9
Scranton–Wilkes-Barre–Hazleton, PA	29	30	5.3	96	585	572	101.4

Table 22. Diagnoses of HIV infection, 2013, and persons living with diagnosed HIV infection (prevalence), year-end 2012, by metropolitan statistical area of residence—United States and Puerto Rico (cont)

MSA of residence	Diagnosis, 2013				Prevalence of diagnosed HIV infection, year-end 2012		
	No.	Estimated ^a		Rank ^b	No.	Estimated ^a	
		No.	Rate			No.	Rate
Seattle–Tacoma–Bellevue, WA	349	361	10.0	70	8,618	8,577	241.4
Seattle Division	291	301	10.8	—	7,607	7,571	276.3
Tacoma Division	58	60	7.3	—	1,011	1,005	123.8
Spokane–Spokane Valley, WA	22	23	4.2	101	431	428	80.4
Springfield, MA	65	98	15.7	41	1,847	1,946	311.0
Stockton–Lodi, CA	80	92	13.1	51	1,182	1,203	171.6
Syracuse, NY	57	59	8.9	75	1,092	1,071	162.0
Tampa–St. Petersburg–Clearwater, FL	597	617	21.5	18	10,387	10,321	362.8
Toledo, OH	47	48	8.0	79	955	952	156.4
Tucson, AZ	109	113	11.3	62	1,993	1,980	199.5
Tulsa, OK	113	153	15.9	38	1,601	1,623	170.5
Virginia Beach–Norfolk–Newport News, VA–NC	332	343	20.1	23	6,638	6,602	388.7
Washington–Arlington–Alexandria, DC–VA–MD–WV	1,482	1,995	33.5	6	32,649	33,595	573.0
Silver Spring Division	232	361	28.7	—	3,966	4,130	332.0
Washington Division	1,250	1,634	34.8	—	28,683	29,465	638.0
Wichita, KS	32	33	5.2	97	807	804	126.4
Winston-Salem, NC	87	89	13.7	48	1,696	1,681	259.8
Worcester, MA–CT	44	64	6.9	88	1,810	1,889	204.7
Youngstown–Warren–Boardman, OH–PA	39	40	7.2	85	675	671	120.1
Subtotal for MSAs (population of ≥500,000)	34,288	39,338	18.3	—	772,703	776,822	365.4
Metropolitan areas (population of 50,000–499,999)	4,937	5,564	9.5	—	97,141	97,218	166.1
Nonmetropolitan areas	2,505	2,809	6.1	—	51,105	51,040	110.0
Total^d	41,994	48,114	15.0	—	928,953	933,295	293.9

Note. Data include persons with a diagnosis of HIV infection regardless of stage of disease at diagnosis. Because of the lack of U.S. information for all U.S. dependent areas, table includes data for only the United States and Puerto Rico.

MSA, metropolitan statistical area.

MSA definitions for this report can be found at <http://www.census.gov/population/metro/>.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting. Rates are per 100,000 population.

^b Based on estimated rate.

^c Counts of diagnoses of HIV infection for the metropolitan divisions do not sum to the MSA total. MSA total includes data from 1 metropolitan division with population of <500,000.

^d Includes persons whose county of residence is unknown. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Table 23. Stage 3 (AIDS), 2013 and cumulative, and persons living with diagnosed HIV infection ever classified as stage 3 (AIDS) (prevalence), year-end 2012, by metropolitan statistical area of residence—United States and Puerto Rico

MSA of residence	Diagnosis, 2013				Diagnosis, cumulative ^a			Prevalence of stage 3 (AIDS), year-end 2012		
	No.	Estimated ^b		Rank ^d	Adults or adolescents No.	Children No.	Total No.	No.	Estimated ^b	
		No.	Rate ^c						No.	Rate ^c
	No.	No.	Rate ^c	Rank ^d	No.	No.	No.	No.	No.	Rate ^c
Akron, OH	13	15	2.1	101	905	1	906	401	403	57.1
Albany–Schenectady–Troy, NY	41	44	5.0	78	2,637	26	2,663	1,167	1,139	130.1
Albuquerque, NM	37	41	4.5	81	1,798	3	1,801	799	799	88.8
Allentown–Bethlehem–Easton, PA–NJ	47	50	6.1	60	1,723	20	1,743	846	828	100.1
Atlanta–Sandy Springs–Roswell, GA	837	1,020	18.5	7	31,680	148	31,828	14,568	14,348	263.1
Augusta–Richmond County, GA–SC	52	64	11.0	24	2,548	23	2,571	1,092	1,076	186.7
Austin–Round Rock, TX	136	150	8.0	40	6,090	26	6,116	2,845	2,871	156.5
Bakersfield, CA	44	49	5.7	66	2,086	9	2,095	1,218	1,227	143.5
Baltimore–Columbia–Towson, MD	468	618	22.3	6	25,666	222	25,888	10,050	10,024	364.0
Baton Rouge, LA	198	208	25.4	3	5,373	24	5,397	2,432	2,436	299.0
Birmingham–Hoover, AL	149	188	16.5	11	3,311	25	3,336	1,325	1,327	117.0
Boise City, ID	8	9	1.4	103	413	—	413	213	213	33.3
Boston–Cambridge–Newton, MA–NH ^e	200	264	5.6	67	16,903	154	17,057	7,603	7,961	171.5
Boston Division	124	164	8.4	—	10,486	93	10,579	4,537	4,749	246.5
Cambridge Division	73	97	4.2	—	6,043	60	6,103	2,916	3,062	133.5
Bridgeport–Stamford–Norwalk, CT	55	60	6.3	56	4,279	58	4,337	1,838	1,846	197.8
Buffalo–Cheektowaga–Niagara Falls, NY	64	67	5.9	64	2,958	21	2,979	1,228	1,192	105.2
Cape Coral–Fort Myers, FL	75	79	12.0	21	2,207	25	2,232	971	969	150.3
Charleston–North Charleston, SC	81	87	12.3	19	2,652	24	2,676	1,222	1,227	175.9
Charlotte–Concord–Gastonia, NC–SC	334	357	15.3	12	5,605	27	5,632	2,738	2,755	120.1
Chattanooga, TN–GA	32	36	6.6	51	1,291	3	1,294	619	601	111.8
Chicago–Naperville–Elgin, IL–IN–WI	796	905	9.5	29	37,743	260	38,003	16,104	16,412	172.5
Chicago Division	717	818	11.1	—	34,576	239	34,814	14,636	14,936	204.2
Elgin Division	16	18	2.8	—	773	8	781	367	371	59.2
Gary Division	42	47	6.6	—	1,383	8	1,391	591	590	83.6
Lake County Division	21	23	2.7	—	1,012	5	1,017	510	515	59.2
Cincinnati, OH–KY–IN	100	107	5.0	77	3,668	20	3,688	1,772	1,779	83.6
Cleveland–Elyria, OH	121	129	6.3	58	5,140	49	5,189	2,319	2,325	112.6
Colorado Springs, CO	17	18	2.6	98	712	5	717	327	327	48.9
Columbia, SC	103	110	13.8	14	4,348	27	4,375	2,264	2,269	289.2

Table 23. Stage 3 (AIDS), 2013 and cumulative, and persons living with diagnosed HIV infection ever classified as stage 3 (AIDS) (prevalence), year-end 2012, by metropolitan statistical area of residence—United States and Puerto Rico (cont)

MSA of residence	Diagnosis, 2013				Diagnosis, cumulative ^a			Prevalence of stage 3 (AIDS), year-end 2012		
	No.	Estimated ^b		Rank ^d	Adults or adolescents No.	Children No.	Total No.	No.	Estimated ^b	
		No.	Rate ^c						No.	No.
	No.	No.	Rate ^c	Rank ^d	No.	No.	No.	No.	No.	Rate ^c
Columbus, OH	157	167	8.5	33	4,515	22	4,537	2,082	2,093	107.6
Dallas–Fort Worth–Arlington, TX	712	805	11.8	23	26,356	63	26,419	11,958	12,228	182.4
Dallas Division	560	638	14.2	—	20,878	37	20,915	9,532	9,778	220.9
Fort Worth Division	152	166	7.2	—	5,478	26	5,504	2,426	2,450	107.7
Dayton, OH	40	43	5.3	71	1,547	14	1,561	739	742	92.4
Deltona–Daytona Beach–Ormond Beach, FL	61	64	10.7	25	2,023	19	2,042	879	876	147.2
Denver–Aurora–Lakewood, CO	133	142	5.2	74	8,387	23	8,410	3,869	3,872	146.3
Des Moines–West Des Moines, IA	31	33	5.5	68	731	4	735	360	360	61.1
Detroit–Warren–Dearborn, MI	322	349	8.1	36	13,079	77	13,156	5,254	5,131	119.5
Detroit Division	237	257	14.5	—	10,256	60	10,316	3,910	3,804	212.2
Warren Division	85	92	3.7	—	2,823	17	2,840	1,344	1,326	53.1
Durham–Chapel Hill, NC	27	28	5.3	73	1,522	10	1,532	666	669	127.3
El Paso, TX	77	81	9.7	28	2,033	10	2,043	1,062	1,071	128.8
Fresno, CA	57	62	6.5	54	2,079	11	2,090	989	997	105.3
Grand Rapids–Wyoming, MI	40	44	4.4	83	1,174	6	1,180	567	559	55.6
Greensboro–High Point, NC	54	57	7.7	41	1,728	19	1,747	786	789	107.2
Greenville–Anderson–Mauldin, SC	57	60	7.1	45	1,965	4	1,969	908	910	108.1
Harrisburg–Carlisle, PA	34	38	6.8	50	1,586	9	1,595	760	736	132.7
Hartford–West Hartford–East Hartford, CT	83	92	7.6	42	5,874	46	5,920	2,349	2,362	194.5
Honolulu (Urban), HI	37	43	4.4	84	2,366	13	2,379	999	1,004	102.9
Houston–The Woodlands–Sugar Land, TX	842	928	14.7	13	33,360	172	33,532	13,821	14,017	227.0
Indianapolis–Carmel–Anderson, IN	125	133	6.8	49	5,149	25	5,174	2,472	2,478	128.5
Jackson, MS	161	170	29.6	2	3,390	30	3,420	1,546	1,526	264.8
Jacksonville, FL	234	246	17.6	9	7,928	78	8,006	3,586	3,575	259.4
Kansas City, MO–KS	123	135	6.6	52	5,901	15	5,916	2,736	2,730	133.9
Knoxville, TN	41	46	5.4	69	1,306	7	1,313	623	605	71.4
Lakeland–Winter Haven, FL	76	81	13.0	17	2,438	22	2,460	1,094	1,090	176.9
Lancaster, PA	14	15	2.8	96	880	22	902	406	396	75.2
Las Vegas–Henderson–Paradise, NV	224	242	11.9	22	6,223	28	6,251	3,028	3,020	151.2
Little Rock–North Little Rock–Conway, AR	69	77	10.6	26	1,797	12	1,809	866	874	121.8

Table 23. Stage 3 (AIDS), 2013 and cumulative, and persons living with diagnosed HIV infection ever classified as stage 3 (AIDS) (prevalence), year-end 2012, by metropolitan statistical area of residence—United States and Puerto Rico (cont)

MSA of residence	Diagnosis, 2013				Diagnosis, cumulative ^a			Prevalence of stage 3 (AIDS), year-end 2012		
	No.	Estimated ^b		Rank ^d	Adults or adolescents	Children	Total	No.	Estimated ^b	
		No.	Rate ^c						No.	No.
	No.	No.	Rate ^c	Rank ^d	No.	No.	No.	No.	No.	Rate ^c
Los Angeles–Long Beach–Anaheim, CA	1,064	1,194	9.1	31	70,713	306	71,019	30,522	31,043	238.1
Anaheim Division	144	160	5.1	—	8,334	44	8,378	3,870	3,933	127.5
Los Angeles Division	920	1,034	10.3	—	62,379	262	62,641	26,652	27,111	272.4
Louisville/Jefferson County, KY–IN	93	103	8.2	35	3,130	27	3,157	1,476	1,469	117.4
Madison, WI	14	15	2.4	99	663	5	668	311	312	50.2
McAllen–Edinburg–Mission, TX	63	69	8.4	34	1,000	12	1,012	557	566	70.3
Memphis, TN–MS–AR	360	420	31.3	1	7,185	19	7,204	3,303	3,224	240.5
Miami–Fort Lauderdale–West Palm Beach, FL	1,362	1,435	24.6	4	67,716	1,002	68,718	28,149	28,058	486.8
Fort Lauderdale Division	443	471	25.6	—	20,737	264	21,002	9,135	9,109	501.9
Miami Division	685	718	27.4	—	35,416	512	35,928	14,152	14,104	544.0
West Palm Beach Division	234	247	18.0	—	11,563	225	11,788	4,862	4,845	357.3
Milwaukee–Waukesha–West Allis, WI	75	78	5.0	76	3,171	20	3,191	1,473	1,476	94.3
Minneapolis–St. Paul–Bloomington, MN–WI	134	142	4.1	87	5,649	24	5,673	2,846	2,847	83.2
Modesto, CA	10	11	2.0	102	827	6	833	388	389	74.6
Nashville–Davidson–Murfreesboro–Franklin, TN	132	149	8.5	32	5,040	23	5,063	2,553	2,488	144.1
New Haven–Milford, CT	64	69	8.0	39	5,162	74	5,236	2,178	2,188	253.3
New Orleans–Metairie, LA	271	286	23.0	5	10,955	75	11,030	4,326	4,320	351.9
New York–Newark–Jersey City, NY–NJ–PA ^e	2,386	2,648	13.3	15	228,961	3,028	231,989	85,111	86,181	434.4
Nassau County Division	123	129	4.5	—	9,332	117	9,449	3,568	3,479	122.2
New York Division	1,948	2,160	15.2	—	193,508	2,539	196,047	73,009	74,055	524.9
Newark Division	298	340	13.6	—	24,093	360	24,453	7,772	7,901	317.7
North Port–Sarasota–Bradenton, FL	52	55	7.5	43	2,407	29	2,436	1,041	1,039	144.3
Ogden–Clearfield, UT	5	5	0.9	104	316	4	320	165	166	27.0
Oklahoma City, OK	88	98	7.4	44	2,907	5	2,912	1,205	1,218	93.9
Omaha–Council Bluffs, NE–IA	33	39	4.4	82	1,360	5	1,365	691	691	78.1
Orlando–Kissimmee–Sanford, FL	357	381	16.8	10	11,418	95	11,513	5,443	5,430	244.2
Oxnard–Thousand Oaks–Ventura, CA	23	26	3.1	95	1,229	5	1,234	538	539	64.6
Palm Bay–Melbourne–Titusville, FL	37	38	7.0	46	1,807	11	1,818	806	802	146.6

Table 23. Stage 3 (AIDS), 2013 and cumulative, and persons living with diagnosed HIV infection ever classified as stage 3 (AIDS) (prevalence), year-end 2012, by metropolitan statistical area of residence—United States and Puerto Rico (cont)

MSA of residence	Diagnosis, 2013				Diagnosis, cumulative ^a			Prevalence of stage 3 (AIDS), year-end 2012		
	No.	Estimated ^b		Rank ^d	Adults or adolescents	Children	Total	No.	Estimated ^b	
		No.	Rate ^c						No.	No.
	No.	No.	Rate ^c	Rank ^d	No.	No.	No.	No.	No.	Rate ^c
Philadelphia–Camden–Wilmington, PA–NJ–DE–MD	621	726	12.0	20	34,504	315	34,819	15,160	15,032	249.7
Camden Division	76	88	7.0	—	3,855	42	3,897	1,554	1,581	126.0
Montgomery County Division	79	88	4.5	—	2,800	11	2,811	1,148	1,112	57.3
Philadelphia Division	391	466	22.0	—	24,206	240	24,446	10,928	10,806	512.3
Wilmington Division	75	85	11.8	—	3,643	22	3,665	1,530	1,533	214.8
Phoenix–Mesa–Scottsdale, AZ	255	269	6.1	59	9,808	32	9,840	4,629	4,626	106.9
Pittsburgh, PA	90	99	4.2	86	3,912	21	3,933	1,664	1,615	68.4
Portland–South Portland, ME	16	20	3.9	90	722	1	723	353	358	69.2
Portland–Vancouver–Hillsboro, OR–WA	99	108	4.7	79	5,838	9	5,847	2,615	2,625	114.7
Providence–Warwick, RI–MA	86	102	6.4	55	4,763	42	4,805	2,113	2,163	135.1
Provo–Orem, UT	2	2	0.4	105	148	3	151	85	85	15.5
Raleigh, NC	93	98	8.1	37	2,925	16	2,941	1,628	1,636	137.7
Richmond, VA	78	85	6.8	48	4,337	34	4,371	2,021	2,036	165.1
Riverside–San Bernardino–Ontario, CA	233	265	6.0	61	10,854	59	10,913	5,136	5,226	120.3
Rochester, NY	55	58	5.3	72	3,690	17	3,707	1,664	1,623	150.0
Sacramento–Roseville–Arden–Arcade, CA	91	102	4.6	80	4,925	28	4,953	2,157	2,170	98.9
St. Louis, MO–IL	158	168	6.0	62	7,527	43	7,570	3,426	3,421	122.3
Salt Lake City, UT	34	37	3.2	93	1,986	9	1,995	1,068	1,069	95.1
San Antonio–New Braunfels, TX	173	183	8.0	38	6,631	31	6,663	3,040	3,069	137.3
San Diego–Carlsbad, CA	267	296	9.2	30	15,463	69	15,532	7,199	7,316	230.3
San Francisco–Oakland–Hayward, CA ^e	393	442	9.8	27	45,839	99	45,939	16,216	16,327	366.6
Oakland Division	180	205	7.7	—	11,925	51	11,976	5,054	5,150	195.6
San Francisco Division	203	226	14.3	—	32,212	44	32,256	10,550	10,567	674.7
San Jose–Sunnyvale–Santa Clara, CA	69	75	3.9	89	4,721	15	4,736	2,271	2,284	120.7
San Juan–Carolina–Caguas, PR	221	290	12.7	18	24,606	278	24,884	7,519	7,254	314.5
Scranton–Wilkes-Barre–Hazleton, PA	11	13	2.3	100	693	7	700	321	310	55.0
Seattle–Tacoma–Bellevue, WA	169	184	5.1	75	10,890	29	10,919	4,865	4,876	137.3
Seattle Division	154	167	6.0	—	9,633	19	9,652	4,309	4,319	157.6
Tacoma Division	15	17	2.0	—	1,257	10	1,267	556	557	68.6
Spokane–Spokane Valley, WA	15	17	3.2	94	650	1	651	269	270	50.7
Springfield, MA	27	36	5.7	65	2,481	28	2,509	1,001	1,053	168.3

Table 23. Stage 3 (AIDS), 2013 and cumulative, and persons living with diagnosed HIV infection ever classified as stage 3 (AIDS) (prevalence), year-end 2012, by metropolitan statistical area of residence—United States and Puerto Rico (cont)

MSA of residence	Diagnosis, cumulative ^a										
	Diagnosis, 2013				Adults or adolescents			Children			Total
	No.	Estimated ^b		Rank ^d	No.	Estimated ^b		No.	Prevalence of stage 3 (AIDS), year-end 2012		
		No.	Rate ^c			No.	No.		No.	Estimated ^b	
								No.	No.	Rate ^c	
Stockton–Lodi, CA	43	48	6.9	47	1,433	14	1,447	682	686	97.9	
Syracuse, NY	27	28	4.2	85	1,555	9	1,564	669	651	98.5	
Tampa–St. Petersburg–Clearwater, FL	360	377	13.1	16	14,129	115	14,244	6,080	6,062	213.1	
Toledo, OH	20	22	3.6	91	1,026	14	1,040	467	469	77.0	
Tucson, AZ	49	53	5.4	70	2,449	10	2,459	1,046	1,044	105.2	
Tulsa, OK	52	57	5.9	63	1,928	10	1,938	824	831	87.3	
Virginia Beach–Norfolk–Newport News, VA–NC	102	112	6.6	53	5,849	64	5,914	2,590	2,600	153.1	
Washington–Arlington–Alexandria, DC–VA–MD–WV	879	1,098	18.5	8	40,026	324	40,351	18,251	18,523	316.0	
Silver Spring Division	172	235	18.7	—	4,020	28	4,048	2,091	2,130	171.3	
Washington Division	707	863	18.4	—	36,006	296	36,303	16,160	16,393	354.9	
Wichita, KS	24	25	3.9	88	1,023	2	1,025	453	453	71.2	
Winston-Salem, NC	39	41	6.3	57	1,286	9	1,295	620	624	96.4	
Worcester, MA–CT	26	33	3.5	92	2,503	22	2,525	1,155	1,203	130.3	
Youngstown–Warren–Boardman, OH–PA	14	15	2.7	97	710	—	710	315	315	56.4	
Subtotal for MSAs (population of ≥500,000)	19,353	21,702	10.1	—	1,030,521	8,518	1,039,039	433,020	435,421	204.8	
Metropolitan areas (population of 50,000–499,999)	2,992	3,349	5.7	—	119,509	828	120,338	52,711	52,556	89.8	
Nonmetropolitan areas	1,662	1,846	4.0	—	62,793	426	63,219	28,181	28,103	60.6	
Total^f	24,181	27,121	8.5	—	1,218,997	9,824	1,228,821	516,934	519,130	163.5	

Note. Because of the lack of U.S. census information for all U.S. dependent areas, table includes data for only the United States and Puerto Rico.

MSA, metropolitan statistical area.

MSA definitions for this report can be found at <http://www.census.gov/population/metro/>.

Reported numbers less than 12, as well as estimated numbers (and accompanying rates and trends) based on these numbers, should be interpreted with caution because the numbers have underlying relative standard errors greater than 30% and are considered unreliable.

^a From the beginning of the epidemic through 2013.

^b Estimated numbers resulted from statistical adjustment that accounted for reporting delays, but not for incomplete reporting.

^c Rates are per 100,000 population.

^d Based on estimated rate.

^e Counts of stage 3 (AIDS) classifications for the metropolitan divisions do not sum to the MSA total. MSA total includes data from 1 metropolitan division with population of <500,000.

^f Includes persons whose county of residence is unknown. Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Web Addresses for Reports of State and Local HIV Surveillance

Alabama	http://www.adph.org/aids/Default.asp?id=984
Alaska	http://www.epi.hss.state.ak.us/bulletins/catlist.jsp?cattype=HIV+(Human+Immunodeficiency+Virus)
Arizona	http://www.azdhs.gov/phs/edc/odis/hiv-epidemiology/reports/index.php
Arkansas	http://www.health.arkansas.gov/programsServices/healthStatistics/Pages/stdHIVRegistry.aspx
California	http://www.cdph.ca.gov/data/statistics/Pages/OAHIVAIDSStatistics.aspx
Chicago	http://www.cityofchicago.org/HealthInformation
Colorado	http://www.colorado.gov/pacific/cdphe/sti-and-hiv-data-and-trends
Connecticut	http://www.ct.gov/dph/cwp/view.asp?a=3135&q=393048
Delaware	http://www.dhss.delaware.gov/dhss/dph/epi/disstatshiv.html
District of Columbia	http://doh.dc.gov/service/hiv-aids-reports-and-publications
Florida	http://www.floridahealth.gov/diseases-and-conditions/aids/surveillance/index.html
Georgia	http://dph.georgia.gov/georgias-hiv-aids-epidemiology-surveillance-section
Guam	http://dphss.guam.gov/content/hiv-surveillance-program
Hawaii	http://health.hawaii.gov/std-aids/hiv-aids/data-and-statistics/
Houston	http://www.houstontx.gov/health/HIV-STD/
Idaho	http://healthandwelfare.idaho.gov/Health/FamilyPlanningSTDHIV/STDStatistics/tabid/393/Default.aspx
Illinois	http://www.idph.state.il.us/aids/stats.htm
Indiana	http://www.in.gov/isdh/23266.htm
Iowa	http://www.idph.state.ia.us/HivStdHep/HIV-AIDS.aspx?prog=Hiv&pg=HivSurv
Kansas	http://www.kdheks.gov/sti_hiv/hiv_std_update_newsletter.html
Kentucky	http://chfs.ky.gov/dph/epi/hiv-aids/surveillance.htm
Los Angeles	http://publichealth.lacounty.gov/phcommon/public/reports/rptspubdisplay.cfm?unit=hiv&ou=ph&prog=hae
Louisiana	http://new.dhh.louisiana.gov/index.cfm/newsroom/category/63
Maine	http://www.maine.gov/dhhs/mecdc/infectious-disease/hiv-std/data/hiv.shtml
Maryland	http://ideha.dhmh.md.gov/OIDEOR/CHSE/SitePages/statistics.aspx
Massachusetts	http://www.mass.gov/eohhs/gov/departments/dph/programs/id/hiv-aids-surveillance/surveillance/public-health-cdc-hiv-aids-surveillance-std-report.html
Michigan	http://michigan.gov/mdch/0,4612,7-132-2944_5320-36307--,00.html
Minnesota	http://www.health.state.mn.us/divs/idepc/diseases/hiv/hivstatistics.html
Mississippi	http://www.msdc.state.ms.us/msdcsite/index.cfm/14,0,150,134,html
Missouri	http://www.dhss.mo.gov/data/hivstdaids/index.php
Montana	http://hivdata.hhs.mt.gov/
Nebraska	http://dhhs.ne.gov/publichealth/Pages/dpc_hiv.aspx
Nevada	http://health.nv.gov/FP_Publications.htm
New Hampshire	http://www.dhhs.nh.gov/data/index.htm
New Jersey	http://www.state.nj.us/health/aids/aidsqr.shtml
New Mexico	http://archive.nmhealth.org/erd/healthdata/hiv_aids.shtml
New York	http://www.health.state.ny.us/diseases/aids/statistics/index.htm
New York City	http://www.nyc.gov/html/doh/html/data/hivepi.shtml
North Carolina	http://epi.publichealth.nc.gov/cd/stds/figures.html
North Dakota	http://www.ndhealth.gov/hiv/
Ohio	http://www.odh.ohio.gov/healthstats/disease/hivdata/hivcov.aspx
Oklahoma	http://www.ok.gov/health/Disease,_Prevention,_Preparedness/HIV_STD_Service/HIV_STD_Statistics/index.html
Oregon	http://public.health.oregon.gov/DiseasesConditions/CommunicableDisease/DiseaseSurveillanceData/HIVData/Pages/index.aspx
Pennsylvania	http://www.portal.state.pa.us/portal/server.pt?open=514&objID=557343&mode=2
Philadelphia	http://www.phila.gov/health/AACO/AACODataResearch.html
Puerto Rico	http://www.salud.gov.pr/Programas/OficEpidemiologia/Pages/ProgramadeVigilanciadeSIDA.aspx
Rhode Island	http://www.health.ri.gov/data/hiv-aids/
San Francisco	http://www.sfdph.org/dph/comupg/oprograms/HIVepiSec/HIVepiSecReports.asp
South Carolina	http://www.dhec.sc.gov/health/disease/stdhiv/surveillance.htm
South Dakota	http://doh.sd.gov/diseases/infectious/HIV-AIDS/Prevention.aspx
Tennessee	http://health.state.tn.us/STD/data.shtml
Texas	http://www.dshs.state.tx.us/hivstd/reports/default.shtm
U.S. Virgin Islands	http://www.healthvi.com/programs/communicable-disease/index.html
Utah	http://health.utah.gov/epi/diseases/hiv-aids/surveillance/index.html
Vermont	http://healthvermont.gov/prevent/aids/quarter/reports.aspx
Virginia	http://www.vdh.virginia.gov/epidemiology/DiseasePrevention/Data/
Washington	http://www.doh.wa.gov/DataandStatisticalReports/DiseasesandChronicConditions/HIVAIDSData.aspx
West Virginia	http://www.dhhr.wv.gov/oeps/std-hiv-hep/HIV_AIDS/Pages/HIVSurveillance.aspx
Wisconsin	http://dhs.wisconsin.gov/aids-hiv/Stats/index.htm
Wyoming	http://www.health.wyo.gov/PHSD/hivsurveillance/index.html

Note. Electronic reports are not available for the following areas: American Samoa, the Northern Mariana Islands, and the Republic of Palau.