

parallel with similar efforts in Europe, strategies need to be developed to protect commercial and pet rabbits.

Tracking the spread of RHDV2 in Australia, in competition with existing field strains, highlights the value of Australia's rabbits and their diseases as a model system for emerging infectious diseases. The point releases of both myxoma virus and RHDV into large naive host populations represent a grand experiment in disease emergence and evolution (10), which provides a unique opportunity to study the virulence evolution of emerging pathogens as well as their complex interactions with each other. It is notable that since the release of RHDV in Australia in 1995, strains of 1 viral lineage dominate the viral population nationwide despite hundreds of deliberate releases of the original virus strain for local rabbit control, which strongly suggests it has a major selective advantage (7). That RHDV2 appeared in a wild rabbit is therefore remarkable, particularly because Australian field strains were spreading simultaneously in the same area. Comparing the epidemiology of this strain in Australia to the epidemiology of its well-documented spread in Europe will provide valuable insights into RHDV epidemiology relevant to both continents.

Acknowledgments

We thank Roslyn Mourant and James Biglia for assistance with sample processing.

J.E.M. is supported by grant DP140103362 from the Australian Research Council. E.C.H. is supported by a National Health and Medical Research Council Australia fellowship (AF30).

References

1. Le Gall-Reculé G, Zwingelstein F, Boucher S, Le Normand B, Plassiart G, Portejoie Y, et al. Detection of a new variant of rabbit haemorrhagic disease virus in France. *Vet Rec.* 2011;168:137–8. <http://dx.doi.org/10.1136/vr.d697>
2. Dalton KP, Nieceza I, Balseiro A, Mugerza MA, Rosell JM, Casais R, et al. Variant rabbit hemorrhagic disease virus in young rabbits, Spain. *Emerg Infect Dis.* 2012;18:2009–12. <http://dx.doi.org/10.3201/eid1812.120341>
3. Le Gall-Reculé G, Lavazza A, Marchandeu S, Bertagnoli S, Zwingelstein F, Cavadini P, et al. Emergence of a new lagovirus related to rabbit haemorrhagic disease virus. *Vet Res.* 2013;44:1–13. <http://dx.doi.org/10.1186/1297-9716-44-81>
4. Abrantes J, Lopes AM, Dalton KP, Melo P, Correia JJ, Ramada M, et al. New variant of rabbit hemorrhagic disease virus, Portugal, 2012–2013. *Emerg Infect Dis.* 2013;19:1900–2. <http://dx.doi.org/10.3201/eid1911.130908>
5. Lopes AM, Correia J, Abrantes J, Melo P, Ramada M, Magalhaes MJ, et al. Is the new variant RHDV replacing genotype 1 in Portuguese wild rabbit populations? *Viruses.* 2015;7:27–36. <http://dx.doi.org/10.3390/v7010027>
6. Elsworth P, Cooke BD, Kovaliski J, Sinclair R, Holmes EC, Strive T. Increased virulence of rabbit haemorrhagic disease virus associated with genetic resistance in wild Australian rabbits (*Oryctolagus cuniculus*). *Virology.* 2014;464–5:415–23. <http://dx.doi.org/10.1016/j.virol.2014.06.037>
7. Eden JS, Kovaliski J, Duckworth JA, Swain G, Mahar JE, Strive T, et al. Comparative phylodynamics of rabbit haemorrhagic disease virus (RHDV) in Australia and New Zealand. *J Virol.* 2015;89:9548–58. <http://dx.doi.org/10.1128/JVI.011100-15>
8. Lopes AM, Dalton KP, Magalhaes MJ, Parra F, Esteves PJ, Holmes EC, et al. Full genomic analysis of new variant rabbit hemorrhagic disease virus revealed multiple recombination events. *J Gen Virol.* 2015;96:1309–19. <http://dx.doi.org/10.1099/vir.0.000070>
9. Almeida T, Lopes AM, Magalhães MJ, Neves F, Pinheiro A, Gonçalves D, et al. Tracking the evolution of the G1/RHDVb recombinant strains introduced from the Iberian Peninsula to the Azores islands, Portugal. *Infect Genet Evol.* 2015;34:307–13. <http://dx.doi.org/10.1016/j.meegid.2015.07.010>
10. Di Giallonardo F, Holmes EC. Viral biocontrol: grand experiments in disease emergence and evolution. *Trends Microbiol.* 2015;23:83–90. <http://dx.doi.org/10.1016/j.tim.2014.10.004>

Address for correspondence: Tanja Strive, CSIRO Health and Biosecurity, Clunies Ross St, Canberra, Australian Capital Territory 2601, Australia; email: tanja.strive@csiro.au

Characteristics of Traveler with Middle East Respiratory Syndrome, China, 2015

Wen Da Guan,¹ Chris Ka Pun Mok,¹ Zi Lin Chen,¹ Li Qiang Feng, Zheng Tu Li, Ji Cheng Huang, Chang Wen Ke, Xilong Deng, Yun Ling, Shi Guan Wu, Xue Feng Niu, Ranawaka A Perera, Yuan Da Xu, Jincun Zhao, Lin Qi Zhang, Yi Min Li, Rong Chang Chen, Malik Peiris, Ling Chen, Nan Shan Zhong

Author affiliations: State Key Laboratory of Respiratory Disease, First Affiliated Hospital of Guangzhou Medical University, Guangzhou, China (W.D. Guan, Z.T. Li, S.G. Wu, X.F. Niu, Y.D. Xu, J. Zhao, Y.M. Li, R.C. Chen, L. Chen, N.S. Zhong); The University of Hong Kong, Hong Kong, China (C.K.P. Mok, R.A. Perera, M. Peiris); Huizhou Municipal Central Hospital, Huizhou, China (Z.L. Chen, Y. Ling); Guanzhou Institute of Biomedicine and Health, Guangzhou (L.Q. Feng, L. Chen); Guangdong Inspection and Quarantine Technology Center, Guangzhou (J.C. Huang); Guangdong Center for Disease Control and Prevention, Guangzhou (C.W. Ke); Guangzhou Eighth People's Hospital, Guangzhou (X. Deng); Tsinghua University School of Medicine, Beijing, China (L.Q. Zhang)

DOI: <http://dx.doi.org/10.3201/eid2112.151232>

To the Editor: A traveler returning from the Middle East initiated an outbreak of Middle East respiratory syndrome (MERS) in South Korea in 2015, which resulted in 186 cases and 36 deaths (1–3). We report a case of

¹These authors contributed equally to this article.

MERS in a 43-year-old man from South Korea who acquired this disease during this outbreak (online Technical Appendix Figure 1, panel A, <http://wwwnc.cdc.gov/EID/article/21/12/15-1232-Techapp1.pdf>) (4).

The National Health and Family Planning Commission of China determined that collection of data for this patient was part of a public health investigation of an emerging outbreak. Therefore, informed consent was not required. This study was approved by the ethical committee of the First Affiliated Hospital of Guangzhou Medical University.

The patient had been receiving thiamazole for 7 years for hyperthyroidism. He had contact with the index case-patient during the outbreak in South Korea on May 16, 2015. On May 25, the patient traveled to Hong Kong and then to Huizhou, China. He was hospitalized in China on May 28 (day 7 of illness). At admission, he had a high fever (temperature 39.5°C) and a dry cough. Chest radiography on day 7 showed mild bilateral ground glass opacities in the lower lung (online Technical Appendix Figure 1, panel B).

The patient was given oseltamivir (150 mg, 2×/day for 2 days) until identified as being infected with Middle East respiratory syndrome coronavirus (MERS-CoV) on day 8 by real-time reverse transcription PCR. He was given ribavirin (2.0 mg on day 8; 0.6 mg 3×/d on days 9–16; and 0.6 mg 2×/d on days 17–19) and 135 µg of peginterferon α -2a by intravenous injection on day 8 (online Technical Appendix Table 2). Thrombocytopenia and a decrease in the hemoglobin level developed, which might have been related to use of ribavirin (online Technical Appendix Table 1).

Chest radiography on June 1 (day 11) showed increased bilateral consolidation of the patient's lower lung (online Technical Appendix Figure 1, panel C). He was given intravenous immunoglobulin, antimicrobial drugs, and thymosin α 1. His body temperature returned to normal on day 14 (online Technical Appendix Figure 2). Chest radiography on day 35 showed resolution of bilateral lung infiltrations (online Technical Appendix Figure 1, panel D). He was discharged on day 36.

Viral RNA was detected in sputum and fecal specimens up to day 26 of illness. Virus load in sputum specimens collected on days 11–15 were lower than in specimens obtained on days 16–18 (online Technical Appendix Figure 3, panel A). Swab samples collected on days 13 and 15 from the patient's palm, mobile telephone, blanket, and bed railings, and from his hospital room floor were negative for viral RNA.

Concentrations of proinflammatory cytokines and chemokines (interferon- α , interferon-inducible protein 10, monocyte chemoattractant protein-1, interleukin 6 [IL-6], IL-10, tumor necrosis factor- α , IL-8, macrophage inflammatory protein- α [MIP-1 α], MIP-1 β , and IL-1 β) were determined for serial serum samples. Interferon- α , interferon-inducible protein 10, monokine induced by interferon- γ , IL-6, monocyte

chemoattractant protein-1, and IL-8 were detected on day 11 of illness but levels decreased as the patient clinically improved (online Technical Appendix Figure 3, panel B).

The peginterferon α 2 the patient was given on day 8 might have influenced his plasma interferon- α levels (6). However, a previous study also showed increased levels of interferon- α in a patient who survived MERS-CoV infection but not in a person who died of MERS (7). Although MERS-CoV evades induction of innate immune responses by cell types, the virus elicits interferon responses in plasmacytoid dendritic cells *in vitro* (8). Levels of tumor necrosis factor- α , MIP-1 α , MIP-1 β , IL-10, and IL-1 β did not increase in any of these specimens.

Peripheral blood mononuclear cells (PBMCs) obtained on day 24 of illness showed a strong specific T-cell response against MERS-CoV spike protein but not against severe acute respiratory syndrome coronavirus (SARS-CoV) spike protein (online Technical Appendix Figure 3, panel C). PBMCs from persons who were infected with SARS-CoV in 2003, as well as healthy persons, showed low-level T-cell responses against MERS-CoV spike protein, although some persons with a history of SARS still had detectable responses to SARS-CoV spike protein. It was reported that T-cell responses to SARS-CoV were directed against spike and nucleocapsid proteins (9). We did not have sufficient PBMCs to test T-cell responses against nucleocapsid protein.

Results for MERS-CoV antibody were negative at day 11 of illness by MERS-CoV spike pseudotype assay (MERS-S ppNT), microneutralization, 50% plaque reduction neutralization test (PRNT₅₀), and S1 ELISA (EUROIMMUN AG, Lübeck, Germany). The patient showed seroconversion by day 14. MERS-S ppNT and PRNT₅₀ provided earlier evidence of seroconversion (day 15) and higher antibody titers than the microneutralization, (day 18) (online Technical Appendix Figure 3, panel D). Potent T-cell responses were elicited to MERS-CoV spike protein. These responses did not show cross-reactivity with SARS-CoV spike protein.

The MERS-S ppNT, which does not require Biosafety Level 3 containment, had sensitivity equivalent with that of PRNT₅₀, which requires containment. Thus, MERS-S ppNT is a sensitive and specific assay for detecting neutralizing antibody against MERS-CoV. The sensitivity and specificity of this assay have been well-documented with serum samples from dromedary camels and other animals (10).

This study was supported by the Science Research Project of the Guangdong Province (2013B020224006); the Municipal Science and Technology Bureau Foundation of Guangzhou (2014Y2-00031); the National Key Project of Clinical Faculty and Facility Construction on Infectious Diseases (2013–2014); and the National Institute of Allergy and Infectious Diseases, National Institutes of Health (HHSN272201400006C).

References


- Cowling BJ, Park M, Fang VJ, Wu P, Leung GM, Wu JT. Preliminary epidemiological assessment of MERS-CoV outbreak in South Korea, May to June 2015. *Euro Surveill.* 2015;20:pii: 21163.
- Park HY, Lee EJ, Ryu YW, Kim Y, Kim H, Lee H, et al. Epidemiological investigation of MERS-CoV spread in a single hospital in South Korea, May to June 2015. *Euro Surveill.* 2015; 20:pii: 21169.
- World Health Organization. Middle East respiratory syndrome coronaviruses (MERS-CoV) [cited 2015 Sep 5]. <http://www.who.int/emergencies/mers-cov/en/>
- Wu J, Yi L, Zou L, Zhong H, Liang L, Song T, et al. Imported case of MERS-CoV infection identified in China, May 2015: detection and lesson learned. *Euro Surveill.* 2015;20:pii:21158.
- Memish ZA, Al-Tawfiq JA, Makhdoom HQ, Assiri A, Alhakeem RF, Albarrak A, et al. Respiratory tract samples, viral load, and genome fraction yield in patients with Middle East respiratory syndrome. *J Infect Dis.* 2014;210:1590–4. <http://dx.doi.org/10.1093/infdis/jiu292>
- El Sabaawy D, El-Haggar S, El-Bahrawy, Waked I, El-Said H. A comparative study of variants of pegylated interferon alpha in treatment of chronic HCV patients. *APMIS.* 2015;123:482–9. <http://dx.doi.org/10.1111/apm.12377>
- Faure E, Poissy J, Goffard A, Fournier C, Kipnis E, Titecat M, et al. Distinct immune response in two MERS-CoV-infected patients: can we go from bench to bedside? *PLoS ONE.* 2014;9:e88716. <http://dx.doi.org/10.1371/journal.pone.0088716>
- Scheuplein VA, Seifried J, Malczyk AH, Miller L, Höcker L, Vergara-Alert J, et al. High secretion of interferons by human plasmacytoid dendritic cells upon recognition of Middle East respiratory syndrome coronavirus. *J Virol.* 2015;89:3859–69. <http://dx.doi.org/10.1128/JVI.03607-14>
- Channappanavar R, Zhao J, Perlman S. T cell-mediated immune response to respiratory coronaviruses. *Immunol Res.* 2014;59:118–28. <http://dx.doi.org/10.1007/s12026-014-8534-z>
- Perera RA, Wang P, Gomaa MR, El-Shesheny R, Kandeil A, Bagato O, et al. Seroepidemiology for MERS coronavirus using microneutralisation and pseudoparticle virus neutralisation assays reveal a high prevalence of antibody in dromedary camels in Egypt, June 2013. *Euro Surveill.* 2013;18:pii=20574.

Address for correspondence: Ling Chen, State Key Laboratory of Respiratory Disease, National Clinical Research Center for Respiratory Disease, First Affiliated Hospital of Guangzhou Medical University, Guangzhou, China; email: chen002@gmail.com

March 2014: Tuberculosis

Including:

- Invasive Fungal Infections after Natural Disasters
- Monitoring Water Sources for Environmental Reservoirs of Toxigenic *Vibrio cholerae* O1, Haiti
- High Level Relatedness among *Mycobacterium abscessus* subsp. massiliense Strains from Widely Separated Outbreaks
- Hendra Virus Vaccine, a One Health Approach to Protecting Horse, Human, and Environmental Health
- Possible Role of Songbirds and Parakeets in Transmission of Influenza A(H7N9) Virus to Humans
- Hantavirus Infections among Overnight Visitors to Yosemite National Park, California, USA, 2012
- Use of Drug-Susceptibility Testing for Management of Drug-Resistant Tuberculosis, Thailand, 2004–2008
- Comparison of Imported *Plasmodium ovale curtisi* and *P. ovale wallikeri* Infections among Patients in Spain, 2005–2011



<http://wwwnc.cdc.gov/eid/articles/issue/20/3/table-of-contents>

Characteristics of Traveler with Middle East Respiratory Syndrome, China, 2015

Technical Appendix

Methods

Detection of Virus in Samples

Throat swab, nasal swab, sputum, urine, serum, fecal, and environmental samples were collected from the patient and placed in viral transport medium. Viral RNA was extracted from samples by using the QIAamp MinElute Virus Spin Kit (QIAGEN, Valencia, CA, USA) according to the manufacturer's instructions. Middle East respiratory syndrome coronavirus (MERS-CoV) was detected by using real-time reverse transcription PCR with primers upE-Fwd (5'-GCAACGCGCGATTTCAGTT-3') and upE-Rev (5'-GCCTCTACACGGGACCCATA-3'), and probe upE-Prb (6-carboxyfluorescein[FAM]-5'-CTCTTCACATAATCGCCCCGAGCTCG-3'-6-carboxy-N,N,N,N'-tetramethylrhodamine [TAMRA]).

Determination of Cytokine Levels

Levels of interleukin 8 (IL-8), interferon-inducible protein 10, interferon- α , macrophage inflammatory protein- α , macrophage inflammatory protein- β , monocyte chemoattractant protein-1, monokine induced by interferon- γ (IFN- γ), IL-1 β , tumor necrosis factor- α , IL-6, and IL-10 in serum samples were determined by using a Cytometric Bead Assay Human Inflammatory Cytokine Kit (BD Biosciences, San Jose, CA, USA) according to the manufacturer's instructions. Data acquisition was performed by using a BD LSR Fortessa Analyzer (BD Biosciences). Data were analyzed by using cytometric bead assay analysis software (BD Biosciences).

IFN- γ ELISPOT Assay

Whole blood samples from healthy donors, health care workers who had been infected with SARS-CoV, and the MERS patient were collected into anticoagulant tubes containing EDTA, and the peripheral blood mononuclear cells (PBMCs) were

isolated by using standard density gradient centrifugation. An IFN- γ ELISPOT assay was performed as follows. A 96-well plate containing Immobilon-P membrane (MSIPS4510; Millipore, Billerica, MA, USA) was incubated with monoclonal anti-monkey IFN- γ (CT610–10; U-Cytech, Utrecht, the Netherlands) at 4°C overnight and blocked with RPMI medium 1640 (no. 11875–093; Gibco, Grand Island, NY, USA) containing 0.05 mmol/L 2-mercaptoethanol, 1 mmol/L sodium pyruvate, 2 mmol/L L-glutamine, 10 mmol/L HEPES, and 10% fetal bovine serum (no. SH30070; HyClone, Logan, UT, USA) at 37°C for 2 h.

PBMCs were plated at a density of 2.5×10^5 cells/well. Recombinant MERS-CoV spike protein (no. 40069-V08B; Sino Biologic Inc., North Wales, PA, USA) and SARS-CoV spike S1 subunit (no. 40150-V08B1; Sino Biologic Inc.) were added to the wells at a final concentration of 10 μ g/mL to stimulate PBMCs. Phosphate-buffered saline was used for mock stimulation. At 24-h poststimulation, IFN- γ was detected by using biotinylated anti-monkey IFN- γ (CT610–10; U-Cytech), and color was developed by using nitro-blue tetrazolium and 5-bromo-4-chloro-3'-indolyphosphate (Pierce, Rockford, IL, USA). Spots were counted by using an ELISPOT reader (Bioreader 4000; BIOSYS, Germany), and the results were reported as number of spot-forming cells/million PBMCs.

Microneutralization Test

Serum samples were heat-inactivated (56°C for 30 min) and mixed with equal volumes of two hundred 50% tissue cultures infectious doses of MERS-CoV. After 1 h of incubation at 37°C, 35 μ L of virus/serum mixture was added in quadruplicate to Vero cell monolayers in 96-well microtiter plates. After 1 h of adsorption, the virus/serum mixture was removed, 150 μ L of culture medium was added to each well, and plates were incubated for 3 days at 37°C in an atmosphere of 5% CO₂ in a humidified incubator. Cytopathic effect was read at day 3 postinfection. The highest serum dilution that completely protected the cells from a cytopathic effect in half of the wells was taken as the neutralizing antibody titer and was estimated by using the Reed-Muench method (1).

MERS-CoV Spike Pseudoparticle Neutralization Assay

MERS CoV pseudoparticles were incubated for 1 h with an equal volume of serum dilution at 4°C. After incubation for 1 h, serum/virus mixture was added to Vero cell monolayers in triplicate in a 96-well format. At day 3 posttransduction,

luciferase activity was measured by using a Microbeta Luminometer (PerkinElmer, Waltham, MA, USA). The highest serum dilution giving 90% reduction of luciferase activity was used as the antibody titer (2).

MERS CoV S1 ELISA

A commercially available (EUROIMMUN AG, Lübeck, Germany) MERS CoV S1 ELISA was used. In brief, serum dilutions, a calibrator, and controls were added to MERS CoV-coated plates and incubated for 30 min. Plates were washed 3 times and peroxidase labeled anti-human IgG was added and incubated for 30 min. Substrate solution was added and incubated for 15 min at room temperature. Reactions were stopped, and absorbance was measured at 450 nm using a reference wave length of 650 nm. The ratio between the extinction values of control/sample and calibrator was calculated. A ratio >0.8 was considered a positive result (3).

Plaque Reduction Neutralization Test

Equal volumes of each serum dilution and 40–60 PFU of MERS CoV/24-well plate were mixed and incubated for 1 h at 37°C in an atmosphere of 5% CO₂ in a humidified incubator. After incubation, virus/serum mixture was added to Vero cell monolayers in a 24-well format in duplicate. Plates were incubated for 1 h to enable absorption before overlaying with agarose. After 3 days, cells were fixed with 10% paraformaldehyde and stained. Titers are expressed as serum dilutions resulting in plaque inhibition \geq 90%.

References

- <jrn>1. Reed LJ, Muench H. A simple method of estimating fifty percent endpoints. American Journal of Hygiene. 1939;27:493–7.</jrn>
- <jrn>2. Hemida MG, Perera RA, Al Jassim RA, Kayali G, Siu LY, Wang P, et al. Seroepidemiology of Middle East respiratory syndrome (MERS) coronavirus in Saudi Arabia (1993) and Australia (2014) and characterisation of assay specificity. Euro Surveill. 2014;19:pii: 20828. [PubMed](#)</jrn>
- <jrn>3. Müller MA, Meyer B, Corman VM, Al-Masri M, Turkestani A, Ritz D, et al. Presence of Middle East respiratory syndrome coronavirus antibodies in Saudi Arabia: a nationwide, cross-sectional, serological study. Lancet Infect Dis. 2015;15:559–64. [PubMed http://dx.doi.org/10.1016/S1473-3099\(15\)70090-3](http://dx.doi.org/10.1016/S1473-3099(15)70090-3)</jrn>

Technical Appendix Table 1. Biochemical parameters for a 43-year-old traveler with Middle East respiratory syndrome, China, 2015*


| Parameter | Reference range | Days after illness onset | | | | | | |
|---|-----------------|--------------------------|------|-----|-----|-----|-----|-----|
| | | 9 | 11 | 16 | 21 | 23 | 29 | 33 |
| Total leukocytes, × 10 ⁹ cells/L | 4.0–10.0 | 3.22 | 3.06 | 3.6 | 4.9 | 4.8 | 4.6 | 7.3 |
| Neutrophils, × 10 ⁹ cells/L | 1.8–6.3 | 2.35 | 1.71 | 2.0 | 2.6 | 2.5 | 2.4 | 4.6 |
| Lymphocytes, × 10 ⁹ cells/L | 1.1–3.2 | 0.65 | 1.02 | 1.0 | 1.7 | 1.7 | 1.7 | 2.0 |
| Platelets, × 10 ⁹ /L | 125–350 | 81 | 113 | 253 | 378 | 362 | 274 | 263 |
| Hemoglobin, g/dL | 130–175 | 142 | 149 | 117 | 119 | 119 | 109 | 106 |
| Aspartate aminotransferase, U/L | 17–59 | 24 | 26 | – | 35 | 34 | 35 | 44 |
| Creatinine, μmol/L | 71–133 | 54 | 65 | – | 52 | 51 | 54 | 59 |
| Creatine kinase, U/L | 55–170 | – | – | – | 21 | 41 | 29 | 41 |
| Creatine kinase isoenzyme, U/L | 0–25 | 8 | 4 | – | 6 | 6 | 6 | 4 |

*–, data not collected.


Technical Appendix Table 2. Clinical characteristics for a 43-year-old traveler with Middle East respiratory syndrome, China, 2015*

| Characteristic | Result or value |
|-------------------------------------|---|
| Clinical features | |
| Fever | Yes |
| Highest temperature, °C | 39.5 |
| Cough | No |
| Sputum production | Yes |
| Shortness of breath | Yes |
| Chills | No |
| Complications | |
| Pneumonia | Yes |
| Acute respiratory distress syndrome | Yes |
| Shock | No |
| Bacterial co-infection | No |
| Treatment | |
| Oxygen therapy | Yes, inhalation oxygen with nasal catheter |
| Mechanical ventilation | No |
| Antivirus | Oseltamivir (150 mg, 2x/d on days 7 and 8); ribavirin (2.0 mg/d on day 8; 0.6 mg 3x/d on days 8–16; 0.6 mg 2x/d on days 17–19); peginterferon-α 2a (135 μg/d iv on day 8) |
| Antimicrobial drug | |
| Ceftriaxone sodium | 2.0 g/d iv on days 10–13 |
| Meropenem | 2.0 g 3x/d iv on days 13–26 |
| Glucocorticoids | No |
| Immune regulation therapy | |
| Immune globulin | 20 g/d iv on days 11–17 |
| Thymosin alpha-1 | 1.6 mg/d iv on days 14–18; 1.6 mg 2x/d iv on days 19–26 |
| Clinical outcome | Discharged |

*iv, intravenous.



Technical Appendix Figure 1. A) Timeline of clinical course of a 43-year-old traveler with Middle East respiratory syndrome, China, 2015. MERS-CoV, Middle East respiratory syndrome coronavirus. B–D) Chest radiographs on days 7, 11, and 35.



Technical Appendix Figure 2. Body temperature of 43-year-old traveler with Middle East respiratory syndrome, by day of hospitalization, China, 2015.


A

The viral load detected from various collection of samples during hospitalization


| Day of onset | MERS-CoV RNA copy numbers (log10) | | | | | | | | | | | | | | | |
|-----------------------|-----------------------------------|------|------|------|------|------|------|------|------|------|------|------|------|--|--|--|
| | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 22 | 25 | 26 | 36 | | | |
| Sputum on demand | 3.0 | 3.9 | 3.6 | 2.0 | 2.4 | N.C. | N.C. | N.C. | 2.8 | 2.2 | N.C. | N.C. | - | | | |
| Expectorated sputum* | N.C. | N.C. | N.C. | N.C. | N.C. | 6.0 | 6.1 | 2.5 | N.C. | N.C. | 2.0 | 3.0 | N.C. | | | |
| Throat Swab | - | - | - | - | - | - | - | N.C. | N.C. | N.C. | N.C. | N.C. | N.C. | | | |
| Nasal Swab | - | - | N.C. | - | - | N.C. | N.C. | N.C. | N.C. | N.C. | N.C. | N.C. | N.C. | | | |
| Serum | - | N.C. | N.C. | - | - | N.C. | - | N.C. | - | N.C. | - | N.C. | N.C. | | | |
| Stool | N.C. | N.C. | N.C. | 2.5 | N.C. | N.C. | N.C. | N.C. | 3.5 | N.C. | - | 2.1 | N.C. | | | |
| Urine | N.C. | N.C. | N.C. | N.C. | N.C. | N.C. | N.C. | - | - | N.C. | - | N.C. | N.C. | | | |
| Environmental samples | N.C. | N.C. | - | N.C. | - | N.C. | N.C. | N.C. | N.C. | N.C. | N.C. | N.C. | N.C. | | | |

Samples with threshold cycle lower than 40 were regarded as negative; N.C. : Not collected; - : Negative detection; Environmental samples : Palm, Pass Box, Bed railings (Day 13); Palm, Bed railings, Blanket, mobile phone, Floor (Day 15); *: A sputum cup was provided for the patient to collect sputum when he coughed. The sputum cup was collected in the morning for analysis.

B



C



D

Detection of antibodies against MERS-CoV by using different assays

| Days after onset of illness | MERS-CoV antibody | | | | |
|-----------------------------|-------------------|----------------------------|-----------------------------------|----------------------|--|
| | ppNT titer* | Micro-neutralization titer | PRNT [†] (90% reduction) | PRNT (50% reduction) | S1-ELISA [‡] (Calibrator/sample extinction ratio) |
| 11 | <1:10 | <1:10 | <1:10 | <1:10 | 0.03 (Negative) |
| 15 | 1:80 | 1:10 | 1:40 | 1:80 | 1.32 (Positive) |
| 18 | 1:320 | 1:40 | 1:80 | 1:160 | 2.25 (Positive) |
| 22 | 1:320 | 1:40 | 1:80 | 1:160 | 2.53 (Positive) |
| 25 | 1:320 | 1:40 | - | - | 2.46 (Positive) |
| 26 | 1:320 | 1:40 | 1:160 | 1:320 | 2.81 (Positive) |
| 34 | 1:640 | 1:80 | 1:160 | 1:320 | 3.1 (Positive) |

*ELISA optical density cut-off for positive result = 0.8; † ppNT: MERS-CoV spike pseudotype assay; ‡PRNT: Plaque reduction neutralization titer.

Technical Appendix Figure 3. Clinical parameters for a 43-year-old traveler with Middle East respiratory syndrome, China, 2015. A) Virus load. MERS-CoV, Middle East respiratory syndrome coronavirus. B) Cytokine levels and antibody titers. MIG, monokine induced by interferon- γ ; IP10, interferon-inducible protein 10; IFN- α , interferon- α ; IL-6, interleukin 6; MCP-1, monocyte chemoattractant protein-1. C) T-cell response of peripheral blood mononuclear cells (PBMCs) after challenge with MERS-CoV spike protein. SARS-CoV, severe acute respiratory syndrome coronavirus; IFN γ , interferon- γ ; SFCs, spot-forming cells. D) Detection of antibodies against MERS-CoV by different assays.