

Isolation of Batborne Neglected Zoonotic Agent Issyk-Kul Virus, Italy

Davide Lelli, Ana Moreno, Sabrina Canziani, Laura Soliani, Maya Carrera, Anna Castelli, Francesca Faccin, Tiziana Trogu, Enrica Sozzi, Gian Luca Cavallari, Matteo Mauri, Fabiana Ferrari, Cristian Salogni, Chiara Garbarino, Chiara Chiapponi, Marco Farioli, Antonio Lavazza

We isolated Issyk-Kul virus (ISKV) from a bat sampled from Italy in 2021 and conducted ISKV-specific surveillance in bats collected in Italy during 2017–2023. ISKV circulation among synanthropic and sedentary species of bat, such as Savi's pipistrelle bat (*Hypsugo savii*) in northern Italy, may have public health implications in this region.

Issyk-Kul virus (ISKV), family *Nairoviridae*, was first isolated in 1970 from a noctule bat (*Nyctalus noctule*) trapped near Lake Issyk-Kul, Kyrgyzstan (1). ISKV was subsequently detected in bats of several countries in central Asia and in *Ixodes vespertilionis* and *Argas vespertilionis* ticks (2). ISKV has caused sporadic outbreaks of illness in humans, characterized by fever, headaches, myalgia, and nausea (2,3). Bats and ticks are assumed to be reservoirs of ISKV; transmission to humans is associated with tick bites and exposure to bat feces and urine (2,4,5). Moreover, *Aedes caspius* mosquitoes, common in Europe and central Asia, may have a role as vectors, having been considered competent through experimental infection (6,7).

Portions of ISKV genome were detected in northern bats (*Eptesicus nilssonii*) in Germany (4) and a Brandt's bat (*Myotis brandtii*) in Sweden (8), suggesting that the ISKV geographic range expanded to Europe. We isolated and performed whole-genome characterization of ISKV detected in a Savi's

pipistrelle bat (*Hypsugo savii*) (hereafter referred to as Savi's bat) in Italy in 2021 and present the results of ISKV-specific surveillance of 415 bats collected during 2017–2023.

Ethics review and approval were waived for this study, which did not involve animal killing or suffering. Samples were collected exclusively from animals that died in wild recovery centers in the context of the regional surveillance plans for wildlife. Therefore, we believe that it does not fall in the provisions of the national law (e.g., DLSG 4/3 2014, n. 26 – Application at national level of the EU Directive 2010/63/UE), and no ethics approval or permit for animal experimentation was required.

The Study

We isolated the virus from an adult female Savi's bat that spontaneously died in a wildlife recovery center in northern Italy. The bat was originally found alive on August 17, 2021, in Bergamo Province, northern Italy, by a private citizen who brought it to the center. Clinically, the bat exhibited lethargy, inappetence, and weight loss. It died 11 days after admission to the center, and no trauma or macroscopic pathologic lesions indicative of infectious disease were observed at necropsy. DNA barcoding confirmed the species as Savi's bat. We collected organ samples (e.g., lung, heart, liver, spleen, intestine, and brain) for laboratory investigations focused mainly on virus detection.

We assessed the brains of all bats for negativity to rabies and related lyssaviruses by using real-time PCR (9). We isolated a virus on MARC 145 cells (fetal monkey kidney) inoculated with a pool of viscera (lung, heart, liver, spleen). Cytopathic effect was noted 5 days after inoculation during the secondary passage and was characterized by cell monolayer degeneration with isolated foci of rounded and aggregated cells (Figure 1, panels A–C). Furthermore, electron microscopy performed on cell culture supernatants

Author affiliations: University of Parma, Parma, Italy (D. Lelli); Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia Romagna, Brescia, Italy (D. Lelli, A. Moreno, S. Canziani, L. Soliani, M. Carrera, A. Castelli, F. Faccin, T. Trogu, E. Sozzi, C. Salogni, C. Garbarino, C. Chiapponi, A. Lavazza); Wildlife Rehabilitation Center WWF of Valpredina, Cenate Sopra, Italy (G.L. Cavallari, M. Mauri); Piacenza Wildlife Rescue Centre, Piacenza, Italy (F. Ferrari); DG Welfare Regione Lombardia, Milano, Italy (M. Farioli)

DOI: <https://doi.org/10.3201/eid3004.231186>




Figure 1. Microscopic appearance of Issyk-Kul virus IT-297348-34/2022, isolated from a *Hypsugo savii* bat, in study of batborne neglected zoonotic agent Issyk-Kul virus, Italy. A) Issyk-Kul-infected MARC 145 cells, mock infection; original magnification $\times 10$. B) Issyk-Kul-infected MARC 145 cells showing cytopathic effect at 120 hours after infection; original magnification $\times 4$. C) Issyk-Kul-infected MARC 145 cells showing cytopathic effect at 120 hours after infection; original magnification $\times 10$. D) Negative-staining electron microscopy performed on cell supernatants (NaPT 2%), showing a viral particle of 55–60 nm morphologically referable to nairovirus; original magnification $\approx 550,000$

revealed distinct viral particles of 55–60 nm, morphologically referable to a nairovirus (Figure 1, panel D).

The complete genome sequence of isolated virus IT-297348-34/2022 (IT-ISKV), obtained through a standardized next-generation sequencing protocol (10), revealed the 3 typical nairovirus genome segments: large (L) (11,978 nt), medium (M) (4,907 nt), and small (S) (1,457 nt). The highest nucleotide identity for each gene segment was to ISKV strains detected in a Brandt's bat (*Myotis brandtii*) and bat-associated ticks in Sweden (strain Sun-2020, k99_1658, k99_589), and in northern bats (*Eptesicus nilssonii*) in Germany (strain PbGER) (Table 1). We submitted the complete genome sequences to GenBank (accession nos. OR583909–11).

The phylogenetic tree, based on the complete L genome sequences of viruses in the genus *Orthonairovirus*, assigned IT-ISKV to the Keterah genogroup. That genogroup includes the few available sequences of ISKVs detected in bats and ticks in Sweden, Germany, and central Asia, as well as other sequences of Keterah virus (detected in bats from Malaysia), Uzun Agach virus (in bats from Kazakhstan), and Gossas virus (in

bats from Senegal) (Figure 1) (11). Phylogenetic trees constructed with complete S and M genes showed similar results with the same topology because of the small number of ISKV sequences available in the genome databases (Appendix Figures 1, 2, <https://wwwnc.cdc.gov/EID/article/30/4/23-1186-App1.pdf>).

After detecting IT-ISKV in the Mediterranean area, we developed and standardized ISKV-specific endpoint reverse transcription PCR targeting the L gene (Appendix) to enhance knowledge of the ISKV ecology and screen its diffusion in bat populations. We used ISKV-specific endpoint reverse transcription PCR to detect viral RNA in the necropsied tissues and to screen cultured cells. We performed Sanger sequencing of generated amplicons to confirm ISKV RNA.

During 2017–2023, we collected 415 bats representing 13 species in the Lombardy and Emilia-Romagna regions of northern Italy. The bats originated mainly from wildlife recovery centers, which usually receive rescued bats (usually near human settlements), or bats found dead during passive surveillance. Most of the bats examined were Savi's bats

Table 1. Highest nucleotide sequence identities for each protein of IT-ISKV isolated in study of batborne neglected zoonotic agent Issyk-Kul virus, Italy*

Gene	% Similarity (query cover, %)	ISKV strain	Host	Country (year)	GenBank accession no.	Reference
Large	95.44 (99)	K_k99_1658_len_12288	Bat-associated tick	Sweden (2020)	OP514654	Unpub. data
	95.34 (99)	LEZ 86–787	<i>Carios</i>	Germany (1986)	KR537441	(11)
	95.31 (99)	Sun-2020	<i>vespertilionis</i> tick			
Medium	81.55 (72)	Sun-2020	<i>Myotis brandtii</i> bat	Sweden (2020)	OP380632	(8)
	81.11 (78)	k99_589	<i>M. brandtii</i> bat	Sweden (2020)	OP380631	(8)
	81.34 (71)	LEIV-315K	Bat-associated tick	Sweden (2020)	OP804626	Unpub. data
			<i>Nyctalus noctula</i> bat	Kyrgyzstan (1973)	KR709220	(3)
Small	97.51 (90)	PbGER	<i>Eptesicus nilssonii</i> bat	Germany (2008–2011)	MW275296	(4)
	89.11 (100)	Sun-2020	<i>M. brandtii</i> bat	Sweden (2020)	OP380630	(8)
	89.04 (100)	LEZ 86–787	<i>C. vespertilionis</i> tick	Germany (1986)	KR537443	(11)

*IT-ISKV, Issyk-Kul virus IT-297348-34/2022.

and Kuhl's bats (*Pipistrellus kuhlii*) (Appendix Table 1). We tested sampled organs (lung, heart, liver, spleen, intestine) for ISKV. We detected 8 bats positive by PCR for ISKV; 7 were Savi's bats and 1 was a whiskered bat (*Myotis mystacinus*) (Table 2), and they were recovered in 2017, 2020, 2021, 2022, and 2023 (Appendix Table 2). We constructed a phylogenetic tree based on the partial L genome sequences with all ISKVs detected in Italy (Figure 2, panel B). None of the ISKV-positive bats had ticks attached, and the ticks (*Ixodes vespertilionis*) found on the analyzed bats were ISKV negative.

Conclusions

Tickborne orthonairoviruses may be agents of human emerging infectious diseases (13). Crimean-Congo hemorrhagic fever virus is the most notable pathogen in the genus *Orthonairovirus* because of its public health effect

with high fatality rates and widespread geographic distribution (14). However, several other emerging and neglected orthonairoviruses, such as ISKV, can cause clinical nonlethal diseases in humans (15).

Our isolation and characterization of IT-ISKV showed high L and S gene identity to the ISKV strains detected in Sweden, Germany, and central Asia (3,4,8). However, the level of M gene nucleotide similarity to the other known ISKV strains (80.98%–81.55%) suggests that IT-ISKV could represent a new ISKV strain from the Mediterranean area, most likely derived from an assortment with a yet unknown virus.

In that context, we conducted ISKV-specific surveillance among bats collected during 2017–2023 with the aim of determining the presence and diffusion of the virus in bat populations in northern Italy. Findings suggest that ISKV in that area seem to be associated with Savi's bats and whiskered bats, which may

Table 2. Issyk-Kul–positive bats and associated GenBank accession number assigned to the gene sequences detected in study of batborne neglected zoonotic agent Issyk-Kul virus, Italy*

Sample	Year	Bat species	Origin	Virus isolation (cell culture)	Sequence	GenBank accession no.	Nucleotide similarity (%)
251170-38/2017	2017	<i>Hypsugo savii</i>	WRC WWF Valpredina, Bergamo, Italy	No	Partial L gene	OR583901	Issyk-Kul virus LEZ 86–787 (98.79)
251170-41/2017	2017	<i>H. savii</i>	WRC WWF Valpredina, Bergamo, Italy	No	Partial L gene	OR583902	Issyk-Kul virus LEZ 86–787 (99.27)
378052-30/2020	2020	<i>H. savii</i>	WRC WWF Valpredina, Bergamo, Italy	No	Partial L gene	OR583903	Issyk-Kul virus K_k99_1658_len_12288 (99.17)
297348-34/2022	2021	<i>H. savii</i>	WRC WWF Valpredina, Bergamo, Italy	Yes	Full genome	OR583909–11, OR583905	Issyk-Kul virus/Prackenbach bat nairovirus (96.65)
297348-26/2022	2021	<i>H. savii</i>	WRC WWF Valpredina, Bergamo, Italy	No	Partial L gene	OR583904	Issyk-Kul virus LEZ 86–787 (99.42)
126482-17/2022	2022	<i>H. savii</i>	WRC Piacenza, Italy	No	Partial L gene	OR583906	Issyk-Kul virus LEZ 86–787 (99)
356061-37/2022	2022	<i>H. savii</i>	WRC WWF Valpredina, Bergamo, Italy	No	Partial L gene	OR583907	Issyk-Kul virus/Prackenbach bat nairovirus (98.42)
24094-8/2023	2023	<i>Myotis mystacinus</i>	WRC Piacenza, Italy	No	Partial L gene	OR583908	Issyk-Kul virus LEZ 86–787 (98.91)

*L, large; WRC, wildlife recovery center; WWF, World Wildlife Fund.




Figure 2. Phylogenetic analysis of isolates from study of batborne neglected zoonotic agent Issyk-Kul virus, Italy A) Phylogeneny nairovirus protein sequences for *Orthonaviridae* large (L) segments, including the complete sequence obtained from *Hypsugo savii* bats, highlighted in yellow. Sequence colors were based on the genogroups proposed by Ozeki et al. 2022 (12). B) Nucleotide alignment magnification of a short PCR-targeted region of the L segment, encompassing all sequences derived from bat surveillance conducted during 2017–2023 identified as Issyk-Kul virus IT-297348-34/2022. Numbers along branches indicate bootstrap values.

represent a previously unrecognized source for ISKV transmission to other wildlife species, ticks, and humans, as has already happened elsewhere (2,3). Savi's bats are a synanthropic and sedentary species that roost in buildings and represent the most common bat species in urban areas, suggesting possible public health implications.

ISKVs identified in bats in Italy were detected from a pool of organs, as in the previously described ISKV PbGER (4), which was found predominantly in the liver, spleen, and lung tissues, indicating systemic infection of bats instead of mere passaging of intestinal tick content. Future investigations may provide information about virus tissue distribution and pathogenesis by using histopathology and may define the infection prevalence in the bat populations through serologic tests.

The successful cell-culture isolation of IT-ISKV suggests the possible shedding of infectious virus particles, which represents a crucial point for assessing viral zoonotic risks that may emerge from synanthropic bats. Such results indicate the emergence of

this neglected zoonotic agent in the Mediterranean area, which might have public health relevance because of its potential transmission to humans. Raising awareness of the risks deriving from this zoonosis should suggest adoption of specific surveillance and prevention programs for ISKV and other nairoviruses at the human–wild animal interface.

Acknowledgments

We thank the technicians of the IZSLER Virology section for their precious technical work.

This study was supported by the Lombardy and Emilia Romagna Wildlife Surveillance Program.

About the Author

Dr. Lelli is a senior researcher virologist at the Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia-Romagna. His research activity involves diagnosing livestock and wildlife viral diseases. The major topics of his studies are emerging viral diseases, arthropodborne viruses, and viruses associated with bats.

References

1. Lvov DK, Karas FR, Timofeev EM, Tsyarkin YM, Vargina SG, Veselovskaya OV, et al. "Issyk-Kul" virus, a new arbovirus isolated from bats and *Argas (Carios) vespertilionis* (Latr., 1802) in the Kirghiz S.S.R. *Arch Gesamte Virusforsch.* 1973;42:207–9. <https://doi.org/10.1007/BF01270841>
2. L'vov DK, Kostiuikov MA, Daniyarov OA, Tukhtaev TM, Sherikov BK. Outbreak of arbovirus infection in the Tadzhik SSR due to the Issyk-Kul virus (Issyk-Kul fever) [in Russian]. *Vopr Virusol.* 1984;29:89–92.
3. Atkinson B, Marston DA, Ellis RJ, Fooks AR, Hewson R. Complete genomic sequence of Issyk-Kul virus. *Genome Announc.* 2015;3:e00662–00715. 10.1128/genomeA.00662-15 <https://doi.org/10.1128/genomeA.00662-15>
4. Brinkmann A, Kohl C, Radonić A, Dabrowski PW, Mühlendorfer K, Nitsche A, et al. First detection of bat-borne Issyk-Kul virus in Europe. *Sci Rep.* 2020;10:22384. 10.1038/s41598-020-79468-8 <https://doi.org/10.1038/s41598-020-79468-8>
5. Vargina SG, Kuchuk LA, Gershtein VI, Karas FR. Transmission of Issyk Kul virus by *Argas vespertilionis* ticks in experiment [in Russian]. *Sborn nauch Tr Inst Virus Im Ivanov Akad Med Nauk SSSR.* 1982;123–7.
6. Bulychev VP, Alekseev AN, Kostiuikov MA, Gordeeva ZE, L'vov DK. Issyk-Kul virus transmission by *Aedes caspius caspius* Pall. mosquitoes via experimental bite [in Russian]. *Med Parazitol (Mosk).* 1979;48:53–6.
7. Kostiuikov MA, Bulychev VP, Lapina TF. Experimental infection of *Aedes caspius caspius* Pall. mosquitoes on dwarf bats, *Vespertilio pipistrellus*, infected with the Issyk-Kul virus and its subsequent transmission to susceptible animals [in Russian]. *Med Parazitol (Mosk).* 1982;51:78–9.
8. Cholleti H, de Jong J, Blomström AL, Berg M. Investigation of the virome and characterization of Issyk-Kul virus from Swedish *Myotis brandtii* bats. *Pathogens.* 2022;12:12. <https://doi.org/10.3390/pathogens12010012>
9. Gigante CM, Dettinger L, Powell JW, Seiders M, Condori REC, Griesser R, et al. Multi-site evaluation of the LN34 pan-lyssavirus real-time RT-PCR assay for post-mortem rabies diagnostics. *PLoS One.* 2018;13:e0197074. <https://doi.org/10.1371/journal.pone.0197074>
10. Lelli D, Prosperi A, Moreno A, Chiapponi C, Gibellini AM, De Benedictis P, et al. Isolation of a novel rhabdovirus from an insectivorous bat (*Pipistrellus kuhlii*) in Italy. *Virol J.* 2018;15:37. <https://doi.org/10.1186/s12985-018-0949-z>
11. Walker PJ, Widen SG, Wood TG, Guzman H, Tesh RB, Vasilakis N. A global genomic characterization of nairoviruses identifies nine discrete genogroups with distinctive structural characteristics and host-vector associations. *Am J Trop Med Hyg.* 2016;94:1107–22. <https://doi.org/10.4269/ajtmh.15-0917>
12. Ozeki T, Abe H, Ushijima Y, Nze-Nkogue C, Akomo-Okoue EF, Ella GWE, et al. Identification of novel orthonairoviruses from rodents and shrews in Gabon, Central Africa. *J Gen Virol.* 2022;103. <https://doi.org/10.1099/jgv.0.001796>
13. Lv X, Liu Z, Li L, Xu W, Yuan Y, Liang X, et al. Yezo virus infection in tick-bitten patient and ticks, northeastern China. *Emerg Infect Dis.* 2023;29:797–800. <https://doi.org/10.3201/eid2904.220885>
14. Shayan S, Bokaeian M, Shahrivar MR, Chinikar S. Crimean-Congo hemorrhagic fever. *Lab Med.* 2015;46:180–9. <https://doi.org/10.1309/LMN1P2FRZ7BKZSCO>
15. Sugimoto S, Suda Y, Nagata N, Fukushima S, Yoshikawa T, Kurosu T, et al. Characterization of Kemerah orthonairovirus and evaluation of therapeutic candidates against Kemerah orthonairovirus infectious disease. *Ticks Tick Borne Dis.* 2022;13:101834. <https://doi.org/10.1016/j.ttbdis.2021.101834>

Address for correspondence: Davide Lelli, Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia Romagna "B. Ubertyni," Via bianchi 9, Brescia 25124, Italy; email: davide.elli@izsler.it

Isolation of Bat-Borne Neglected Zoonotic Agent Issyk-Kul Virus, Italy

Appendix

Materials and Methods

Sampling

From January 2017 to January 2023, 415 bats were analyzed in the context of surveillance implemented in Northern Italy regarding emerging viruses of bats, which was launched in 2010 (1,2). Dead bats of different species were collected for virological investigations from wild animal rescue/rehabilitation centers. The bat species were taxonomically identified by both morphologic characteristics according to the European bat identification keys (3) and by a molecular method (4). The anamnestic data were gathered when available. The bat carcasses were necropsied, the tissue samples were collected, and subjected to a diagnostic protocol broadly targeting viral agents.

Virus isolation and Electron Microscopy

After necropsy, organ samples (lungs, heart, kidney, brain, and intestines) were homogenized in minimal essential medium (1 g/10 mL), which contained antibiotics, and were centrifuged at 3000 g for 15 min. Samples were inoculated in confluent monolayers of MARC 145 and VERO (African green monkey) cells, incubated at 37°C with 5% CO₂, and observed daily for 7 days for cytopathic effects (CPEs). In the absence of CPEs, the cryo-lysates were subcultured twice onto fresh monolayers. The supernatant from cell cultures showing CPEs was submitted for viral identification with negative-staining electron microscopy (nsEM) by using the Airfuge method (5) (Airfuge, Beckman Coulter, Inc. Life Sciences, Indianapolis, Indiana, USA). The examination was done using a Tecnai G2 Spirit Biotwin transmission electron microscope (TEM; FEI, Hillsboro, Oregon, OR, USA) operating at 85 kV. The observation was

done at 13,500–43,000x for at least 15 min before being considered negative. Identification and recognition of the observed viral particles were based on their morphology.

End-point one-step RT-PCR for ISKV detection

An endpoint one-step RT-PCR was developed using the complete viral genome sequenced. Primers IZSLER-ISKV F (5'- CTAGCTCTGCTGATTATGAG –3') and IZSLER-ISKV R (5'- GCCAAGATAGTTGCGTCAATC –3') were designed on a conserved 844 bp region of Katerah Genogroup L gene, which encodes for the RNA-dependent RNA polymerase of the ISKV. Total RNA was extracted from samples using QIAasympyphony DSP Virus/Pathogen Mini Kit, following the manufacturer's instructions. The PCR assay was performed with the QIAGEN® OneStep RT-PCRmaster mix (Qiagen, Hilden, Germany). The reaction mixture contained 0.6 pmol of sense and antisense primer, Qiagen buffer 1X, 400 µM of each dNTPs, Qiagen OneStep RT-PCR Enzyme mix 1µl, 8U of RNase inhibitor and 5 µl of the RNA template in a final volume of 25 µl. PCR conditions were optimized using homogenized bat organs and cell culture samples, which were known to be positive and negative for ISKV infections. The thermal cycling conditions consisted of 30 min at 50°C for reverse transcription, 15 min at 95°C for the initial enzyme activation, and 40 thermal cycles of 94°C for 60 s; 59°C for 60 s; and 72°C for 30 s, with a final elongation step of 72°C for 10 min. Amplicons were visualized on 2% agarose electrophoretic gels stained with EuroSafe Fluorescent Nucleic Acid Stain (Euroclone, Milan, Italy) used at a 1× concentration.

Positive samples were confirmed by Sanger sequencing using the IZSLER ISKV forward and reverse primers.

Phylogenetic Analysis

A collection of nairovirus complete protein sequences was downloaded based on the classification provided by the International Committee on Taxonomy of Viruses (ICTV), including all defined member species of the genus Orthonairovirus, available as of July 1, 2023. The sequences were aligned using the CLUSTAL algorithm (6). For the L segment, a nucleotide alignment of a targeted region (positions 368–760) was performed using the same algorithm. A maximum-likelihood tree was generated using Iqtree-1.6.12 (7), with the best-fit model automatically selected by ModelFinder. The statistical significance of the tree topology was

assessed using ultrafast bootstrap with 1000 replicates. Tree visualizations were generated using FigTree v.1.1.4.

References

1. Lelli D, Moreno A, Lavazza A, Bresola M, Canelli E, Boniotti MB, et al. Identification of Mammalian orthoreovirus type 3 in Italian bats. *Zoonoses Public Health*. 2013;60:84–92. [PubMed](#) <https://doi.org/10.1111/zph.12001>
2. Lelli D, Papetti A, Sabelli C, Rosti E, Moreno A, Boniotti MB. Detection of coronaviruses in bats of various species in Italy. *Viruses*. 2013;5:2679–89. [PubMed](#) <https://doi.org/10.3390/v5112679>
3. Dietz C, von Helversen O. Illustrated identification key to the bats of Europe. Electronic publication. Version 1.0. Released 15.12.2004. Tuebingen & Erlangen (Germany).
4. De Benedictis P, Leopardi S, Markotter W, Velasco-Villa A. The Importance of Accurate Host Species Identification in the Framework of Rabies Surveillance, Control and Elimination. *Viruses*. 2022;14:492. [PubMed](#) <https://doi.org/10.3390/v14030492>
5. Lavazza A, Pascucci S, Gelmetti D. Rod-shaped virus-like particles in intestinal contents of three avian species. *Vet Rec*. 1990;126:581. [PubMed](#)
6. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol Syst Biol*. 2011;7:539. [PubMed](#) <https://doi.org/10.1038/msb.2011.75>
7. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol*. 2015;32:268–74. [PubMed](#) <https://doi.org/10.1093/molbev/msu300>
8. Ozeki T, Abe H, Ushijima Y, Nze-Nkogue C, Akomo-Okoue EF, Ella GWE, et al. Identification of novel orthonairoviruses from rodents and shrews in Gabon, Central Africa. *J Gen Virol*. 2022;103. [PubMed](#) <https://doi.org/10.1099/jgv.0.001796>

Appendix Table 1. List of bat species analyzed in the study

Species	N°	%
<i>Eptesicus serotinus</i>	1	0.2
<i>Hypsugo savii</i>	170	41
<i>Myotis emarginatus</i>	1	0.2
<i>Myotis crypticus</i>	1	0.2
<i>Myotis daubentonii</i>	9	2.2
<i>Myotis mystacinus</i>	4	1.0
<i>Nyctalus leisleri</i>	3	0.7
<i>Pipistrellus kuhlii</i>	151	36.4
<i>Pipistrellus nathusii</i>	6	1.5
<i>Pipistrellus pipistrellus</i>	15	3.6
<i>Plecotus uratus</i>	3	0.7
<i>Plecotus macrobullaris</i>	1	0.2
<i>Tadarida teniotis</i>	4	1.0
<i>Plecotus sp.</i>	2	0.5
<i>Vespertilio murinus</i>	1	0.2
<i>Pipistrellus spp.</i>	43	10.4
Total	415	100


Appendix Table 2. Number of bats analyzed in the study divided by year of sampling

Year	N° tested bats	N° positive bats
2017	98	2
2018	42	0
2019	30	0
2020	117	1
2021	58	2
2022	52	1
2023	18	2
Total	415	8

Appendix Table 3. Details of the ISKV-positive bats in the study: clinical signs, organs tested and positive

N° Identification	Year	Species	Age	Sex	Clinical signs	Sampled organs	Positive samples
251170/38	2017	<i>Hypsugo savii</i>	J	n.a.	No signs	I + V	I + V
251170/41	2017	<i>Hypsugo savii</i>	J	n.a.	No signs	I + V	I + V
378052/30	2020	<i>Hypsugo savii</i>	SA	F	No signs	I + V	I + V
297348/26	2021	<i>Hypsugo savii</i>	SA	M	Dehydration	I + V	I + V
297348/34	2021	<i>Hypsugo savii</i>	SA	F	Apathy, inappetence weight loss	I + V	I + V
126482/17	2022	<i>Hypsugo savii</i>	J	F	No signs	I + V	V
356061/37	2023	<i>Hypsugo savii</i>	n.a.	n.a.	No signs	I + V	I
24094/8	2023	<i>Myotis mystacinus</i>	A	M	Traumatic lesions (cat aggression)	I + V	I + V

A = adult, SA = sub-adult (this-year-born individuals, usually ranging in age from 1 to 10–11 mo) J = juvenile, - recently born, incapable of flight, M = male, F = female, I = intestine, V = pool of viscera (heart + lung + spleen + liver)
n.a. = not available.



Appendix Figure 2. Phylogenetic analysis of nairovirus protein sequences for S segments, including our complete sequence obtained from *Hypsugo savii*, highlighted. The sequences were colored based on the genogroups proposed by Ozeki *et al.* 2022 (8).