

# TABLES

TABLE 1. Sample Sizes and Response Rates, Selected U.S. Sites: School Health Profiles, Principal and Lead Health Education Teacher Surveys, 2004

Site	Principal surveys		Teacher surveys	
	Sample size	Response rate (%)	Sample size	Response rate (%)
<b>STATE SURVEYS</b>				
Alaska	291	82	250	70
Arizona	317	75	303	71
Arkansas	212	75	211	74
Connecticut	258	78	242	74
Delaware	58	81	58	81
Idaho	202	91	186	84
Iowa	275	79	254	73
Maine	193	85	187	83
Massachusetts	607	85	581	82
Michigan	324	80	318	79
Minnesota	278	71	NA*	NA
Missouri	353	82	331	77
Montana	313	93	311	92
Nebraska	282	81	259	74
New Hampshire	168	79	167	79
New York	313	71	306	70
North Carolina	281	70	280	70
North Dakota	203	85	196	82
Oklahoma	351	89	340	86
Oregon	249	75	235	71
Pennsylvania	330	74	318	72
South Carolina	240	77	228	73
Tennessee	314	81	307	79
Texas**	376	75	NA	NA
Utah	208	86	185	76
Washington	265	80	243	73
Wisconsin	313	74	317	75
<b>LOCAL SURVEYS</b>				
Chicago	262	78	258	77
Dallas	50	98	50	98
District of Columbia	32	74	31	72
Los Angeles	96	76	98	78
Memphis	55	87	50	79
Miami	96	92	95	91
New Orleans	33	73	33	73
Orange County	41	82	40	80
Philadelphia	125	84	113	76
San Bernardino	16	94	NA	NA
San Diego	42	98	42	98

\* NA = not available.  
 \*\* Survey did not include schools from two of the state's largest school districts.

TABLE 2. Percentage of Schools That Required Health Education in Any of Grades 6–12, and Among Those Schools, the Percentage That Taught Required Health Education in a Combined Course or in Another Course, the Percentage That Required Students to Take Only One Course or Two or More Courses, and the Percentage That Required Students Who Fail a Required Health Education Course to Repeat It, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2004

Site	Required health education	Taught required health education in a combined health education and physical education course	Taught required health education in another course*	Required only one health education course	Required two or more health education courses	Required students who fail a required health education course to repeat it
<b>STATE SURVEYS</b>						
Alaska	89.2	63.2	39.6	46.8	41.2	87.0
Arizona	59.9	56.9	45.0	53.3	25.7	51.7
Arkansas	97.3	62.5	16.6	57.1	41.0	82.6
Connecticut	89.5	49.3	20.9	30.3	59.6	42.1
Delaware	100.0	69.4	9.1	41.1	50.7	53.4
Idaho	95.8	46.0	10.8	52.9	44.4	60.1
Iowa	81.4	40.2	39.0	43.1	47.6	62.1
Maine	97.9	43.0	16.5	51.3	46.5	67.1
Massachusetts	86.3	53.1	17.2	34.7	59.0	39.6
Michigan	85.9	49.5	27.8	55.7	34.3	53.3
Minnesota	98.4	62.1	16.8	26.3	69.5	69.5
Missouri	91.5	68.0	29.8	42.3	49.6	56.3
Montana	94.6	96.4	20.8	12.6	76.7	63.8
Nebraska	91.8	77.1	35.1	48.1	46.7	58.7
New Hampshire	84.1	33.5	13.6	43.8	50.4	54.7
New York	98.1	33.5	9.3	43.6	54.0	60.5
North Carolina	95.0	88.8	15.1	39.4	48.2	48.1
North Dakota	94.3	69.3	36.6	20.8	75.4	56.6
Oklahoma	27.3	62.0	60.4	47.6	22.8	18.6
Oregon	98.3	52.5	22.5	19.7	75.2	54.1
Pennsylvania	96.8	68.2	14.8	28.8	67.8	58.2
South Carolina	85.9	68.7	26.4	44.3	36.2	37.1
Tennessee	73.0	82.1	38.3	56.1	28.0	66.2
Texas**	78.9	40.1	18.2	77.0	16.0	83.0
Utah	98.0	35.2	12.8	76.2	22.7	53.8
Washington	92.3	58.6	28.4	54.4	36.4	63.2
Wisconsin	96.3	40.9	22.4	45.3	49.5	72.9
<b>State Median</b>	<b>92.3</b>	<b>58.6</b>	<b>20.9</b>	<b>44.3</b>	<b>47.6</b>	<b>58.2</b>
<b>State Range</b>	<b>27.3 – 100.0</b>	<b>33.5 – 96.4</b>	<b>9.1 – 60.4</b>	<b>12.6 – 77.0</b>	<b>16.0 – 76.7</b>	<b>18.6 – 87.0</b>
<b>LOCAL SURVEYS</b>						
Chicago	79.1	83.6	59.4	34.7	28.1	31.9
Dallas	63.2	52.0	30.1	89.4	3.5	76.2
District of Columbia	90.5	100.0	4.5	79.5	17.4	80.8
Los Angeles	99.0	15.0	45.5	77.6	17.7	47.5
Memphis	83.7	86.1	27.7	53.1	33.0	60.9
Miami	69.2	69.5	56.8	63.1	19.4	81.6
New Orleans	81.6	100.0	9.7	49.7	46.4	74.4
Orange County	68.5	65.9	43.9	56.0	28.6	63.6
Philadelphia	86.6	79.5	30.3	36.1	46.4	37.7
San Bernardino	81.3	60.0	75.0	33.3	16.7	50.0
San Diego	45.2	70.9	94.6	NA <sup>§</sup>	NA	NA
<b>Local Median</b>	<b>81.3</b>	<b>70.9</b>	<b>43.9</b>	<b>54.6</b>	<b>23.8</b>	<b>62.3</b>
<b>Local Range</b>	<b>45.2 – 99.0</b>	<b>15.0 – 100.0</b>	<b>4.5 – 94.6</b>	<b>33.3 – 89.4</b>	<b>3.5 – 46.4</b>	<b>31.9 – 81.6</b>

\* e.g., science, social studies, home economics, or English.

\*\* Survey did not include schools from two of the state's largest school districts.

§ NA = data not available.

TABLE 3. Among Schools That Required a Health Education Course in Any of Grades 6–12, the Percentage That Taught a Required Health Education Course in Each Grade, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2004

Site	6th grade	7th grade	8th grade	9th grade	10th grade	11th grade	12th grade
<b>STATE SURVEYS</b>							
Alaska	47.2	51.9	54.3	84.0	61.4	52.8	49.9
Arizona	76.0	84.9	79.9	75.9	58.8	39.3	42.9
Arkansas	40.4	68.6	25.9	81.5	62.3	49.9	52.2
Connecticut	95.7	87.7	92.6	84.9	71.7	55.1	54.0
Delaware	84.6	96.4	89.4	77.4	55.0	21.1	15.8
Idaho	45.0	65.8	73.4	15.1	60.1	44.2	15.7
Iowa	51.5	61.9	60.8	69.8	46.4	35.0	26.0
Maine	75.2	81.1	75.8	68.2	49.2	10.9	9.0
Massachusetts	85.9	87.7	90.3	82.6	63.5	35.1	24.6
Michigan	47.2	73.0	55.6	75.3	33.7	16.2	12.4
Minnesota	61.7	71.9	77.8	47.3	87.1	15.8	12.3
Missouri	68.4	85.1	80.4	81.6	30.8	19.7	19.7
Montana	72.8	91.7	93.3	93.6	89.6	11.1	8.9
Nebraska	54.3	69.7	62.2	65.6	36.8	10.9	9.5
New Hampshire	78.8	90.6	85.3	61.0	53.9	18.9	13.8
New York	49.8	71.2	57.2	30.7	68.9	54.2	48.8
North Carolina	93.2	94.3	93.8	96.9	19.8	10.9	10.0
North Dakota	82.8	81.5	77.2	64.2	36.0	11.2	12.0
Oklahoma	78.6	71.5	53.5	34.1	38.5	27.1	24.2
Oregon	76.9	90.0	89.6	60.8	62.2	43.2	12.4
Pennsylvania	76.8	73.1	80.9	61.1	55.0	57.8	28.8
South Carolina	97.5	93.9	90.9	89.3	55.8	42.8	46.1
Tennessee	75.5	80.3	77.4	89.0	38.7	19.8	20.4
Texas*	39.1	35.2	42.0	87.3	60.1	51.4	48.7
Utah	66.5	47.7	58.7	11.6	91.1	23.3	18.9
Washington	65.4	65.4	74.1	72.2	51.3	25.6	26.3
Wisconsin	49.4	59.5	63.4	59.4	40.7	20.6	12.4
<b>State Median</b>	<b>72.8</b>	<b>73.1</b>	<b>77.2</b>	<b>72.2</b>	<b>55.0</b>	<b>25.6</b>	<b>19.7</b>
<b>State Range</b>	<b>39.1 – 97.5</b>	<b>35.2 – 96.4</b>	<b>25.9 – 93.8</b>	<b>11.6 – 96.9</b>	<b>19.8 – 91.1</b>	<b>10.9 – 57.8</b>	<b>8.9 – 54.0</b>
<b>LOCAL SURVEYS</b>							
Chicago	84.6	89.5	88.0	97.6	53.8	29.2	28.9
Dallas	NA**	100.0	83.3	76.2	71.4	75.0	65.0
District of Columbia	42.9	62.5	75.0	73.6	100.0	25.0	25.0
Los Angeles	35.5	97.7	NA	97.4	23.1	8.8	8.8
Memphis	86.7	91.0	82.8	100.0	50.0	37.5	37.5
Miami	56.9	46.3	46.3	35.2	91.7	26.5	23.5
New Orleans	100.0	84.8	84.8	93.4	76.9	46.2	46.2
Orange County	73.2	100.0	86.6	94.4	50.0	50.0	50.0
Philadelphia	93.8	92.6	94.5	70.7	84.4	64.5	45.2
San Bernardino	100.0	100.0	100.0	75.0	66.7	33.3	33.3
San Diego	NA	NA	NA	NA	NA	NA	NA
<b>Local Median</b>	<b>84.6</b>	<b>91.8</b>	<b>84.8</b>	<b>84.8</b>	<b>69.1</b>	<b>35.4</b>	<b>35.4</b>
<b>Local Range</b>	<b>35.5 – 100.0</b>	<b>46.3 – 100.0</b>	<b>46.3 – 100.0</b>	<b>35.2 – 100.0</b>	<b>23.1 – 100.0</b>	<b>8.8 – 75.0</b>	<b>8.8 – 65.0</b>

\* Survey did not include schools from two of the state's largest school districts.

\*\* NA = data not available.

TABLE 4. Percentage of Schools That Required Teachers to Use Specific Materials in a Required Health Education Course, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	National health education standards	State, district, or school curriculum, guidelines, or framework	Materials from health organizations	Commercially-developed student textbook	Commercially-developed teacher's guide
<b>STATE SURVEYS</b>					
Alaska	44.7	95.5	25.0	48.9	43.4
Arizona	51.9	95.4	45.2	55.1	53.4
Arkansas	43.4	99.4	31.3	81.4	73.1
Connecticut	55.7	97.1	38.9	23.4	22.9
Delaware	84.5	98.2	21.6	27.2	19.8
Idaho	41.2	97.4	29.6	68.5	53.1
Iowa	44.5	97.0	28.2	46.3	38.0
Maine	30.3	96.3	15.5	24.2	22.5
Massachusetts	46.9	97.6	40.8	33.9	39.1
Michigan	32.1	95.9	35.6	36.1	34.6
Missouri	47.3	99.0	38.0	64.1	58.6
Montana	47.1	95.5	30.0	48.2	46.6
Nebraska	31.0	92.0	30.2	59.7	54.7
New Hampshire	58.7	88.8	29.1	31.3	31.3
New York	65.9	98.5	36.3	39.3	31.7
North Carolina	43.8	98.9	48.5	68.9	65.4
North Dakota	46.4	84.4	33.1	57.2	54.8
Oklahoma	31.5	98.7	51.5	58.4	57.0
Oregon	43.7	96.2	27.5	53.5	47.2
Pennsylvania	63.6	96.8	49.6	63.9	57.9
South Carolina	58.3	96.6	36.3	71.6	61.6
Tennessee	49.2	98.3	45.8	79.9	75.0
Utah	31.8	100.0	27.7	58.3	50.5
Washington	32.6	91.3	33.7	54.1	53.3
Wisconsin	54.1	95.6	34.4	41.0	36.0
<b>State Median</b>	<b>46.4</b>	<b>96.8</b>	<b>33.7</b>	<b>54.1</b>	<b>50.5</b>
<b>State Range</b>	<b>30.3 – 84.5</b>	<b>84.4 – 100.0</b>	<b>15.5 – 51.5</b>	<b>23.4 – 81.4</b>	<b>19.8 – 75.0</b>
<b>LOCAL SURVEYS</b>					
Chicago	50.7	96.3	52.0	54.9	48.2
Dallas	60.8	100.0	87.8	74.8	62.4
District of Columbia	58.8	100.0	56.9	53.2	46.1
Los Angeles	37.7	93.9	30.1	61.1	40.8
Memphis	79.9	100.0	45.1	84.3	81.1
Miami	51.2	100.0	58.0	72.1	62.8
New Orleans	58.8	100.0	58.8	72.2	65.3
Orange County	50.0	100.0	51.2	87.7	58.7
Philadelphia	48.3	100.0	44.3	46.3	42.2
San Diego	NA*	NA	NA	NA	NA
<b>Local Median</b>	<b>51.2</b>	<b>100.0</b>	<b>52.0</b>	<b>72.1</b>	<b>58.7</b>
<b>Local Range</b>	<b>37.7 – 79.9</b>	<b>93.9 – 100.0</b>	<b>30.1 – 87.8</b>	<b>46.3 – 87.7</b>	<b>40.8 – 81.1</b>

\* NA = data not available.

TABLE 5a. Percentage of Schools That Tried to Increase Student Knowledge on a Specific Health-Related Topic in a Required Health Education Course During the 2003–2004 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Accident or injury prevention	Alcohol or other drug use prevention	Consumer health	CPR*	Death and dying	Dental and oral health	Emotional and mental health
<b>STATE SURVEYS</b>							
Alaska	82.3	94.0	79.6	51.2	59.8	73.4	87.6
Arizona	89.7	95.7	79.0	57.2	59.8	69.5	87.0
Arkansas	96.2	99.5	88.4	69.7	71.3	84.5	98.4
Connecticut	86.5	99.5	72.1	42.6	48.0	46.2	94.7
Delaware	90.3	100.0	83.9	56.8	40.1	41.9	98.2
Idaho	92.4	100.0	83.9	65.1	74.3	66.5	99.4
Iowa	82.7	96.8	78.8	62.3	60.7	61.8	93.2
Maine	89.0	95.6	88.5	54.3	46.2	42.0	95.6
Massachusetts	87.2	98.3	79.1	52.7	50.7	52.2	97.0
Michigan	77.6	96.4	74.8	47.0	44.9	56.8	86.4
Missouri	95.3	98.4	89.5	73.2	64.0	77.7	94.4
Montana	91.9	98.4	85.2	71.4	57.0	69.9	91.2
Nebraska	85.7	99.1	77.2	67.8	59.4	63.6	92.6
New Hampshire	90.8	98.4	87.3	55.1	43.8	50.0	95.2
New York	88.6	99.4	81.3	58.6	64.8	60.6	96.0
North Carolina	93.9	99.6	80.5	75.7	58.7	61.6	96.1
North Dakota	95.6	99.3	87.0	73.5	60.6	83.7	97.5
Oklahoma	90.1	98.5	83.2	58.0	56.5	88.7	74.9
Oregon	89.6	99.5	83.9	64.5	61.0	59.2	97.2
Pennsylvania	89.3	98.9	74.9	70.2	48.1	58.3	93.1
South Carolina	93.1	96.2	74.3	50.5	60.1	72.5	88.8
Tennessee	93.7	98.3	82.1	74.6	64.4	74.9	90.0
Utah	92.6	99.4	85.2	70.9	76.5	58.2	98.7
Washington	79.9	97.9	74.2	59.1	42.6	51.9	88.0
Wisconsin	86.1	99.7	80.3	57.9	59.8	56.8	94.9
<b>State Median</b>	<b>89.7</b>	<b>98.5</b>	<b>81.3</b>	<b>59.1</b>	<b>59.8</b>	<b>61.6</b>	<b>94.7</b>
<b>State Range</b>	<b>77.6 – 96.2</b>	<b>94.0 – 100.0</b>	<b>72.1 – 89.5</b>	<b>42.6 – 75.7</b>	<b>40.1 – 76.5</b>	<b>41.9 – 88.7</b>	<b>74.9 – 99.4</b>
<b>LOCAL SURVEYS</b>							
Chicago	89.1	96.9	73.6	62.8	54.5	79.9	82.9
Dallas	100.0	100.0	87.4	91.7	79.8	67.9	100.0
District of Columbia	96.3	89.5	66.7	67.8	63.7	79.0	85.8
Los Angeles	86.0	97.6	89.3	56.8	64.2	79.0	93.0
Memphis	96.9	100.0	91.1	75.8	75.8	84.5	94.0
Miami	83.9	100.0	81.5	81.3	63.2	72.7	86.2
New Orleans	96.6	100.0	72.4	55.0	64.5	66.1	89.6
Orange County	96.3	100.0	71.0	95.7	79.0	52.5	100.0
Philadelphia	87.7	96.2	75.7	60.0	49.8	73.9	86.3
San Diego	NA**	NA	NA	NA	NA	NA	NA
<b>Local Median</b>	<b>96.3</b>	<b>100.0</b>	<b>75.7</b>	<b>67.8</b>	<b>64.2</b>	<b>73.9</b>	<b>89.6</b>
<b>Local Range</b>	<b>83.9 – 100.0</b>	<b>89.5 – 100.0</b>	<b>66.7 – 91.1</b>	<b>55.0 – 95.7</b>	<b>49.8 – 79.8</b>	<b>52.5 – 84.5</b>	<b>82.9 – 100.0</b>

\* CPR = cardiopulmonary resuscitation.  
 \*\* NA = data not available.

TABLE 5b. Percentage of Schools That Tried to Increase Student Knowledge on a Specific Health-Related Topic in a Required Health Education Course During the 2003–2004 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Environmental health	First aid	Growth and development	HIV* prevention	Human sexuality	Immunization and vaccinations	Nutrition and dietary behavior
<b>STATE SURVEYS</b>							
Alaska	76.6	66.3	84.4	82.1	75.6	66.1	92.5
Arizona	79.8	68.9	90.7	86.4	72.9	59.3	98.5
Arkansas	82.2	83.7	91.6	94.3	82.3	71.8	98.3
Connecticut	56.3	56.5	92.9	98.8	93.3	53.8	97.0
Delaware	57.2	72.7	88.8	95.6	95.7	49.3	97.7
Idaho	72.0	78.2	88.8	99.4	79.6	69.7	98.8
Iowa	73.9	69.2	91.7	96.4	84.9	67.5	97.8
Maine	64.0	70.4	88.8	97.7	93.2	52.6	98.8
Massachusetts	60.8	64.8	91.5	95.9	89.3	51.6	98.2
Michigan	68.1	59.7	87.5	96.5	82.2	47.3	95.1
Missouri	77.4	88.0	93.1	97.8	86.2	76.5	98.9
Montana	70.0	76.8	90.5	94.6	85.0	61.5	100.0
Nebraska	71.4	76.7	92.6	95.8	85.7	59.4	98.5
New Hampshire	59.4	65.5	90.5	98.2	93.0	56.1	96.8
New York	69.8	72.4	94.8	98.9	92.9	70.4	99.6
North Carolina	74.6	85.9	84.2	95.2	81.0	62.3	99.2
North Dakota	80.8	83.4	95.6	96.6	86.7	74.5	99.4
Oklahoma	83.0	79.1	87.4	93.9	78.6	74.2	97.0
Oregon	67.9	77.6	94.1	98.0	93.2	64.7	98.9
Pennsylvania	60.1	80.2	91.9	98.1	90.9	60.4	97.7
South Carolina	73.9	72.3	93.2	97.1	93.1	62.3	99.5
Tennessee	79.3	84.2	89.6	95.3	86.2	76.2	97.8
Utah	69.7	76.1	96.2	98.0	91.2	66.0	99.4
Washington	64.2	70.5	85.7	96.7	91.6	45.9	97.9
Wisconsin	63.4	70.6	94.0	98.1	93.8	56.1	99.3
<b>State Median</b>	<b>70.0</b>	<b>72.7</b>	<b>91.5</b>	<b>96.6</b>	<b>86.7</b>	<b>62.3</b>	<b>98.5</b>
<b>State Range</b>	<b>56.3 – 83.0</b>	<b>56.5 – 88.0</b>	<b>84.2 – 96.2</b>	<b>82.1 – 99.4</b>	<b>72.9 – 95.7</b>	<b>45.9 – 76.5</b>	<b>92.5 – 100.0</b>
<b>LOCAL SURVEYS</b>							
Chicago	72.8	75.5	93.0	92.6	83.1	67.3	97.4
Dallas	92.0	95.9	95.9	100.0	91.7	80.0	100.0
District of Columbia	71.5	92.9	89.1	100.0	96.5	76.0	100.0
Los Angeles	75.4	69.4	96.4	97.6	95.2	78.4	98.8
Memphis	91.0	91.0	97.0	100.0	93.9	78.1	100.0
Miami	79.3	77.2	93.1	100.0	97.6	68.1	92.8
New Orleans	75.9	86.1	96.6	100.0	89.6	65.8	100.0
Orange County	56.2	79.6	100.0	100.0	100.0	53.1	100.0
Philadelphia	73.4	79.3	92.7	93.7	88.3	64.0	97.4
San Diego	NA**	NA	NA	NA	NA	NA	NA
<b>Local Median</b>	<b>75.4</b>	<b>79.6</b>	<b>95.9</b>	<b>100.0</b>	<b>93.9</b>	<b>68.1</b>	<b>100.0</b>
<b>Local Range</b>	<b>56.2 – 92.0</b>	<b>69.4 – 95.9</b>	<b>89.1 – 100.0</b>	<b>92.6 – 100.0</b>	<b>83.1 – 100.0</b>	<b>53.1 – 80.0</b>	<b>92.8 – 100.0</b>

\* HIV = human immunodeficiency virus.

\*\* NA = data not available.

TABLE 5c. Percentage of Schools That Tried to Increase Student Knowledge on a Specific Health-Related Topic in a Required Health Education Course During the 2003–2004 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Personal hygiene	Physical activity and fitness	Pregnancy prevention	STD* prevention	Suicide prevention	Sun safety or skin cancer prevention	Tobacco use prevention	Violence prevention
<b>STATE SURVEYS</b>								
Alaska	84.5	94.8	75.3	80.0	73.6	54.7	93.1	84.0
Arizona	88.9	99.3	70.1	73.1	59.2	72.7	94.8	82.5
Arkansas	93.3	100.0	84.2	89.9	84.5	79.7	100.0	91.2
Connecticut	81.1	98.9	86.3	93.5	74.7	59.6	98.4	90.9
Delaware	78.3	100.0	89.4	95.7	66.3	65.0	100.0	84.8
Idaho	85.9	98.2	75.7	91.1	84.2	77.8	100.0	88.7
Iowa	80.2	97.6	83.0	91.1	77.0	73.7	98.3	80.0
Maine	76.0	98.7	90.4	94.4	73.0	73.8	99.4	83.5
Massachusetts	83.3	98.2	84.1	91.5	74.8	76.9	98.5	94.0
Michigan	81.7	96.2	71.9	89.0	60.9	62.8	97.4	81.4
Missouri	93.9	98.8	85.8	92.6	74.3	82.0	99.0	87.1
Montana	92.1	100.0	77.3	88.7	70.5	75.1	99.2	88.6
Nebraska	86.6	99.0	79.7	88.1	70.7	72.4	98.5	81.5
New Hampshire	81.1	97.5	87.2	94.2	75.0	65.1	98.4	87.6
New York	84.4	99.6	90.5	96.4	86.2	80.1	99.2	88.9
North Carolina	79.8	100.0	83.1	90.2	72.7	76.7	100.0	90.7
North Dakota	94.0	98.7	77.7	87.7	82.5	86.5	100.0	90.9
Oklahoma	95.7	100.0	73.1	81.6	65.8	63.6	98.4	90.1
Oregon	81.8	99.4	89.0	94.2	76.7	74.6	99.6	88.6
Pennsylvania	83.2	98.9	84.7	94.5	67.3	71.9	98.6	81.8
South Carolina	90.0	100.0	89.9	94.6	62.6	66.5	96.3	83.1
Tennessee	92.2	99.4	90.2	93.2	75.7	78.0	98.2	88.6
Utah	82.9	99.4	75.9	94.6	94.2	75.7	100.0	93.1
Washington	78.7	98.5	82.1	91.9	74.4	55.7	96.8	86.8
Wisconsin	80.7	98.2	90.3	95.3	83.2	71.9	99.2	84.4
<b>State Median</b>	<b>83.3</b>	<b>98.9</b>	<b>84.1</b>	<b>91.9</b>	<b>74.4</b>	<b>73.7</b>	<b>98.6</b>	<b>87.6</b>
<b>State Range</b>	<b>76.0 – 95.7</b>	<b>94.8 – 100.0</b>	<b>70.1 – 90.5</b>	<b>73.1 – 96.4</b>	<b>59.2 – 94.2</b>	<b>54.7 – 86.5</b>	<b>93.1 – 100.0</b>	<b>80.0 – 94.0</b>
<b>LOCAL SURVEYS</b>								
Chicago	93.2	100.0	78.7	82.8	50.2	55.9	95.0	90.1
Dallas	84.1	100.0	87.3	91.7	87.6	79.8	100.0	100.0
District of Columbia	89.5	100.0	100.0	100.0	64.4	43.8	92.0	93.2
Los Angeles	89.5	98.8	95.2	96.4	75.1	87.0	100.0	87.0
Memphis	97.0	100.0	97.0	100.0	79.1	66.5	100.0	96.9
Miami	84.0	87.9	95.4	100.0	79.3	74.7	100.0	79.5
New Orleans	96.5	100.0	82.4	96.5	55.0	48.0	100.0	86.4
Orange County	84.0	100.0	100.0	100.0	91.4	72.8	100.0	92.0
Philadelphia	85.5	97.6	78.4	92.3	53.9	51.2	94.8	90.7
San Diego	NA**	NA	NA	NA	NA	NA	NA	NA
<b>Local Median</b>	<b>89.5</b>	<b>100.0</b>	<b>95.2</b>	<b>96.5</b>	<b>75.1</b>	<b>66.5</b>	<b>100.0</b>	<b>90.7</b>
<b>Local Range</b>	<b>84.0 – 97.0</b>	<b>87.9 – 100.0</b>	<b>78.4 – 100.0</b>	<b>82.8 – 100.0</b>	<b>50.2 – 91.4</b>	<b>43.8 – 87.0</b>	<b>92.0 – 100.0</b>	<b>79.5 – 100.0</b>

\* STD = sexually transmitted disease.  
 \*\* NA = data not available.

TABLE 6. Percentage of Schools That Tried To Improve Specific Student Skills in a Required Health Education Course During the 2003–2004 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Accessing valid health information, products, and services	Advocating for personal, family, and community health	Analysis of media messages	Communication	Decision making	Goal setting	Conflict resolution	Resisting peer pressure for unhealthy behaviors	Stress management
<b>STATE SURVEYS</b>									
Alaska	76.3	72.4	66.7	82.7	90.7	85.6	84.7	89.2	78.4
Arizona	79.1	81.7	77.4	94.2	95.5	92.7	86.5	92.8	83.5
Arkansas	90.0	84.8	77.3	96.4	100.0	96.3	91.4	99.0	94.1
Connecticut	88.9	81.3	90.5	95.5	97.6	92.1	87.8	97.8	89.5
Delaware	91.8	94.2	92.4	96.0	100.0	96.0	90.9	97.9	85.7
Idaho	89.3	82.6	87.3	96.3	98.2	93.7	93.0	98.7	94.9
Iowa	87.1	86.0	83.6	91.6	97.7	90.9	85.3	94.9	88.1
Maine	92.7	83.4	92.8	95.6	97.7	95.0	87.3	93.3	92.1
Massachusetts	85.2	79.3	90.6	96.0	98.8	94.4	92.1	98.5	88.4
Michigan	80.9	79.5	85.0	89.0	96.6	92.1	87.4	97.1	84.1
Missouri	90.8	85.0	83.0	93.3	98.7	94.8	90.0	97.0	91.6
Montana	88.4	85.6	84.6	92.8	98.4	94.3	89.2	96.8	83.0
Nebraska	82.4	79.4	80.0	85.5	97.0	91.8	82.9	95.0	89.8
New Hampshire	84.6	82.4	94.6	91.4	96.7	88.8	86.1	98.4	88.0
New York	93.9	89.3	92.4	95.6	98.5	95.6	91.4	99.2	96.3
North Carolina	89.0	82.6	80.6	90.3	96.4	94.2	90.8	97.2	92.6
North Dakota	86.9	82.1	87.5	95.1	98.8	94.5	92.1	97.4	93.4
Oklahoma	79.0	75.7	54.5	76.1	93.0	94.3	85.9	97.0	78.4
Oregon	87.6	81.8	93.2	96.4	99.2	93.5	93.9	98.5	92.9
Pennsylvania	85.2	80.0	85.7	92.3	97.8	95.1	83.9	95.4	86.1
South Carolina	86.4	87.1	81.7	90.2	94.3	95.7	92.1	93.5	89.1
Tennessee	84.6	82.1	76.4	92.6	97.5	95.5	90.3	96.0	91.7
Utah	87.3	85.0	91.7	97.5	99.4	97.0	93.2	100.0	98.0
Washington	82.2	76.3	80.4	91.1	93.1	94.8	87.6	94.1	85.5
Wisconsin	86.2	87.7	90.3	94.7	97.2	92.2	86.5	97.2	92.5
<b>State Median</b>	<b>86.9</b>	<b>82.4</b>	<b>85.0</b>	<b>93.3</b>	<b>97.7</b>	<b>94.3</b>	<b>89.2</b>	<b>97.1</b>	<b>89.5</b>
<b>State Range</b>	<b>76.3 – 93.9</b>	<b>72.4 – 94.2</b>	<b>54.5 – 94.6</b>	<b>76.1 – 97.5</b>	<b>90.7 – 100.0</b>	<b>85.6 – 97.0</b>	<b>82.9 – 93.9</b>	<b>89.2 – 100.0</b>	<b>78.4 – 98.0</b>
<b>LOCAL SURVEYS</b>									
Chicago	74.8	79.3	61.6	84.2	92.3	90.1	90.7	93.8	74.8
Dallas	100.0	91.6	76.1	96.0	100.0	100.0	100.0	100.0	95.9
District of Columbia	89.5	78.6	88.7	93.2	100.0	96.6	100.0	96.6	75.6
Los Angeles	90.5	83.6	90.5	92.8	94.1	86.1	90.7	92.7	81.0
Memphis	87.9	93.8	78.9	97.0	100.0	100.0	100.0	100.0	94.1
Miami	81.7	77.0	77.0	90.8	95.5	90.8	79.4	95.3	83.9
New Orleans	86.4	82.7	72.2	83.0	96.5	93.0	93.1	100.0	75.8
Orange County	84.0	87.7	87.7	100.0	100.0	100.0	100.0	95.7	100.0
Philadelphia	85.0	74.2	79.8	89.8	94.9	96.2	91.3	95.1	77.4
San Diego	NA*	NA	NA	NA	NA	NA	NA	NA	NA
<b>Local Median</b>	<b>86.4</b>	<b>82.7</b>	<b>78.9</b>	<b>92.8</b>	<b>96.5</b>	<b>96.2</b>	<b>93.1</b>	<b>95.7</b>	<b>81.0</b>
<b>Local Range</b>	<b>74.8 – 100.0</b>	<b>74.2 – 93.8</b>	<b>61.6 – 90.5</b>	<b>83.0 – 100.0</b>	<b>92.3 – 100.0</b>	<b>86.1 – 100.0</b>	<b>79.4 – 100.0</b>	<b>92.7 – 100.0</b>	<b>74.8 – 100.0</b>

\* NA = data not available.

TABLE 7a. Percentage of Schools That Taught About Health Outcomes and Risks of Tobacco Use in a Required Health Education Course During the 2003–2004 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Addictive effects of nicotine	Benefits of not smoking cigarettes	Benefits of not using smokeless tobacco	Health consequences of cigarette smoking	Health consequences of using smokeless tobacco	Health effects of ETS*	Number of illnesses and deaths related to tobacco use	Risks of cigar or pipe smoking
<b>STATE SURVEYS</b>								
Alaska	90.3	92.7	85.9	92.8	88.6	88.6	85.6	76.2
Arizona	91.2	91.9	88.4	91.9	88.6	89.2	89.6	83.1
Arkansas	98.9	99.5	99.4	100.0	99.4	98.4	97.1	94.2
Connecticut	95.5	96.3	92.1	95.9	93.7	95.1	95.6	87.2
Delaware	98.1	100.0	96.2	100.0	98.1	100.0	100.0	88.0
Idaho	98.2	99.4	97.6	99.4	98.8	97.5	98.2	92.4
Iowa	95.2	96.7	91.8	96.8	94.7	94.6	93.6	83.8
Maine	95.5	97.1	86.3	97.2	91.4	93.7	91.4	74.6
Massachusetts	97.0	97.4	92.0	97.7	94.5	95.7	94.9	88.0
Michigan	94.9	94.6	91.7	96.2	93.3	95.2	92.8	87.5
Missouri	97.2	98.1	96.3	98.1	95.9	96.6	96.0	92.8
Montana	96.7	97.1	94.6	97.5	95.4	94.6	93.9	84.8
Nebraska	97.0	96.4	96.3	96.9	96.9	95.9	91.5	85.8
New Hampshire	95.1	96.0	89.2	96.8	92.3	92.6	93.6	81.1
New York	98.1	97.7	94.4	99.2	96.6	97.5	96.5	91.6
North Carolina	98.0	98.4	95.9	99.6	99.2	97.7	96.2	92.2
North Dakota	99.3	99.4	96.3	100.0	98.8	98.7	97.5	86.5
Oklahoma	92.7	98.4	89.9	98.4	92.7	94.1	89.9	85.3
Oregon	99.0	98.4	95.5	99.0	98.5	97.8	95.9	87.5
Pennsylvania	96.3	95.8	93.5	97.2	95.3	94.7	94.1	87.8
South Carolina	94.2	94.6	90.4	95.1	91.8	89.7	90.9	82.8
Tennessee	96.8	96.8	94.1	98.2	96.2	94.8	95.3	88.0
Utah	99.4	100.0	95.2	100.0	97.6	98.1	96.2	89.5
Washington	93.4	93.8	90.5	95.1	93.9	91.0	89.7	79.9
Wisconsin	98.4	97.6	95.1	98.5	96.8	97.0	95.9	89.3
<b>State Median</b>	<b>96.8</b>	<b>97.1</b>	<b>94.1</b>	<b>97.7</b>	<b>95.4</b>	<b>95.2</b>	<b>94.9</b>	<b>87.5</b>
<b>State Range</b>	<b>90.3 – 99.4</b>	<b>91.9 – 100.0</b>	<b>85.9 – 99.4</b>	<b>91.9 – 100.0</b>	<b>88.6 – 99.4</b>	<b>88.6 – 100.0</b>	<b>85.6 – 100.0</b>	<b>74.6 – 94.2</b>
<b>LOCAL SURVEYS</b>								
Chicago	84.9	87.2	74.3	91.3	74.8	85.2	88.5	79.0
Dallas	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
District of Columbia	77.9	81.1	59.3	84.7	59.7	73.0	66.7	63.5
Los Angeles	96.3	98.8	95.2	98.8	95.2	95.1	96.3	91.5
Memphis	100.0	100.0	96.9	100.0	96.9	94.1	97.0	94.0
Miami	92.8	97.6	87.8	97.6	81.0	92.8	88.1	88.1
New Orleans	100.0	100.0	92.9	100.0	92.9	100.0	100.0	93.0
Orange County	100.0	100.0	95.7	100.0	100.0	96.3	95.7	92.6
Philadelphia	92.1	94.6	87.2	94.6	90.9	90.7	90.8	87.0
San Diego	NA**	NA	NA	NA	NA	NA	NA	NA
<b>Local Median</b>	<b>96.3</b>	<b>98.8</b>	<b>92.9</b>	<b>98.8</b>	<b>92.9</b>	<b>94.1</b>	<b>95.7</b>	<b>91.5</b>
<b>Local Range</b>	<b>77.9 – 100.0</b>	<b>81.1 – 100.0</b>	<b>59.3 – 100.0</b>	<b>84.7 – 100.0</b>	<b>59.7 – 100.0</b>	<b>73.0 – 100.0</b>	<b>66.7 – 100.0</b>	<b>63.5 – 100.0</b>

\* ETS = environmental tobacco smoke.  
 \*\* NA = data not available.

TABLE 7b. Percentage of Schools That Taught About the External Influences on Tobacco Use in a Required Health Education Course During the 2003–2004 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Influence of families on tobacco use	Influence of the media on tobacco use	Social or cultural influences on tobacco use	How many young people use tobacco
<b>STATE SURVEYS</b>				
Alaska	82.5	83.7	82.4	83.5
Arizona	83.7	87.3	84.9	85.4
Arkansas	96.6	97.1	94.4	96.7
Connecticut	91.3	95.2	86.1	93.9
Delaware	98.1	96.2	89.8	98.1
Idaho	97.1	98.2	89.9	93.1
Iowa	90.1	95.2	86.4	87.8
Maine	87.1	90.3	79.6	85.1
Massachusetts	91.8	96.1	88.8	92.8
Michigan	91.0	93.8	87.7	90.8
Missouri	92.9	94.6	88.7	92.7
Montana	92.6	93.7	91.1	91.5
Nebraska	91.5	91.0	86.9	87.8
New Hampshire	88.7	95.9	91.0	89.6
New York	94.5	96.2	90.4	93.8
North Carolina	96.5	96.6	93.6	95.4
North Dakota	98.1	96.9	92.0	96.2
Oklahoma	87.2	90.0	87.5	88.8
Oregon	93.5	98.2	93.0	93.1
Pennsylvania	93.1	94.5	90.0	91.2
South Carolina	86.7	87.1	83.3	87.4
Tennessee	92.6	93.9	90.9	93.4
Utah	97.0	99.3	91.6	92.2
Washington	85.9	88.1	81.7	82.9
Wisconsin	91.1	98.0	91.2	93.0
<b>State Median</b>	<b>91.8</b>	<b>95.2</b>	<b>89.8</b>	<b>92.2</b>
<b>State Range</b>	<b>82.5 – 98.1</b>	<b>83.7 – 99.3</b>	<b>79.6 – 94.4</b>	<b>82.9 – 98.1</b>
<b>LOCAL SURVEYS</b>				
Chicago	81.8	80.9	82.7	86.9
Dallas	100.0	95.9	95.7	100.0
District of Columbia	67.8	77.9	66.7	66.7
Los Angeles	91.4	97.5	96.4	93.9
Memphis	96.9	96.9	94.0	94.1
Miami	85.7	83.4	88.1	90.4
New Orleans	93.0	100.0	96.4	96.5
Orange County	95.7	100.0	96.3	95.7
Philadelphia	88.2	90.7	80.7	91.0
San Diego	NA*	NA	NA	NA
<b>Local Median</b>	<b>91.4</b>	<b>95.9</b>	<b>94.0</b>	<b>93.9</b>
<b>Local Range</b>	<b>67.8 – 100.0</b>	<b>77.9 – 100.0</b>	<b>66.7 – 96.4</b>	<b>66.7 – 100.0</b>

\* NA = data not available.

TABLE 7c. Percentage of Schools That Taught Skills to Avoid and to Stop Using Tobacco in a Required Health Education Course During the 2003–2004 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	How to say no to tobacco use	Making a personal commitment not to use tobacco	How students can influence or support others to prevent tobacco use	How to find valid information or services related to tobacco use cessation	How students can influence or support others in efforts to quit using tobacco
<b>STATE SURVEYS</b>					
Alaska	87.6	69.3	79.4	69.4	78.3
Arizona	87.9	74.0	84.7	80.1	84.9
Arkansas	98.9	80.2	93.3	76.9	94.4
Connecticut	91.7	72.0	85.2	71.8	82.4
Delaware	94.0	83.9	94.1	85.8	94.1
Idaho	98.3	76.3	92.5	80.1	91.3
Iowa	91.3	71.0	84.1	69.4	79.8
Maine	87.0	62.4	78.3	74.3	76.1
Massachusetts	94.4	76.2	89.0	77.9	87.4
Michigan	94.2	75.4	88.1	75.5	86.4
Missouri	96.0	75.2	91.2	78.0	90.5
Montana	92.2	74.3	85.4	74.4	83.5
Nebraska	91.9	66.1	86.8	71.3	84.2
New Hampshire	88.6	69.5	83.0	71.3	77.2
New York	95.8	77.2	90.5	85.1	89.9
North Carolina	97.1	84.8	93.0	79.8	91.4
North Dakota	97.5	77.9	93.4	76.3	90.2
Oklahoma	94.2	84.4	92.7	77.8	87.1
Oregon	94.7	72.3	89.3	79.1	85.3
Pennsylvania	94.0	74.7	86.9	76.4	83.2
South Carolina	93.0	79.5	83.2	72.9	80.7
Tennessee	95.3	80.7	90.0	72.8	89.6
Utah	98.7	80.8	92.0	72.0	92.6
Washington	91.5	65.8	83.7	74.4	84.0
Wisconsin	95.5	71.8	88.2	76.4	88.8
<b>State Median</b>	<b>94.2</b>	<b>75.2</b>	<b>88.2</b>	<b>76.3</b>	<b>86.4</b>
<b>State Range</b>	<b>87.0 – 98.9</b>	<b>62.4 – 84.8</b>	<b>78.3 – 94.1</b>	<b>69.4 – 85.8</b>	<b>76.1 – 94.4</b>
<b>LOCAL SURVEYS</b>					
Chicago	87.8	79.0	87.0	66.2	83.9
Dallas	100.0	83.4	100.0	95.7	100.0
District of Columbia	71.5	57.6	67.8	64.4	68.1
Los Angeles	96.4	89.0	87.9	87.9	87.9
Memphis	100.0	90.9	97.0	76.0	97.0
Miami	90.4	78.8	85.7	76.3	83.4
New Orleans	100.0	85.7	100.0	89.4	96.5
Orange County	96.3	71.6	88.3	92.0	96.3
Philadelphia	90.7	83.4	87.1	79.6	85.8
San Diego	NA*	NA	NA	NA	NA
<b>Local Median</b>	<b>96.3</b>	<b>83.4</b>	<b>87.9</b>	<b>79.6</b>	<b>87.9</b>
<b>Local Range</b>	<b>71.5 – 100.0</b>	<b>57.6 – 90.9</b>	<b>67.8 – 100.0</b>	<b>64.4 – 95.7</b>	<b>68.1 – 100.0</b>

\* NA = data not available.

TABLE 8a. Percentage of Schools That Taught HIV\* Transmission and Prevention Topics in a Required Health Education Course During the 2003–2004 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Abstinence as the most effective method to avoid HIV infection	How HIV is transmitted	How HIV affects the human body	Condom efficacy
<b>STATE SURVEYS</b>				
Alaska	79.2	80.5	78.9	57.2
Arizona	77.5	78.5	76.2	46.9
Arkansas	92.1	91.2	89.7	60.7
Connecticut	97.5	97.7	95.3	80.8
Delaware	95.5	91.3	91.3	78.5
Idaho	97.5	96.1	93.6	65.9
Iowa	92.3	91.8	89.6	74.4
Maine	96.5	95.4	93.7	83.8
Massachusetts	94.0	94.2	91.8	75.6
Michigan	92.6	92.8	91.3	56.9
Missouri	95.2	95.0	93.7	68.7
Montana	91.1	90.8	88.8	62.9
Nebraska	90.8	90.4	86.9	56.0
New Hampshire	95.6	95.9	93.3	75.1
New York	98.5	98.1	98.1	87.2
North Carolina	92.4	91.6	89.7	58.2
North Dakota	89.4	91.6	88.3	54.5
Oklahoma	86.4	89.6	85.3	55.7
Oregon	94.9	95.8	95.3	79.9
Pennsylvania	95.4	97.6	96.3	75.9
South Carolina	94.5	93.9	91.8	63.0
Tennessee	92.8	93.3	93.3	67.5
Utah	95.6	96.1	96.1	35.5
Washington	95.6	95.5	95.0	75.3
Wisconsin	98.0	97.7	96.1	83.0
<b>State Median</b>	<b>94.5</b>	<b>93.9</b>	<b>91.8</b>	<b>67.5</b>
<b>State Range</b>	<b>77.5 – 98.5</b>	<b>78.5 – 98.1</b>	<b>76.2 – 98.1</b>	<b>35.5 – 87.2</b>
<b>LOCAL SURVEYS</b>				
Chicago	84.0	85.9	84.4	60.0
Dallas	100.0	100.0	100.0	75.5
District of Columbia	100.0	100.0	100.0	96.3
Los Angeles	97.5	97.5	96.3	85.4
Memphis	100.0	100.0	100.0	76.1
Miami	100.0	100.0	100.0	95.4
New Orleans	96.4	96.4	100.0	68.5
Orange County	100.0	100.0	100.0	87.0
Philadelphia	92.2	93.6	93.6	79.8
San Diego	NA**	NA	NA	NA
<b>Local Median</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>79.8</b>
<b>Local Range</b>	<b>84.0 – 100.0</b>	<b>85.9 – 100.0</b>	<b>84.4 – 100.0</b>	<b>60.0 – 96.3</b>

\* HIV = human immunodeficiency virus.

\*\* NA = data not available.

TABLE 8b. Percentage of Schools That Taught About the External Influences on HIV\* Risk Behavior and Skills to Avoid HIV Infection in a Required Health Education Course During the 2003–2004 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Number of young people who get HIV	Influence of alcohol and other drugs on HIV-related risk behaviors	Social or cultural influences on HIV-related risk behaviors	How to find valid information or services related to HIV or HIV testing	How to correctly use a condom	Compassion for persons living with HIV or AIDS**
<b>STATE SURVEYS</b>						
Alaska	71.2	71.8	67.1	63.8	35.1	61.8
Arizona	66.6	67.3	63.2	64.4	25.2	63.0
Arkansas	81.2	88.8	79.9	73.1	23.7	74.8
Connecticut	89.6	93.0	81.8	81.0	55.3	88.8
Delaware	82.9	91.1	81.0	82.6	57.7	74.8
Idaho	87.6	95.2	82.2	81.9	20.4	83.1
Iowa	76.2	88.1	77.7	71.2	47.1	72.6
Maine	82.3	93.6	82.8	84.6	60.0	82.9
Massachusetts	85.2	91.5	78.6	80.7	50.4	86.8
Michigan	84.8	89.9	81.7	78.7	28.6	76.9
Missouri	83.9	91.1	82.6	75.5	31.8	75.3
Montana	81.4	86.7	77.4	71.5	28.9	75.9
Nebraska	79.5	85.7	75.4	71.3	25.4	75.2
New Hampshire	90.0	95.6	83.3	81.0	57.9	86.3
New York	94.5	98.5	86.4	92.1	59.1	90.8
North Carolina	83.5	88.4	82.4	72.5	17.2	75.2
North Dakota	79.8	88.7	76.3	74.0	23.5	78.1
Oklahoma	79.3	82.0	77.6	65.8	25.0	70.7
Oregon	90.8	93.4	81.2	82.2	48.8	82.5
Pennsylvania	89.9	92.9	85.0	82.4	43.1	84.7
South Carolina	86.0	89.7	82.2	77.8	42.6	77.2
Tennessee	86.3	87.3	85.5	83.4	37.6	77.6
Utah	86.4	94.9	87.9	73.4	8.8	87.8
Washington	85.6	94.3	81.9	81.1	45.5	79.7
Wisconsin	89.7	95.2	83.9	79.9	55.0	84.1
<b>State Median</b>	<b>84.8</b>	<b>91.1</b>	<b>81.8</b>	<b>78.7</b>	<b>37.6</b>	<b>77.6</b>
<b>State Range</b>	<b>66.6 – 94.5</b>	<b>67.3 – 98.5</b>	<b>63.2 – 87.9</b>	<b>63.8 – 92.1</b>	<b>8.8 – 60.0</b>	<b>61.8 – 90.8</b>
<b>LOCAL SURVEYS</b>						
Chicago	80.2	76.2	73.1	70.2	42.0	76.8
Dallas	95.7	100.0	95.7	87.1	43.7	95.7
District of Columbia	96.6	96.6	100.0	96.3	85.8	85.4
Los Angeles	91.5	90.4	86.9	86.9	71.9	91.7
Memphis	93.8	94.1	100.0	90.9	42.6	97.0
Miami	100.0	93.3	93.1	95.3	88.5	90.7
New Orleans	92.7	92.7	89.2	85.4	47.8	88.7
Orange County	100.0	100.0	91.4	100.0	59.3	100.0
Philadelphia	89.8	91.2	83.8	86.2	57.9	87.6
San Diego	NA <sup>§</sup>	NA	NA	NA	NA	NA
<b>Local Median</b>	<b>93.8</b>	<b>93.3</b>	<b>91.4</b>	<b>87.1</b>	<b>57.9</b>	<b>90.7</b>
<b>Local Range</b>	<b>80.2 – 100.0</b>	<b>76.2 – 100.0</b>	<b>73.1 – 100.0</b>	<b>70.2 – 100.0</b>	<b>42.0 – 88.5</b>	<b>76.8 – 100.0</b>

\* HIV = human immunodeficiency virus.

\*\* AIDS = acquired immunodeficiency syndrome.

§ NA = data not available.

TABLE 9. Percentage of Schools That Taught Required HIV\* Prevention Units or Lessons in Specific Courses, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Science	Home economics or family and consumer education	Physical education	Family life education or life skills	Special education	Social studies
<b>STATE SURVEYS</b>						
Alaska	33.4	12.9	23.9	36.2	16.1	9.9
Arizona	35.8	10.0	21.5	25.5	11.7	8.3
Arkansas	42.4	47.4	31.2	51.4	30.7	13.5
Connecticut	38.8	16.0	20.1	44.2	21.3	7.2
Delaware	18.2	21.6	30.6	42.5	16.6	1.8
Idaho	28.7	23.1	14.5	24.5	10.3	3.0
Iowa	39.9	43.7	11.8	40.6	14.7	4.4
Maine	24.3	15.9	9.1	31.2	9.0	2.3
Massachusetts	27.3	14.8	17.9	22.1	12.3	4.1
Michigan	26.9	23.3	21.6	40.6	19.3	4.0
Missouri	39.4	50.6	45.0	39.7	24.6	6.2
Montana	33.9	28.7	73.5	40.1	16.0	6.9
Nebraska	37.3	46.2	36.3	43.1	13.4	6.7
New Hampshire	29.3	20.9	4.4	27.5	6.1	4.0
New York	48.8	20.1	19.6	38.5	28.5	11.8
North Carolina	27.9	21.4	61.9	42.6	16.8	6.2
North Dakota	36.4	45.3	31.2	52.9	12.1	9.0
Oklahoma	56.2	38.9	19.6	37.0	22.3	9.5
Oregon	27.0	11.6	24.6	24.7	17.9	9.1
Pennsylvania	26.4	19.2	36.1	27.3	20.8	6.4
South Carolina	39.4	20.7	47.9	32.1	21.2	8.8
Tennessee	41.6	29.4	44.7	44.8	19.7	6.9
Utah	25.4	34.1	12.6	47.0	18.7	4.5
Washington	41.5	26.4	27.1	33.3	17.9	11.3
Wisconsin	32.3	30.7	14.4	36.7	15.5	8.7
<b>State Median</b>	<b>33.9</b>	<b>23.1</b>	<b>23.9</b>	<b>38.5</b>	<b>16.8</b>	<b>6.9</b>
<b>State Range</b>	<b>18.2 – 56.2</b>	<b>10.0 – 50.6</b>	<b>4.4 – 73.5</b>	<b>22.1 – 52.9</b>	<b>6.1 – 30.7</b>	<b>1.8 – 13.5</b>
<b>LOCAL SURVEYS</b>						
Chicago	49.4	11.6	45.2	52.6	21.2	14.3
Dallas	47.8	40.9	30.4	40.9	19.2	6.5
District of Columbia	62.9	31.2	93.9	45.3	43.1	24.0
Los Angeles	53.2	9.0	7.8	27.9	39.5	4.4
Memphis	47.8	32.8	58.5	66.8	37.6	23.5
Miami	82.1	27.2	31.8	48.5	42.2	19.6
New Orleans	53.2	21.1	81.3	55.1	36.7	7.3
Orange County	65.7	15.9	17.3	57.4	42.2	8.0
Philadelphia	27.3	7.0	51.2	34.0	29.9	5.1
San Diego	69.3	2.4	12.5	41.9	12.8	20.7
<b>Local Median</b>	<b>53.2</b>	<b>18.5</b>	<b>38.5</b>	<b>46.9</b>	<b>37.2</b>	<b>11.2</b>
<b>Local Range</b>	<b>27.3 – 82.1</b>	<b>2.4 – 40.9</b>	<b>7.8 – 93.9</b>	<b>27.9 – 66.8</b>	<b>12.8 – 43.1</b>	<b>4.4 – 24.0</b>

\* HIV = human immunodeficiency virus.

TABLE 10a. Percentage of Schools That Taught About Choosing Healthful Foods in a Required Health Education Course During the 2003–2004 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Benefits of healthy eating	Using food labels	Identifying Food Guide Pyramid food groups and serving recommendations	Choosing a variety of grains daily*	Choosing a variety of fruits and vegetables daily	Choosing a low fat diet**	Choosing and preparing foods with less salt	Moderating intake of sugars	Choosing more calcium-rich foods
<b>STATE SURVEYS</b>									
Alaska	89.2	76.9	80.7	73.1	77.8	76.8	67.1	81.4	71.7
Arizona	97.0	94.9	90.6	89.3	91.3	87.7	79.3	87.5	80.5
Arkansas	97.2	90.4	92.1	86.7	90.8	89.8	82.7	89.2	84.2
Connecticut	96.4	87.2	88.6	84.2	88.0	88.2	72.6	86.9	76.5
Delaware	91.6	79.3	81.2	75.1	82.5	83.2	67.8	87.4	74.8
Idaho	98.8	95.7	96.3	92.0	95.7	96.3	85.9	97.0	89.6
Iowa	96.1	84.6	92.0	81.8	88.2	88.0	72.8	87.1	79.1
Maine	97.2	90.3	91.5	89.3	92.0	92.5	80.1	92.0	81.4
Massachusetts	94.4	86.8	87.1	85.3	89.0	88.5	76.3	88.2	81.8
Michigan	94.0	85.7	89.8	86.2	87.2	84.5	76.2	85.9	79.4
Missouri	98.4	96.1	96.9	94.9	97.1	96.4	86.3	94.5	87.2
Montana	98.0	83.2	90.2	85.8	89.8	87.7	73.7	90.3	81.2
Nebraska	97.5	83.5	91.2	84.1	86.6	87.1	76.6	85.1	76.1
New Hampshire	94.4	87.8	87.2	88.3	89.2	88.3	71.6	88.7	82.2
New York	97.3	87.2	87.5	86.8	91.0	93.6	79.4	91.7	82.2
North Carolina	97.9	94.5	94.5	90.5	94.6	93.9	90.1	93.1	87.4
North Dakota	97.5	90.8	93.8	92.5	94.3	91.3	81.2	91.6	82.4
Oklahoma	94.3	84.2	91.4	77.6	82.8	78.8	74.9	84.8	76.9
Oregon	98.4	93.9	91.9	89.7	93.1	92.0	78.8	91.6	78.9
Pennsylvania	92.7	83.7	83.6	80.8	86.3	88.0	76.4	87.2	76.7
South Carolina	98.3	92.9	90.9	86.4	93.1	94.5	84.6	91.8	86.9
Tennessee	97.3	93.1	95.7	91.8	94.2	94.2	86.3	92.3	88.3
Utah	99.4	93.2	94.8	91.9	94.9	95.5	86.9	92.5	84.2
Washington	96.8	87.1	88.9	85.5	89.5	89.8	72.7	86.5	75.4
Wisconsin	96.0	86.5	88.5	86.0	88.1	89.0	78.6	89.7	79.6
<b>State Median</b>	<b>97.2</b>	<b>87.2</b>	<b>90.9</b>	<b>86.4</b>	<b>89.8</b>	<b>89.0</b>	<b>78.6</b>	<b>89.2</b>	<b>81.2</b>
<b>State Range</b>	<b>89.2 – 99.4</b>	<b>76.9 – 96.1</b>	<b>80.7 – 96.9</b>	<b>73.1 – 94.9</b>	<b>77.8 – 97.1</b>	<b>76.8 – 96.4</b>	<b>67.1 – 90.1</b>	<b>81.4 – 97.0</b>	<b>71.7 – 89.6</b>
<b>LOCAL SURVEYS</b>									
Chicago	96.6	86.2	92.3	78.2	87.2	81.0	71.9	85.6	76.0
Dallas	100.0	100.0	100.0	100.0	100.0	100.0	96.0	100.0	100.0
District of Columbia	96.4	85.8	96.0	85.3	85.3	71.9	57.3	75.6	60.7
Los Angeles	98.8	97.6	96.4	92.6	96.3	91.4	83.3	95.2	86.9
Memphis	100.0	94.0	94.0	94.0	94.0	91.0	82.1	91.1	82.0
Miami	92.8	88.2	88.2	88.2	85.8	83.4	78.8	85.7	83.4
New Orleans	100.0	85.7	92.5	77.8	88.1	84.6	75.0	88.7	75.1
Orange County	100.0	96.3	96.3	100.0	100.0	100.0	92.0	100.0	95.7
Philadelphia	96.1	90.1	91.0	86.0	93.8	87.3	86.2	88.9	80.4
San Diego	NA <sup>§</sup>	NA	NA	NA	NA	NA	NA	NA	NA
<b>Local Median</b>	<b>98.8</b>	<b>90.1</b>	<b>94.0</b>	<b>88.2</b>	<b>93.8</b>	<b>87.3</b>	<b>82.1</b>	<b>88.9</b>	<b>82.0</b>
<b>Local Range</b>	<b>92.8 – 100.0</b>	<b>85.7 – 100.0</b>	<b>88.2 – 100.0</b>	<b>77.8 – 100.0</b>	<b>85.3 – 100.0</b>	<b>71.9 – 100.0</b>	<b>57.3 – 96.0</b>	<b>75.6 – 100.0</b>	<b>60.7 – 100.0</b>

\* Especially whole grains.

\*\* Low in saturated fat and cholesterol and moderate in total fat.

§ NA = data not available.

TABLE 10b. Percentage of Schools That Taught About Food Safety and Behaviors That Contribute to Maintaining a Healthy Weight in a Required Health Education Course During the 2003–2004 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Keeping food safe to eat	Preparing healthy meals and snacks	Aiming for a healthy weight	Risks of unhealthy weight control practices	Accepting body size differences	Eating disorders
<b>STATE SURVEYS</b>						
Alaska	73.4	76.7	77.7	78.3	74.0	77.8
Arizona	80.8	88.8	96.3	89.9	86.9	85.0
Arkansas	81.4	88.0	96.2	95.1	87.9	93.4
Connecticut	67.1	78.1	93.5	90.2	89.3	84.1
Delaware	61.4	71.9	81.6	87.7	77.5	85.8
Idaho	81.3	88.5	98.1	96.9	95.2	96.2
Iowa	73.7	79.0	89.2	90.5	87.3	87.7
Maine	74.9	81.4	96.0	93.1	89.6	91.4
Massachusetts	67.5	82.8	91.5	90.6	91.4	88.9
Michigan	74.5	82.2	91.2	90.4	87.3	87.3
Missouri	83.1	88.6	97.8	95.8	90.4	92.3
Montana	72.4	83.8	96.4	91.4	85.4	89.4
Nebraska	72.6	81.4	92.6	91.4	85.3	90.3
New Hampshire	67.3	81.5	89.2	89.4	89.5	90.7
New York	74.3	82.7	92.8	93.8	88.5	93.6
North Carolina	81.3	93.6	97.0	97.4	93.0	94.8
North Dakota	86.3	90.6	94.9	93.8	91.9	91.7
Oklahoma	77.3	76.3	89.9	86.0	80.7	82.9
Oregon	69.7	85.0	95.0	93.4	92.8	93.3
Pennsylvania	66.8	80.5	91.0	89.5	86.8	86.5
South Carolina	80.6	87.9	96.4	96.4	90.6	90.6
Tennessee	85.5	89.1	96.7	95.3	90.0	91.6
Utah	75.6	89.7	97.5	96.9	94.9	99.4
Washington	70.0	80.9	91.0	89.2	88.0	88.5
Wisconsin	69.9	78.5	92.7	91.3	89.4	93.8
<b>State Median</b>	<b>74.3</b>	<b>82.7</b>	<b>93.5</b>	<b>91.4</b>	<b>89.3</b>	<b>90.6</b>
<b>State Range</b>	<b>61.4 – 86.3</b>	<b>71.9 – 93.6</b>	<b>77.7 – 98.1</b>	<b>78.3 – 97.4</b>	<b>74.0 – 95.2</b>	<b>77.8 – 99.4</b>
<b>LOCAL SURVEYS</b>						
Chicago	77.5	81.3	90.4	84.8	81.4	74.4
Dallas	87.9	96.0	100.0	95.7	91.9	96.0
District of Columbia	69.8	81.7	86.1	78.2	85.1	76.4
Los Angeles	79.6	91.5	96.4	96.4	87.9	90.3
Memphis	88.1	94.0	97.0	97.0	88.1	97.0
Miami	81.1	90.5	90.4	90.4	88.1	88.2
New Orleans	80.9	84.8	88.7	84.6	75.1	84.3
Orange County	84.0	88.3	100.0	100.0	95.7	96.3
Philadelphia	85.1	88.8	93.7	91.0	86.2	87.4
San Diego	NA*	NA	NA	NA	NA	NA
<b>Local Median</b>	<b>81.1</b>	<b>88.8</b>	<b>93.7</b>	<b>91.0</b>	<b>87.9</b>	<b>88.2</b>
<b>Local Range</b>	<b>69.8 – 88.1</b>	<b>81.3 – 96.0</b>	<b>86.1 – 100.0</b>	<b>78.2 – 100.0</b>	<b>75.1 – 95.7</b>	<b>74.4 – 97.0</b>

\* NA = data not available.

TABLE 11a. Percentage of Schools That Taught About the Benefits of Physical Activity in a Required Health Education Course During the 2003–2004 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Physical, psychological, or social benefits	Health-related fitness	Phases of a workout	How much physical activity is enough	Decreasing sedentary activities
<b>STATE SURVEYS</b>					
Alaska	86.5	84.2	77.6	68.0	76.8
Arizona	96.2	91.8	85.9	88.1	87.3
Arkansas	97.9	95.3	90.9	87.0	90.5
Connecticut	90.3	78.9	63.9	67.0	79.1
Delaware	91.9	89.2	89.7	86.0	89.6
Idaho	95.2	94.7	87.1	89.5	90.1
Iowa	93.0	88.4	81.1	80.1	81.6
Maine	93.1	83.7	78.5	77.0	77.4
Massachusetts	93.2	85.8	77.3	77.5	84.4
Michigan	90.8	84.5	74.8	78.2	82.0
Missouri	94.7	94.5	89.5	90.0	90.7
Montana	97.6	98.4	95.7	91.2	86.7
Nebraska	94.4	92.0	89.6	86.1	85.2
New Hampshire	87.4	80.9	73.0	74.1	78.1
New York	94.4	81.7	70.0	74.4	84.2
North Carolina	96.6	97.1	95.0	90.7	92.3
North Dakota	96.2	93.8	91.9	87.2	92.4
Oklahoma	92.6	92.9	92.7	89.9	87.2
Oregon	96.1	92.2	86.9	84.5	83.5
Pennsylvania	94.2	90.6	84.0	82.3	84.6
South Carolina	96.5	93.3	86.6	86.8	91.9
Tennessee	96.5	95.5	94.6	90.5	89.9
Utah	97.0	92.1	87.5	86.3	91.9
Washington	94.4	92.6	80.4	77.5	82.9
Wisconsin	90.3	84.3	67.5	73.9	82.6
<b>State Median</b>	<b>94.4</b>	<b>92.0</b>	<b>86.6</b>	<b>86.0</b>	<b>85.2</b>
<b>State Range</b>	<b>86.5 – 97.9</b>	<b>78.9 – 98.4</b>	<b>63.9 – 95.7</b>	<b>67.0 – 91.2</b>	<b>76.8 – 92.4</b>
<b>LOCAL SURVEYS</b>					
Chicago	92.7	95.1	88.0	80.2	84.6
Dallas	100.0	87.4	87.0	82.6	87.4
District of Columbia	89.1	96.3	96.3	78.3	89.1
Los Angeles	91.4	86.6	73.4	79.3	86.6
Memphis	100.0	100.0	100.0	96.9	90.9
Miami	78.2	62.5	63.9	67.1	78.8
New Orleans	92.7	92.7	82.1	72.2	82.1
Orange County	95.8	87.2	81.9	77.9	95.1
Philadelphia	92.6	93.9	90.4	83.5	85.0
San Diego	NA*	NA	NA	NA	NA
<b>Local Median</b>	<b>92.7</b>	<b>92.7</b>	<b>87.0</b>	<b>79.3</b>	<b>86.6</b>
<b>Local Range</b>	<b>78.2 – 100.0</b>	<b>62.5 – 100.0</b>	<b>63.9 – 100.0</b>	<b>67.1 – 96.9</b>	<b>78.8 – 95.1</b>

\* NA = data not available.

Table 11b. Percentage of Schools That Taught About the Challenges to Engaging in Physical Activity in a Required Health Education Course During the 2003–2004 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Overcoming barriers to physical activity	Developing an individualized physical activity plan	Monitoring progress toward reaching goals	Opportunities for physical activity in the community	Preventing injury during physical activity	Weather-related safety	Dangers of using performance-enhancing drugs
<b>STATE SURVEYS</b>							
Alaska	63.5	54.1	55.0	72.9	76.3	77.6	69.5
Arizona	77.4	70.9	63.9	77.1	87.6	89.8	80.5
Arkansas	80.1	74.7	68.9	83.0	94.0	90.3	94.6
Connecticut	61.0	48.7	45.4	62.8	68.2	66.1	86.4
Delaware	73.0	71.4	62.6	78.4	87.3	79.2	97.7
Idaho	73.5	73.3	62.3	72.5	89.1	86.2	95.1
Iowa	65.7	64.2	59.9	70.1	76.1	77.9	87.8
Maine	67.4	69.1	57.4	74.0	77.2	73.2	83.8
Massachusetts	68.2	60.3	58.2	70.1	75.6	75.9	90.9
Michigan	64.6	62.9	60.3	66.2	74.1	73.8	85.9
Missouri	78.1	75.1	69.6	78.7	89.4	85.5	89.3
Montana	78.8	71.4	72.9	79.1	93.9	88.9	91.9
Nebraska	67.3	69.8	66.6	78.6	84.0	82.4	89.0
New Hampshire	69.5	58.7	55.2	71.3	71.2	72.6	85.3
New York	63.2	52.1	50.9	66.7	71.4	77.1	93.8
North Carolina	84.2	79.5	76.6	85.8	92.6	89.4	92.0
North Dakota	73.0	63.7	59.7	78.1	88.8	89.3	87.8
Oklahoma	77.0	74.3	74.1	76.9	91.1	88.5	92.7
Oregon	70.0	72.4	64.1	74.4	87.2	80.4	88.2
Pennsylvania	70.4	68.7	62.8	71.1	86.1	75.1	88.1
South Carolina	79.4	73.6	67.0	84.7	87.0	86.3	86.3
Tennessee	81.5	78.6	76.9	85.2	91.5	88.2	90.2
Utah	72.8	79.3	74.0	74.0	86.4	83.1	93.7
Washington	61.0	68.4	66.3	68.8	78.9	66.8	80.3
Wisconsin	69.5	55.6	51.8	69.0	75.3	74.1	88.6
<b>State Median</b>	<b>70.4</b>	<b>69.8</b>	<b>62.8</b>	<b>74.0</b>	<b>86.4</b>	<b>80.4</b>	<b>88.6</b>
<b>State Range</b>	<b>61.0 – 84.2</b>	<b>48.7 – 79.5</b>	<b>45.4 – 76.9</b>	<b>62.8 – 85.8</b>	<b>68.2 – 94.0</b>	<b>66.1 – 90.3</b>	<b>69.5 – 97.7</b>
<b>LOCAL SURVEYS</b>							
Chicago	79.4	61.1	63.1	79.7	89.6	75.3	75.6
Dallas	83.3	74.8	73.9	83.4	91.7	100.0	100.0
District of Columbia	74.9	67.8	67.8	85.6	92.5	82.4	85.8
Los Angeles	70.0	64.9	59.8	70.6	72.2	83.0	82.9
Memphis	94.0	87.9	82.0	94.0	100.0	87.9	94.0
Miami	64.8	53.3	55.6	64.7	62.4	71.7	78.2
New Orleans	78.4	55.1	62.0	79.2	85.4	85.0	89.1
Orange County	81.9	73.2	77.9	81.9	87.2	74.8	90.5
Philadelphia	72.3	75.5	71.9	76.0	89.0	77.3	80.0
San Diego	NA*	NA	NA	NA	NA	NA	NA
<b>Local Median</b>	<b>78.4</b>	<b>67.8</b>	<b>67.8</b>	<b>79.7</b>	<b>89.0</b>	<b>82.4</b>	<b>85.8</b>
<b>Local Range</b>	<b>64.8 – 94.0</b>	<b>53.3 – 87.9</b>	<b>55.6 – 82.0</b>	<b>64.7 – 94.0</b>	<b>62.4 – 100.0</b>	<b>71.7 – 100.0</b>	<b>75.6 – 100.0</b>

\* NA = data not available.

TABLE 12. Percentage of Schools That Taught All 17 Tobacco Use Prevention Topics, All 10 HIV\* Prevention Topics, All 15 Nutrition and Dietary Topics, or All 12 Physical Activity Topics in a Required Health Education Course During the 2003–2004 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Taught all 17 tobacco use prevention topics	Taught all 10 HIV prevention topics	Taught all 15 nutrition and dietary topics	Taught all 12 physical activity topics
<b>STATE SURVEYS</b>				
Alaska	50.8	29.6	50.6	39.2
Arizona	59.9	21.1	66.7	44.9
Arkansas	62.7	21.5	68.6	57.5
Connecticut	49.8	44.2	50.1	28.8
Delaware	61.4	50.0	51.7	53.2
Idaho	58.7	19.1	68.2	45.6
Iowa	54.8	38.6	55.7	44.2
Maine	43.3	43.5	56.1	41.6
Massachusetts	58.6	39.7	57.4	42.2
Michigan	56.8	25.7	61.1	42.9
Missouri	58.3	28.0	69.5	53.5
Montana	54.3	23.8	58.2	47.1
Nebraska	53.7	20.7	54.2	48.8
New Hampshire	51.0	48.6	53.6	37.2
New York	63.3	51.0	57.7	34.6
North Carolina	69.0	14.9	73.4	59.5
North Dakota	63.3	20.2	71.6	44.3
Oklahoma	69.5	23.5	65.3	60.1
Oregon	56.4	41.7	56.5	43.4
Pennsylvania	59.8	36.8	55.4	47.6
South Carolina	58.4	34.9	67.5	53.9
Tennessee	64.6	34.7	73.1	62.7
Utah	54.3	7.0	64.3	48.2
Washington	50.0	38.6	55.3	38.3
Wisconsin	50.6	46.4	57.0	35.4
<b>State Median</b>	<b>58.3</b>	<b>34.7</b>	<b>57.7</b>	<b>44.9</b>
<b>State Range</b>	<b>43.3 – 69.5</b>	<b>7.0 – 51.0</b>	<b>50.1 – 73.4</b>	<b>28.8 – 62.7</b>
<b>LOCAL SURVEYS</b>				
Chicago	52.4	38.5	51.8	43.7
Dallas	78.2	43.7	83.9	43.5
District of Columbia	39.3	71.5	42.7	50.2
Los Angeles	79.4	61.6	67.7	48.2
Memphis	69.8	42.6	76.0	67.0
Miami	66.9	79.0	71.8	48.6
New Orleans	71.5	42.4	50.1	48.4
Orange County	67.9	54.9	69.1	60.4
Philadelphia	63.3	52.0	67.8	51.8
San Diego	NA**	NA	NA	NA
<b>Local Median</b>	<b>67.9</b>	<b>52.0</b>	<b>67.8</b>	<b>48.6</b>
<b>Local Range</b>	<b>39.3 – 79.4</b>	<b>38.5 – 79.0</b>	<b>42.7 – 83.9</b>	<b>43.5 – 67.0</b>

\* HIV = human immunodeficiency virus.  
 \*\* NA = data not available.

TABLE 13. Percentage of Schools That Used Specific Teaching Methods in a Required Health Education Course During the 2003–2004 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Group discussions	Cooperative group activities	Role play, simulations, or practice	Language, performing, or visual arts	Pledges or contracts for behavior change	Peer educators	The Internet	Computer-assisted instruction
<b>STATE SURVEYS</b>								
Alaska	90.0	82.5	69.8	63.0	48.9	59.8	79.2	49.6
Arizona	92.6	91.3	72.0	65.0	50.3	58.5	78.4	52.2
Arkansas	98.9	91.4	72.9	51.8	40.0	64.5	76.1	49.4
Connecticut	100.0	98.8	87.8	68.5	43.9	55.5	91.5	67.1
Delaware	97.9	97.9	95.7	77.3	48.1	56.2	86.0	57.9
Idaho	98.6	98.7	79.4	69.6	40.0	63.4	85.8	57.4
Iowa	98.9	97.1	80.8	61.3	42.7	54.9	87.1	51.8
Maine	99.4	95.5	85.1	60.0	35.7	52.0	95.0	58.4
Massachusetts	99.6	97.4	87.6	73.9	42.7	54.2	83.2	57.3
Michigan	98.0	96.0	84.3	68.9	46.8	60.0	82.3	59.9
Missouri	97.9	96.4	77.3	70.3	44.0	65.7	82.7	58.4
Montana	99.6	94.4	72.4	67.2	40.8	70.6	87.3	63.8
Nebraska	99.0	94.1	74.0	56.1	41.1	53.0	83.6	59.0
New Hampshire	97.5	99.2	91.7	79.3	33.9	59.3	80.7	51.0
New York	98.6	98.5	89.9	73.8	55.8	62.1	91.1	65.3
North Carolina	97.1	94.4	82.3	73.0	57.4	70.2	79.2	66.5
North Dakota	100.0	96.3	82.7	65.5	42.6	64.8	93.0	64.6
Oklahoma	91.4	86.0	55.0	56.4	41.5	53.3	84.7	63.8
Oregon	98.8	98.2	87.5	68.5	56.0	63.7	86.1	55.2
Pennsylvania	98.4	97.1	87.0	70.0	47.6	65.2	82.4	60.4
South Carolina	98.8	96.5	77.5	63.4	57.7	65.3	81.3	61.8
Tennessee	99.0	94.9	82.4	61.6	58.0	68.0	73.9	55.7
Utah	99.4	97.7	90.0	70.8	50.4	68.0	79.0	55.0
Washington	93.7	90.6	77.6	68.8	46.3	59.3	82.8	43.5
Wisconsin	98.6	95.4	82.4	68.6	46.2	61.6	86.6	60.7
<b>State Median</b>	<b>98.6</b>	<b>96.3</b>	<b>82.4</b>	<b>68.5</b>	<b>46.2</b>	<b>61.6</b>	<b>83.2</b>	<b>58.4</b>
<b>State Range</b>	<b>90.0 – 100.0</b>	<b>82.5 – 99.2</b>	<b>55.0 – 95.7</b>	<b>51.8 – 79.3</b>	<b>33.9 – 58.0</b>	<b>52.0 – 70.6</b>	<b>73.9 – 95.0</b>	<b>43.5 – 67.1</b>
<b>LOCAL SURVEYS</b>								
Chicago	97.7	90.9	77.6	75.8	57.1	57.8	66.9	54.9
Dallas	100.0	100.0	95.9	70.5	54.3	83.3	95.9	65.1
District of Columbia	100.0	100.0	92.9	71.5	60.7	85.4	82.0	64.4
Los Angeles	97.6	95.2	89.4	79.8	67.7	61.5	84.5	61.0
Memphis	100.0	100.0	100.0	84.8	71.7	85.0	85.2	73.0
Miami	100.0	91.0	91.0	72.6	65.7	72.6	81.7	55.7
New Orleans	96.5	96.5	93.0	78.0	54.1	68.1	68.8	55.4
Orange County	96.3	92.0	88.3	96.3	43.8	71.0	87.7	59.3
Philadelphia	94.9	97.7	87.1	75.7	47.2	64.8	72.5	57.4
San Diego	NA*	NA	NA	NA	NA	NA	NA	NA
<b>Local Median</b>	<b>97.7</b>	<b>96.5</b>	<b>91.0</b>	<b>75.8</b>	<b>57.1</b>	<b>71.0</b>	<b>82.0</b>	<b>59.3</b>
<b>Local Range</b>	<b>94.9 – 100.0</b>	<b>90.9 – 100.0</b>	<b>77.6 – 100.0</b>	<b>70.5 – 96.3</b>	<b>43.8 – 71.7</b>	<b>57.8 – 85.4</b>	<b>66.9 – 95.9</b>	<b>54.9 – 73.0</b>

\* NA = data not available.

TABLE 14. Percentage of Schools That Used Teaching Methods to Highlight Diversity or the Values of Various Cultures in a Required Health Education Course During the 2003–2004 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Using textbooks or curricular materials reflective of various cultures	Using textbooks or curricular materials designed for students with limited English proficiency	Asking students to share their own cultural experiences related to health topics	Teaching about cultural differences and similarities	Modifying teaching methods to match students' learning styles, health beliefs, or cultural values
<b>STATE SURVEYS</b>					
Alaska	71.4	32.3	79.3	79.0	90.1
Arizona	66.9	40.2	72.0	78.6	82.1
Arkansas	67.2	31.9	72.1	81.5	87.3
Connecticut	53.1	22.3	72.9	73.6	92.3
Delaware	66.6	21.3	83.5	74.6	94.2
Idaho	64.2	30.6	75.0	76.6	85.5
Iowa	71.2	20.3	66.8	68.0	84.9
Maine	48.5	16.3	62.4	67.3	88.5
Massachusetts	61.6	18.6	78.8	78.3	92.2
Michigan	61.0	19.5	65.3	68.7	83.4
Missouri	75.6	24.8	80.0	79.4	88.5
Montana	56.7	12.9	60.2	66.8	85.2
Nebraska	76.6	20.4	71.6	77.1	87.3
New Hampshire	56.8	18.4	60.9	70.2	89.1
New York	63.6	21.5	79.7	78.0	93.0
North Carolina	69.0	43.2	81.1	76.9	90.3
North Dakota	65.8	15.0	57.3	74.5	83.4
Oklahoma	64.3	29.0	52.7	66.7	73.3
Oregon	69.1	34.9	71.2	73.7	88.0
Pennsylvania	65.4	20.0	69.0	70.7	88.1
South Carolina	75.1	29.9	80.9	78.5	92.1
Tennessee	72.0	32.9	75.0	82.0	88.7
Utah	69.2	41.4	71.2	79.4	89.7
Washington	71.9	23.6	64.6	66.3	89.2
Wisconsin	61.9	22.2	66.6	70.3	91.2
<b>State Median</b>	<b>66.6</b>	<b>22.3</b>	<b>71.6</b>	<b>74.6</b>	<b>88.5</b>
<b>State Range</b>	<b>48.5 – 76.6</b>	<b>12.9 – 43.2</b>	<b>52.7 – 83.5</b>	<b>66.3 – 82.0</b>	<b>73.3 – 94.2</b>
<b>LOCAL SURVEYS</b>					
Chicago	76.4	42.1	84.0	87.7	85.4
Dallas	100.0	74.8	87.9	79.8	100.0
District of Columbia	67.4	50.2	82.0	86.1	92.9
Los Angeles	86.9	78.7	89.5	88.4	90.4
Memphis	96.9	50.0	94.1	97.0	96.9
Miami	83.9	65.8	88.5	83.9	86.3
New Orleans	68.9	20.7	82.7	75.9	72.7
Orange County	64.2	59.9	83.3	91.4	96.3
Philadelphia	75.0	33.8	75.9	84.7	87.7
San Diego	NA*	NA	NA	NA	NA
<b>Local Median</b>	<b>76.4</b>	<b>50.2</b>	<b>84.0</b>	<b>86.1</b>	<b>90.4</b>
<b>Local Range</b>	<b>64.2 – 100.0</b>	<b>20.7 – 78.7</b>	<b>75.9 – 94.1</b>	<b>75.9 – 97.0</b>	<b>72.7 – 100.0</b>

\* NA = data not available.

TABLE 15. Percentage of Schools With a Health Education Coordinator and, Among those Schools, the Percentage in Which Specific Staff Serve as the Health Education Coordinator, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2004

Site	Have a health education coordinator	District administrator or district health education or curriculum coordinator	School administrator	Health education teacher	School nurse	Someone else
<b>STATE SURVEYS</b>						
Alaska	87.8	21.8	32.5	31.6	1.5	12.6
Arizona	86.9	22.7	25.3	25.5	14.0	12.5
Arkansas	98.2	20.1	25.3	53.1	0.0	1.5
Connecticut	92.2	37.8	18.9	35.3	1.7	6.2
Delaware	100.0	26.6	15.9	54.1	1.7	1.8
Idaho	99.5	26.1	16.7	56.6	0.0	0.5
Iowa	96.9	30.2	17.3	46.9	1.6	4.0
Maine	95.3	34.2	9.3	51.6	2.7	2.2
Massachusetts	93.2	49.0	12.7	29.2	3.6	5.6
Michigan	94.5	39.2	15.9	39.9	1.5	3.5
Minnesota	98.3	24.9	16.1	54.0	0.5	4.5
Missouri	97.4	32.9	18.5	43.5	3.9	1.3
Montana	97.0	23.2	15.0	59.3	0.3	2.1
Nebraska	96.3	24.2	19.4	49.6	2.6	4.1
New Hampshire	93.3	15.7	18.9	44.3	14.1	7.0
New York	97.5	39.3	23.9	32.9	1.7	2.3
North Carolina	95.1	31.0	10.4	53.0	1.1	4.5
North Dakota	94.9	20.9	30.0	46.5	1.6	1.0
Oklahoma	74.2	16.9	33.7	20.7	14.9	13.9
Oregon	97.4	21.0	22.7	52.6	0.4	3.3
Pennsylvania	98.1	36.7	18.1	40.5	0.6	4.1
South Carolina	97.0	25.1	21.2	43.2	2.6	7.9
Tennessee	90.8	22.1	29.2	35.9	5.4	7.3
Texas*	93.6	27.0	25.0	36.3	5.4	6.3
Utah	100.0	18.6	20.6	59.8	0.5	0.5
Washington	95.2	25.1	15.3	46.3	3.2	10.0
Wisconsin	96.6	27.0	12.5	56.4	0.6	3.5
<b>State Median</b>	<b>96.3</b>	<b>25.1</b>	<b>18.9</b>	<b>46.3</b>	<b>1.7</b>	<b>4.1</b>
<b>State Range</b>	<b>74.2 – 100.0</b>	<b>15.7 – 49.0</b>	<b>9.3 – 33.7</b>	<b>20.7 – 59.8</b>	<b>0.0 – 14.9</b>	<b>0.5 – 13.9</b>
<b>LOCAL SURVEYS</b>						
Chicago	83.7	1.9	17.6	43.8	6.8	29.9
Dallas	81.4	39.6	10.0	32.5	2.6	15.2
District of Columbia	93.6	13.7	6.8	76.4	3.1	0.0
Los Angeles	95.6	11.1	28.5	52.8	0.0	7.6
Memphis	94.3	22.4	23.7	39.9	1.9	12.0
Miami	91.7	25.4	25.2	30.1	1.2	18.1
New Orleans	96.9	16.0	0.0	71.0	9.7	3.3
Orange County	97.3	15.4	33.4	34.2	0.0	17.0
Philadelphia	94.9	10.7	19.6	56.1	3.7	9.9
San Bernardino	85.7	50.0	33.3	8.3	0.0	8.3
San Diego	95.0	48.9	7.7	0.0	0.0	43.4
<b>Local Median</b>	<b>94.3</b>	<b>16.0</b>	<b>19.6</b>	<b>39.9</b>	<b>1.9</b>	<b>12.0</b>
<b>Local Range</b>	<b>81.4 – 97.3</b>	<b>1.9 – 50.0</b>	<b>0.0 – 33.4</b>	<b>0.0 – 76.4</b>	<b>0.0 – 9.7</b>	<b>0.0 – 43.4</b>

\* Survey did not include schools from two of the state's largest school districts.

TABLE 16. Percentage of Schools in Which Health Education Staff Worked on Health Education Activities With Other School Staff or Community Members During the 2003–2004 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Physical education staff	Teachers in other subject areas	School health services staff	School mental health or social services staff	Food service staff	Community members
<b>STATE SURVEYS</b>						
Alaska	51.1	64.7	48.7	50.8	15.8	56.2
Arizona	55.1	57.6	50.6	55.7	22.9	36.0
Arkansas	79.6	62.6	72.1	53.9	20.4	46.1
Connecticut	70.3	60.0	70.9	69.8	24.1	58.8
Delaware	91.7	65.0	83.5	69.2	25.0	58.2
Idaho	80.7	62.2	61.4	60.2	20.5	55.4
Iowa	63.2	62.5	67.9	51.9	19.8	55.0
Maine	80.8	60.9	82.5	64.4	31.1	53.4
Massachusetts	76.9	63.6	78.5	69.5	27.4	56.7
Michigan	65.5	53.9	34.9	50.2	23.3	46.7
Missouri	81.7	59.8	78.9	58.4	26.9	45.6
Montana	81.9	60.0	55.9	58.7	28.4	55.7
Nebraska	72.4	57.8	67.5	43.6	18.4	41.5
New Hampshire	68.0	62.8	80.7	72.1	30.0	60.2
New York	70.3	71.9	64.5	67.3	21.9	62.9
North Carolina	87.2	58.7	72.5	66.5	24.9	55.2
North Dakota	73.1	62.2	37.6	59.8	35.0	54.6
Oklahoma	51.3	47.1	38.9	39.5	26.3	26.1
Oregon	72.0	63.4	56.5	60.7	16.8	54.7
Pennsylvania	87.7	60.6	79.5	61.0	20.3	50.7
South Carolina	77.2	55.5	67.9	53.5	23.4	43.0
Tennessee	74.0	61.0	65.5	60.5	31.2	52.3
Utah	75.1	65.7	48.1	67.7	11.6	66.5
Washington	64.3	55.4	65.5	62.8	18.0	46.8
Wisconsin	76.5	64.1	69.2	62.6	24.6	59.4
<b>State Median</b>	<b>74.0</b>	<b>61.0</b>	<b>67.5</b>	<b>60.5</b>	<b>23.4</b>	<b>54.7</b>
<b>State Range</b>	<b>51.1 – 91.7</b>	<b>47.1 – 71.9</b>	<b>34.9 – 83.5</b>	<b>39.5 – 72.1</b>	<b>11.6 – 35.0</b>	<b>26.1 – 66.5</b>
<b>LOCAL SURVEYS</b>						
Chicago	66.4	61.5	53.6	57.4	25.6	33.9
Dallas	50.9	54.9	74.5	54.1	19.9	22.0
District of Columbia	86.9	74.1	83.2	77.4	37.5	77.4
Los Angeles	51.6	68.1	69.1	56.5	24.5	54.7
Memphis	88.2	74.0	82.2	68.0	36.0	64.1
Miami	47.8	45.6	28.2	50.6	22.1	41.7
New Orleans	94.0	61.5	93.8	78.0	53.4	54.9
Orange County	60.6	63.6	48.2	60.0	15.9	68.5
Philadelphia	87.3	68.9	77.5	56.6	34.5	34.8
San Diego	46.4	83.6	62.5	82.7	5.1	27.2
<b>Local Median</b>	<b>63.5</b>	<b>65.9</b>	<b>71.8</b>	<b>58.7</b>	<b>25.1</b>	<b>48.2</b>
<b>Local Range</b>	<b>46.4 – 94.0</b>	<b>45.6 – 83.6</b>	<b>28.2 – 93.8</b>	<b>50.6 – 82.7</b>	<b>5.1 – 53.4</b>	<b>22.0 – 77.4</b>

TABLE 17. Percentage of Schools in Which the Lead Health Education Teacher Had Professional Preparation in Each Specific Discipline, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Health and physical education combined	Health education only	Physical education only	Other education degree	Kinesiology, exercise science or exercise physiology, home economics or family and consumer science, or science	Nursing or counseling	Public health or other discipline
<b>STATE SURVEYS*</b>							
Alaska	10.3	2.7	7.4	28.4	19.8	10.6	20.9
Arizona	19.5	6.1	9.3	22.8	10.9	19.0	12.4
Arkansas	68.5	8.0	14.0	2.3	5.6	0.6	1.1
Connecticut	48.6	20.6	9.1	4.5	8.5	4.7	4.1
Delaware	77.1	9.6	9.6	0.0	0.0	1.7	2.0
Idaho	61.1	5.9	15.6	7.8	7.1	0.6	1.8
Iowa	37.2	6.4	14.1	4.5	32.8	4.6	0.4
Maine	39.5	22.6	10.9	5.6	15.3	5.0	1.0
Massachusetts	47.6	20.1	7.0	3.1	8.5	9.9	3.7
Michigan	33.2	7.9	17.3	9.6	25.7	2.2	4.1
Missouri	57.2	1.8	16.7	3.8	16.4	2.6	1.4
Montana	70.7	1.1	10.3	11.8	2.5	0.8	2.9
Nebraska	41.0	2.1	27.9	4.0	19.0	3.9	2.1
New Hampshire	19.9	18.6	17.4	3.5	17.4	17.1	6.1
New York	40.7	40.5	8.5	1.9	4.8	2.6	1.1
North Carolina	58.8	6.6	23.6	1.0	4.7	3.4	1.8
North Dakota	40.1	4.0	13.3	5.9	24.4	4.0	8.2
Oklahoma	38.7	1.4	6.8	11.8	15.6	17.4	8.3
Oregon	46.0	16.2	7.8	11.4	12.2	2.1	4.4
Pennsylvania	86.8	3.3	4.2	1.1	2.3	0.4	1.8
South Carolina	45.1	6.3	28.5	4.0	6.1	7.6	2.3
Tennessee	56.5	2.3	11.2	10.0	7.6	10.3	2.1
Utah	45.1	16.6	20.6	6.6	8.0	1.3	1.7
Washington	22.5	5.6	15.2	15.5	21.1	7.6	12.4
Wisconsin	55.2	8.9	13.8	7.8	10.6	2.4	1.3
<b>State Median</b>	<b>45.1</b>	<b>6.4</b>	<b>13.3</b>	<b>5.6</b>	<b>10.6</b>	<b>3.9</b>	<b>2.1</b>
<b>State Range</b>	<b>10.3 – 86.8</b>	<b>1.1 – 40.5</b>	<b>4.2 – 28.5</b>	<b>0.0 – 28.4</b>	<b>0.0 – 32.8</b>	<b>0.4 – 19.0</b>	<b>0.4 – 20.9</b>
<b>LOCAL SURVEYS*</b>							
Chicago	32.1	4.3	22.0	9.4	10.6	14.7	6.9
Dallas	39.6	8.7	4.4	0.0	47.3	0.0	0.0
District of Columbia	73.9	3.9	3.9	0.0	11.3	3.5	3.5
Los Angeles	9.8	30.6	1.3	7.7	45.5	1.3	3.8
Memphis	63.2	4.5	6.8	4.5	9.1	9.4	2.4
Miami	24.8	13.7	12.4	5.0	34.1	5.0	5.0
New Orleans	72.5	0.0	3.4	6.9	3.4	13.9	0.0
Orange County	24.4	26.8	15.9	8.1	21.9	0.0	2.8
Philadelphia	84.6	2.0	4.9	2.1	0.9	4.3	1.1
San Diego	2.8	0.0	2.6	0.0	2.8	91.9	0.0
<b>Local Median</b>	<b>35.9</b>	<b>4.4</b>	<b>4.7</b>	<b>4.8</b>	<b>11.0</b>	<b>4.7</b>	<b>2.6</b>
<b>Local Range</b>	<b>2.8 – 84.6</b>	<b>0.0 – 30.6</b>	<b>1.3 – 22.0</b>	<b>0.0 – 9.4</b>	<b>0.9 – 47.3</b>	<b>0.0 – 91.9</b>	<b>0.0 – 6.9</b>

\* Percentages for each row might not add up to 100.0 because of rounding.

TABLE 18. Percentage of Schools in Which a Newly Hired Health Education Teacher Is Required to Be Certified\* in Health Education, the Lead Health Education Teacher Holds State Certification\* in Health Education, and the Lead Health Education Teacher Had Taught Health Education for a Specific Number of Years, Selected U.S. Sites: School Health Profiles, Principal Surveys and Lead Health Education Teacher Surveys, 2004

Site	Newly hired health education teacher required to be certified in health education	Lead health education teacher holds current teaching certificate in health education	Years Lead Health Education Teacher Taught Health Education				
			1 year	2 to 5 years	6 to 9 years	10 to 14 years	15 years or more
<b>STATE SURVEYS</b>							
Alaska	31.8	32.9	18.7	33.8	14.8	11.1	21.6
Arizona	50.8	32.1	16.8	32.3	18.7	11.1	21.1
Arkansas	98.0	90.2	10.5	24.2	13.6	14.5	37.2
Connecticut	94.1	81.3	6.6	22.9	16.4	16.8	37.2
Delaware	87.3	89.6	3.4	29.5	22.3	15.9	28.9
Idaho	89.5	89.7	7.9	20.9	15.9	16.1	39.1
Iowa	84.3	80.7	6.6	27.4	15.8	16.4	33.9
Maine	91.2	85.5	8.2	24.2	13.0	19.5	35.0
Massachusetts	92.6	80.0	5.0	22.8	16.6	18.7	36.8
Michigan	84.5	71.9	9.5	28.5	17.0	13.6	31.5
Minnesota	98.3	NA**	NA	NA	NA	NA	NA
Missouri	87.5	83.3	10.8	32.0	22.3	13.2	21.7
Montana	83.9	91.6	6.0	25.7	14.6	16.9	36.8
Nebraska	57.0	61.5	5.7	21.9	21.7	14.0	36.8
New Hampshire	83.1	49.2	7.8	29.1	18.5	15.9	28.8
New York	96.0	88.8	5.7	24.2	16.9	9.4	43.9
North Carolina	82.3	71.3	2.5	26.1	12.1	18.9	40.3
North Dakota	85.4	NA	7.2	18.1	18.5	19.0	37.2
Oklahoma	75.6	72.3	13.3	23.7	16.6	19.0	27.4
Oregon	82.1	77.2	9.7	20.4	19.0	14.8	36.1
Pennsylvania	94.0	94.7	5.1	13.2	12.2	13.1	56.4
South Carolina	71.2	56.6	8.1	25.8	14.2	14.6	37.2
Tennessee	81.8	75.1	8.6	25.4	14.7	14.7	36.6
Texas <sup>§</sup>	90.7	NA	NA	NA	NA	NA	NA
Utah	98.5	89.3	8.9	20.8	17.9	17.8	34.5
Washington	67.6	58.5	10.2	34.7	13.3	14.7	27.2
Wisconsin	90.0	85.8	5.9	15.0	17.5	15.3	46.4
<b>State Median</b>	<b>85.4</b>	<b>80.4</b>	<b>7.9</b>	<b>24.2</b>	<b>16.6</b>	<b>15.3</b>	<b>36.6</b>
<b>State Range</b>	<b>31.8 – 98.5</b>	<b>32.1 – 94.7</b>	<b>2.5 – 18.7</b>	<b>13.2 – 34.7</b>	<b>12.1 – 22.3</b>	<b>9.4 – 19.5</b>	<b>21.1 – 56.4</b>
<b>LOCAL SURVEYS</b>							
Chicago	63.6	50.1	9.4	23.7	14.9	16.5	35.4
Dallas	91.6	63.5	6.9	25.2	25.1	15.9	27.0
District of Columbia	97.2	97.0	0.0	19.8	13.1	6.4	60.6
Los Angeles	90.1	70.9	6.5	37.7	17.2	11.7	26.9
Memphis	83.3	73.3	6.5	26.4	13.2	11.2	42.7
Miami	86.3	58.3	9.9	26.0	18.5	21.1	24.6
New Orleans	96.8	93.6	6.2	9.0	3.0	12.1	69.8
Orange County	68.5	61.5	13.2	28.7	20.5	10.5	27.1
Philadelphia	85.4	95.2	5.8	18.8	5.6	10.8	59.0
San Bernardino	86.7	NA	NA	NA	NA	NA	NA
San Diego	80.1	39.8	63.5	14.5	0.0	2.5	19.5
<b>Local Median</b>	<b>86.3</b>	<b>67.2</b>	<b>6.7</b>	<b>24.5</b>	<b>14.1</b>	<b>11.5</b>	<b>31.3</b>
<b>Local Range</b>	<b>63.6 – 97.2</b>	<b>39.8 – 97.0</b>	<b>0.0 – 63.5</b>	<b>9.0 – 37.7</b>	<b>0.0 – 25.1</b>	<b>2.5 – 21.1</b>	<b>19.5 – 69.8</b>

\* Certification, licensure, or endorsement by the state.

\*\* NA = data not available.

§ Survey did not include schools from two of the state's largest school districts.

TABLE 19a. Percentage of Schools in Which the Lead Health Education Teacher Received Staff Development During the Two Years Preceding the Survey on Specific Health Education Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Accident or injury prevention	Alcohol or other drug use prevention	Consumer health	CPR*	Death and dying	Dental and oral health	Emotional and mental health
<b>STATE SURVEYS</b>							
Alaska	45.4	43.4	18.8	41.9	20.0	16.4	38.0
Arizona	50.8	50.0	16.6	59.2	16.5	18.7	38.3
Arkansas	40.6	46.8	13.4	59.0	11.2	9.4	24.4
Connecticut	28.7	46.3	8.6	55.7	7.7	5.7	38.6
Delaware	33.8	66.2	18.2	64.6	10.7	9.6	46.6
Idaho	33.5	60.7	14.8	45.1	13.9	8.7	50.2
Iowa	33.2	38.7	16.5	47.4	11.0	6.7	31.4
Maine	27.4	38.9	11.4	45.8	10.3	5.3	32.3
Massachusetts	30.7	50.4	15.1	64.2	12.6	7.9	43.0
Michigan	32.3	43.2	18.1	54.1	10.2	10.3	29.8
Missouri	42.5	43.0	17.4	70.2	13.2	14.4	29.7
Montana	52.0	53.7	19.0	72.8	20.5	14.5	31.9
Nebraska	35.5	37.7	11.2	65.4	16.6	4.8	19.9
New Hampshire	32.9	56.0	17.2	59.1	14.7	11.5	50.2
New York	39.0	45.7	15.0	67.0	12.3	4.8	35.7
North Carolina	41.3	36.0	10.3	61.4	6.0	5.8	21.8
North Dakota	41.5	65.7	21.6	56.2	13.5	17.7	38.2
Oklahoma	60.5	53.2	14.3	65.8	10.4	18.7	30.4
Oregon	38.7	48.2	9.2	69.4	9.7	7.9	28.0
Pennsylvania	35.8	57.4	12.9	72.4	9.7	6.0	29.6
South Carolina	44.4	35.3	17.0	62.9	12.0	15.3	26.0
Tennessee	59.3	50.5	21.1	74.8	17.2	19.5	32.6
Utah	35.4	48.7	11.9	64.4	10.1	4.6	32.6
Washington	37.9	34.4	11.2	77.5	11.0	9.8	31.0
Wisconsin	36.5	51.4	16.7	66.5	9.8	5.2	34.4
<b>State Median</b>	<b>37.9</b>	<b>48.2</b>	<b>15.1</b>	<b>64.2</b>	<b>11.2</b>	<b>9.4</b>	<b>32.3</b>
<b>State Range</b>	<b>27.4 – 60.5</b>	<b>34.4 – 66.2</b>	<b>8.6 – 21.6</b>	<b>41.9 – 77.5</b>	<b>6.0 – 20.5</b>	<b>4.6 – 19.5</b>	<b>19.9 – 50.2</b>
<b>LOCAL SURVEYS</b>							
Chicago	42.3	45.4	26.4	45.9	16.4	25.6	36.0
Dallas	49.8	56.7	24.2	58.4	14.4	8.4	36.4
District of Columbia	48.2	69.4	26.4	51.2	35.0	30.0	44.8
Los Angeles	36.9	61.7	15.6	41.9	5.3	7.3	26.7
Memphis	82.4	61.6	39.0	88.2	28.9	28.3	53.4
Miami	22.9	29.7	9.8	53.5	6.4	7.5	17.1
New Orleans	39.5	53.2	16.2	62.5	22.0	33.3	50.2
Orange County	30.2	37.7	12.8	55.1	8.4	0.0	28.3
Philadelphia	34.6	41.1	17.4	48.0	7.5	9.6	24.3
San Diego	33.3	92.6	24.5	34.3	38.9	4.9	65.8
<b>Local Median</b>	<b>38.2</b>	<b>55.0</b>	<b>20.8</b>	<b>52.4</b>	<b>15.4</b>	<b>9.0</b>	<b>36.2</b>
<b>Local Range</b>	<b>22.9 – 82.4</b>	<b>29.7 – 92.6</b>	<b>9.8 – 39.0</b>	<b>34.3 – 88.2</b>	<b>5.3 – 38.9</b>	<b>0.0 – 33.3</b>	<b>17.1 – 65.8</b>

\* CPR = cardiopulmonary resuscitation.

TABLE 19b. Percentage of Schools in Which the Lead Health Education Teacher Received Staff Development During the Two Years Preceding the Survey on Specific Health Education Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Environmental health	First aid	Growth and development	HIV* prevention	Human sexuality	Immunization and vaccinations	Nutrition and dietary behavior
<b>STATE SURVEYS</b>							
Alaska	22.0	53.3	23.9	33.7	18.0	25.3	22.4
Arizona	23.0	55.5	28.7	35.1	22.0	28.4	33.1
Arkansas	16.4	53.9	15.7	25.0	16.3	11.4	27.1
Connecticut	9.6	51.1	23.0	41.7	29.4	10.4	30.2
Delaware	20.2	46.9	31.1	48.7	43.1	12.4	32.0
Idaho	12.2	46.7	15.6	55.9	40.7	12.6	48.1
Iowa	14.9	36.9	23.2	35.3	25.6	12.2	26.6
Maine	8.1	43.6	16.8	51.2	34.4	9.2	36.3
Massachusetts	10.7	48.1	25.4	38.0	36.6	12.7	38.7
Michigan	15.1	45.9	26.9	55.1	45.0	15.5	35.1
Missouri	18.4	59.9	21.7	35.9	24.0	20.1	35.8
Montana	15.8	72.5	21.5	51.7	29.5	17.0	37.6
Nebraska	9.9	47.0	15.8	21.8	18.6	15.7	22.5
New Hampshire	13.8	48.0	34.3	42.7	43.4	13.8	46.4
New York	18.3	57.3	26.2	50.0	35.9	17.8	29.8
North Carolina	12.3	55.7	19.2	38.9	27.1	11.7	23.2
North Dakota	18.6	44.2	25.9	46.0	22.2	24.6	35.5
Oklahoma	21.5	65.4	25.4	77.4	20.0	31.4	23.1
Oregon	9.1	71.3	16.7	43.7	30.1	16.9	23.5
Pennsylvania	12.0	52.7	19.4	40.3	29.3	11.9	35.7
South Carolina	20.1	56.7	25.7	45.4	37.2	21.8	26.8
Tennessee	22.0	66.1	26.1	48.0	24.4	28.1	37.7
Utah	4.9	55.8	19.2	35.0	32.0	11.5	21.4
Washington	14.7	78.2	21.4	56.9	30.7	16.5	26.9
Wisconsin	10.1	55.4	34.4	39.4	31.3	14.6	34.9
<b>State Median</b>	<b>14.9</b>	<b>53.9</b>	<b>23.2</b>	<b>42.7</b>	<b>29.5</b>	<b>15.5</b>	<b>32.0</b>
<b>State Range</b>	<b>4.9 – 23.0</b>	<b>36.9 – 78.2</b>	<b>15.6 – 34.4</b>	<b>21.8 – 77.4</b>	<b>16.3 – 45.0</b>	<b>9.2 – 31.4</b>	<b>21.4 – 48.1</b>
<b>LOCAL SURVEYS</b>							
Chicago	28.3	45.8	35.3	48.2	35.4	33.0	38.2
Dallas	22.8	58.6	50.7	46.5	50.6	20.7	36.4
District of Columbia	25.6	53.9	49.8	69.4	77.1	38.7	33.4
Los Angeles	23.9	42.4	21.5	71.7	36.8	22.6	25.8
Memphis	47.4	92.2	51.5	65.5	41.2	41.1	59.5
Miami	13.8	34.2	45.8	72.7	45.3	23.3	26.5
New Orleans	31.5	59.5	29.1	60.7	44.1	37.6	39.3
Orange County	15.9	44.5	26.0	87.2	61.2	10.2	48.1
Philadelphia	15.2	45.7	23.5	41.6	37.3	14.6	50.3
San Diego	26.8	31.9	57.3	85.6	90.4	29.1	32.3
<b>Local Median</b>	<b>24.8</b>	<b>45.8</b>	<b>40.6</b>	<b>67.5</b>	<b>44.7</b>	<b>26.2</b>	<b>37.3</b>
<b>Local Range</b>	<b>13.8 – 47.4</b>	<b>31.9 – 92.2</b>	<b>21.5 – 57.3</b>	<b>41.6 – 87.2</b>	<b>35.4 – 90.4</b>	<b>10.2 – 41.1</b>	<b>25.8 – 59.5</b>

\* HIV = human immunodeficiency virus.

TABLE 19c. Percentage of Schools in Which the Lead Health Education Teacher Had Received Staff Development During the Two Years Preceding the Survey on Specific Health Education Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Personal hygiene	Physical activity and fitness	Pregnancy prevention	STD* prevention	Suicide prevention	Sun safety or skin cancer prevention	Tobacco use prevention	Violence prevention
<b>STATE SURVEYS</b>								
Alaska	16.7	29.9	14.5	21.3	34.1	5.8	30.0	47.2
Arizona	20.7	36.9	21.0	27.4	27.2	24.2	38.6	53.9
Arkansas	12.8	46.9	16.9	22.1	16.3	13.3	45.5	52.4
Connecticut	10.1	45.9	22.8	33.1	18.9	3.6	23.1	55.7
Delaware	13.8	61.6	46.3	45.4	23.1	12.3	54.7	61.1
Idaho	9.6	43.3	25.5	51.0	32.5	11.1	35.2	49.2
Iowa	8.3	29.5	18.5	29.2	18.1	8.0	22.5	44.7
Maine	5.3	40.1	28.1	37.1	38.2	12.4	32.1	49.6
Massachusetts	9.7	44.8	26.7	35.5	23.0	32.3	30.1	62.9
Michigan	16.6	43.0	28.1	49.2	20.6	10.3	36.4	49.4
Missouri	18.3	49.5	22.4	31.6	21.0	10.1	31.4	47.1
Montana	15.7	55.3	27.7	40.1	27.9	12.6	40.8	52.3
Nebraska	9.6	32.6	15.6	20.4	14.8	6.4	25.1	42.7
New Hampshire	9.1	48.4	34.3	39.3	25.0	15.4	35.0	64.4
New York	9.4	41.9	32.1	38.7	21.9	10.2	32.7	57.3
North Carolina	8.2	51.2	28.2	36.6	9.4	8.2	29.1	36.1
North Dakota	16.1	50.0	15.8	33.3	31.1	12.1	54.1	59.7
Oklahoma	19.3	31.7	21.4	44.1	25.7	8.8	36.7	74.3
Oregon	7.2	32.5	23.4	34.2	24.9	7.4	30.5	52.2
Pennsylvania	10.1	53.0	21.6	31.3	21.8	10.8	36.6	59.7
South Carolina	19.3	51.2	33.7	37.6	11.4	14.4	28.5	38.2
Tennessee	22.9	52.2	24.8	34.5	26.6	15.7	37.6	61.7
Utah	6.5	34.9	20.3	29.2	21.8	10.7	34.6	38.5
Washington	10.2	35.3	17.3	37.3	27.4	8.4	32.1	52.1
Wisconsin	7.6	42.4	24.4	27.4	25.7	8.3	33.9	42.1
<b>State Median</b>	<b>10.1</b>	<b>43.3</b>	<b>23.4</b>	<b>34.5</b>	<b>23.1</b>	<b>10.7</b>	<b>33.9</b>	<b>52.2</b>
<b>State Range</b>	<b>5.3 – 22.9</b>	<b>29.5 – 61.6</b>	<b>14.5 – 46.3</b>	<b>20.4 – 51.0</b>	<b>9.4 – 38.2</b>	<b>3.6 – 32.3</b>	<b>22.5 – 54.7</b>	<b>36.1 – 74.3</b>
<b>LOCAL SURVEYS</b>								
Chicago	30.9	49.4	32.2	42.2	18.6	15.0	32.7	52.3
Dallas	16.7	39.4	41.3	56.6	31.0	18.7	42.3	56.2
District of Columbia	32.3	61.3	67.0	67.4	38.7	20.5	64.6	73.8
Los Angeles	6.3	23.9	31.9	55.9	22.6	8.4	68.4	49.5
Memphis	44.0	75.8	43.2	48.4	35.2	31.0	55.4	81.9
Miami	19.2	27.0	27.8	48.5	22.1	22.1	32.0	37.9
New Orleans	35.7	57.8	31.3	51.6	31.2	9.3	50.2	59.5
Orange County	15.5	20.2	38.8	64.3	15.1	10.5	23.6	47.9
Philadelphia	16.0	56.8	20.0	37.0	16.0	13.2	37.4	62.6
San Diego	22.1	16.9	82.8	92.6	80.4	4.9	90.3	73.7
<b>Local Median</b>	<b>20.7</b>	<b>44.4</b>	<b>35.5</b>	<b>53.8</b>	<b>26.8</b>	<b>14.1</b>	<b>46.3</b>	<b>57.9</b>
<b>Local Range</b>	<b>6.3 – 44.0</b>	<b>16.9 – 75.8</b>	<b>20.0 – 82.8</b>	<b>37.0 – 92.6</b>	<b>15.1 – 80.4</b>	<b>4.9 – 31.0</b>	<b>23.6 – 90.3</b>	<b>37.9 – 81.9</b>

\* STD = sexually transmitted disease.

TABLE 20a. Percentage of Schools in Which the Lead Health Education Teacher Wanted to Receive Staff Development on Specific Health Education Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Accident or injury prevention	Alcohol or other drug use prevention	Consumer health	CPR*	Death and dying	Dental and oral health	Emotional and mental health
<b>STATE SURVEYS</b>							
Alaska	39.8	61.9	41.5	55.6	46.8	32.0	58.6
Arizona	54.5	73.3	47.9	66.1	50.4	47.2	65.6
Arkansas	58.8	73.1	51.9	70.3	52.5	45.2	59.3
Connecticut	38.6	74.7	54.5	46.9	57.3	28.8	73.7
Delaware	49.0	69.0	58.2	54.9	61.8	36.7	71.0
Idaho	48.9	65.0	48.0	58.0	50.9	31.5	63.8
Iowa	33.8	60.4	44.0	50.3	52.0	26.8	57.3
Maine	32.7	55.3	42.7	49.2	43.7	19.5	57.9
Massachusetts	44.8	71.7	50.8	58.8	61.7	36.0	75.6
Michigan	41.2	62.6	46.8	60.3	47.3	33.7	61.0
Missouri	51.3	70.9	44.9	68.5	48.6	37.4	57.3
Montana	43.9	67.7	41.2	59.1	53.8	34.5	65.6
Nebraska	38.5	59.2	35.0	55.1	45.3	27.3	53.3
New Hampshire	40.2	66.3	53.6	46.7	46.4	30.2	71.7
New York	50.5	72.6	55.4	55.6	62.4	37.8	69.6
North Carolina	48.3	71.1	42.4	67.9	50.7	37.0	62.8
North Dakota	42.0	56.8	42.3	60.6	43.5	29.3	56.6
Oklahoma	57.7	74.7	44.4	74.5	47.2	42.9	65.2
Oregon	27.0	60.6	38.5	33.7	45.9	21.9	53.9
Pennsylvania	51.0	78.7	52.4	64.8	58.7	37.3	72.8
South Carolina	45.2	65.0	41.6	60.3	47.3	37.3	57.9
Tennessee	53.0	66.8	46.3	66.6	52.8	44.8	64.5
Utah	37.4	67.0	46.8	50.1	53.2	32.4	66.3
Washington	37.7	67.1	47.5	47.5	49.6	26.8	61.5
Wisconsin	40.8	65.2	49.3	49.2	53.6	28.4	66.3
<b>State Median</b>	<b>43.9</b>	<b>67.0</b>	<b>46.8</b>	<b>58.0</b>	<b>50.7</b>	<b>33.7</b>	<b>63.8</b>
<b>State Range</b>	<b>27.0 – 58.8</b>	<b>55.3 – 78.7</b>	<b>35.0 – 58.2</b>	<b>33.7 – 74.5</b>	<b>43.5 – 62.4</b>	<b>19.5 – 47.2</b>	<b>53.3 – 75.6</b>
<b>LOCAL SURVEYS</b>							
Chicago	66.1	75.6	60.4	85.0	64.2	56.6	73.3
Dallas	38.2	74.8	39.4	74.0	48.9	46.7	68.0
District of Columbia	70.7	73.8	66.2	81.1	61.3	61.0	77.4
Los Angeles	63.9	81.1	65.0	79.0	69.3	55.1	77.0
Memphis	75.7	85.7	72.9	89.8	82.0	69.5	84.0
Miami	64.1	78.2	57.1	79.0	63.4	50.4	68.3
New Orleans	72.7	81.0	71.7	78.0	74.9	65.5	72.6
Orange County	36.1	72.1	51.8	62.0	62.0	38.4	70.2
Philadelphia	73.1	80.4	66.5	86.7	79.3	67.0	81.6
San Diego	62.7	70.7	60.4	78.2	90.1	47.9	90.1
<b>Local Median</b>	<b>65.1</b>	<b>76.9</b>	<b>62.7</b>	<b>79.0</b>	<b>66.8</b>	<b>55.9</b>	<b>75.2</b>
<b>Local Range</b>	<b>36.1 – 75.7</b>	<b>70.7 – 85.7</b>	<b>39.4 – 72.9</b>	<b>62.0 – 89.8</b>	<b>48.9 – 90.1</b>	<b>38.4 – 69.5</b>	<b>68.0 – 90.1</b>

\* CPR = cardiopulmonary resuscitation.

TABLE 20b. Percentage of Schools in Which the Lead Health Education Teacher Wanted to Receive Staff Development on Specific Health Education Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Environmental health	First aid	Growth and development	HIV* prevention	Human sexuality	Immunization and vaccinations	Nutrition and dietary behavior
<b>STATE SURVEYS</b>							
Alaska	44.3	58.5	42.4	53.2	40.9	35.9	54.5
Arizona	52.1	64.9	56.3	59.2	50.4	47.8	62.8
Arkansas	50.7	69.5	44.4	69.8	52.4	44.8	70.6
Connecticut	53.5	49.5	59.3	67.6	70.5	38.7	69.4
Delaware	53.4	54.6	61.8	70.0	79.1	40.0	66.8
Idaho	49.2	61.9	47.5	62.3	51.0	40.6	64.6
Iowa	42.6	47.1	40.3	55.6	51.0	38.4	53.7
Maine	49.6	47.6	33.7	52.3	50.1	29.6	53.5
Massachusetts	56.7	57.2	57.5	67.5	69.3	41.4	69.4
Michigan	50.6	62.1	50.3	54.8	51.2	38.7	64.8
Missouri	50.5	71.1	47.1	64.2	50.5	45.8	63.6
Montana	46.7	62.3	44.5	59.7	53.5	40.4	63.6
Nebraska	37.6	49.4	33.9	51.9	42.0	31.8	53.2
New Hampshire	58.5	46.8	50.8	58.8	59.5	38.7	75.5
New York	57.5	54.9	60.9	73.0	70.3	48.4	69.7
North Carolina	46.3	71.1	41.1	62.4	54.4	37.3	69.9
North Dakota	40.1	59.2	41.1	54.0	45.0	36.1	54.0
Oklahoma	49.6	72.2	46.8	62.7	44.0	46.0	64.9
Oregon	40.1	37.0	38.2	49.8	46.2	28.7	54.2
Pennsylvania	54.1	63.5	60.6	75.5	72.2	45.3	73.9
South Carolina	45.9	58.4	49.1	60.2	54.3	37.8	66.0
Tennessee	53.3	68.6	49.3	65.4	55.1	41.4	69.9
Utah	47.2	57.7	46.4	68.3	62.4	45.4	66.5
Washington	48.0	47.3	42.8	55.8	49.6	32.0	62.7
Wisconsin	49.9	53.8	55.0	60.9	60.6	35.1	60.3
<b>State Median</b>	<b>49.6</b>	<b>58.4</b>	<b>47.1</b>	<b>60.9</b>	<b>52.4</b>	<b>38.7</b>	<b>64.8</b>
<b>State Range</b>	<b>37.6 – 58.5</b>	<b>37.0 – 72.2</b>	<b>33.7 – 61.8</b>	<b>49.8 – 75.5</b>	<b>40.9 – 79.1</b>	<b>28.7 – 48.4</b>	<b>53.2 – 75.5</b>
<b>LOCAL SURVEYS</b>							
Chicago	59.5	80.2	61.3	70.2	63.3	56.8	72.3
Dallas	57.5	68.7	70.9	79.1	72.2	42.5	60.1
District of Columbia	70.4	77.8	64.3	80.8	87.2	71.0	80.5
Los Angeles	74.5	72.6	69.1	79.7	78.7	61.5	76.7
Memphis	75.5	83.8	75.7	82.1	76.3	75.6	77.8
Miami	66.2	83.5	70.4	72.9	68.2	51.1	74.3
New Orleans	74.1	84.2	74.8	84.2	81.0	70.8	74.8
Orange County	40.8	64.9	62.4	73.7	70.9	31.9	71.9
Philadelphia	69.1	85.4	76.8	84.8	83.8	65.0	74.1
San Diego	57.6	73.4	62.3	73.1	67.5	45.6	80.7
<b>Local Median</b>	<b>67.7</b>	<b>79.0</b>	<b>69.8</b>	<b>79.4</b>	<b>74.3</b>	<b>59.2</b>	<b>74.6</b>
<b>Local Range</b>	<b>40.8 – 75.5</b>	<b>64.9 – 85.4</b>	<b>61.3 – 76.8</b>	<b>70.2 – 84.8</b>	<b>63.3 – 87.2</b>	<b>31.9 – 75.6</b>	<b>60.1 – 80.7</b>

\* HIV = human immunodeficiency virus.

TABLE 20c. Percentage of Schools in Which the Lead Health Education Teacher Wanted to Receive Staff Development on Specific Health Education Topics, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Personal hygiene	Physical activity and fitness	Pregnancy prevention	STD* prevention	Suicide prevention	Sun safety or skin cancer prevention	Tobacco use prevention	Violence prevention
<b>STATE SURVEYS</b>								
Alaska	33.8	51.8	48.2	50.6	65.0	31.9	58.6	70.0
Arizona	50.8	60.2	57.7	60.1	67.4	57.8	61.1	77.5
Arkansas	45.5	65.6	56.2	65.0	68.4	50.6	66.2	72.4
Connecticut	38.9	56.5	64.5	68.4	71.4	48.2	58.9	73.3
Delaware	41.6	67.3	63.8	71.3	74.6	58.4	61.3	85.7
Idaho	34.5	54.4	53.9	61.7	74.5	42.7	56.7	77.8
Iowa	29.8	47.6	54.2	55.9	62.0	40.2	50.2	71.5
Maine	24.3	49.0	46.0	47.4	63.1	33.3	43.3	71.1
Massachusetts	39.7	60.2	63.2	72.7	79.9	52.2	59.2	80.2
Michigan	37.4	57.6	50.1	57.3	67.4	41.2	55.1	68.8
Missouri	43.7	61.8	56.8	61.7	69.7	52.1	63.6	71.5
Montana	38.5	60.5	51.2	57.0	74.4	45.8	61.0	75.9
Nebraska	30.2	49.1	46.5	50.5	58.3	39.7	49.5	66.4
New Hampshire	34.3	56.4	52.9	63.3	69.7	46.8	59.3	73.8
New York	42.6	57.7	64.9	70.8	74.0	52.7	65.1	76.0
North Carolina	39.4	67.9	55.7	63.5	67.2	49.0	61.2	73.6
North Dakota	30.8	56.3	43.2	52.3	60.3	49.0	51.5	65.9
Oklahoma	47.9	61.2	53.7	59.1	68.0	54.4	65.9	75.8
Oregon	25.7	45.7	44.7	51.9	64.4	29.8	47.4	63.8
Pennsylvania	41.2	74.4	68.1	74.8	74.8	52.1	67.8	79.4
South Carolina	41.7	59.3	57.9	58.6	60.7	48.3	63.1	75.6
Tennessee	48.8	69.7	59.2	62.9	69.6	51.2	58.2	75.1
Utah	34.1	51.4	56.0	63.5	77.2	47.1	57.8	81.0
Washington	35.8	52.1	49.8	57.3	64.3	40.0	54.3	66.7
Wisconsin	32.9	56.5	57.0	60.4	70.9	44.5	54.8	73.2
<b>State Median</b>	<b>38.5</b>	<b>57.6</b>	<b>55.7</b>	<b>60.4</b>	<b>68.4</b>	<b>48.2</b>	<b>58.9</b>	<b>73.6</b>
<b>State Range</b>	<b>24.3 – 50.8</b>	<b>45.7 – 74.4</b>	<b>43.2 – 68.1</b>	<b>47.4 – 74.8</b>	<b>58.3 – 79.9</b>	<b>29.8 – 58.4</b>	<b>43.3 – 67.8</b>	<b>63.8 – 85.7</b>
<b>LOCAL SURVEYS</b>								
Chicago	63.5	66.6	62.9	68.2	71.5	56.6	66.6	84.2
Dallas	39.4	52.2	65.9	81.2	72.2	51.8	70.1	81.5
District of Columbia	64.6	69.8	77.4	77.8	71.7	61.3	71.0	77.4
Los Angeles	54.7	69.5	73.5	75.3	85.9	61.7	71.0	88.5
Memphis	72.3	74.2	82.0	82.1	85.8	69.4	77.7	88.1
Miami	58.9	65.0	64.1	71.9	74.6	58.7	71.3	81.7
New Orleans	68.7	74.0	77.9	87.4	78.0	64.4	84.2	87.4
Orange County	43.7	45.7	64.3	71.3	71.3	53.9	62.8	73.4
Philadelphia	64.7	78.4	78.5	83.0	81.5	65.5	72.7	84.5
San Diego	50.4	60.2	68.3	67.5	87.7	47.9	65.9	83.3
<b>Local Median</b>	<b>61.2</b>	<b>68.1</b>	<b>70.9</b>	<b>76.6</b>	<b>76.3</b>	<b>60.0</b>	<b>71.0</b>	<b>83.8</b>
<b>Local Range</b>	<b>39.4 – 72.3</b>	<b>45.7 – 78.4</b>	<b>62.9 – 82.0</b>	<b>67.5 – 87.4</b>	<b>71.3 – 87.7</b>	<b>47.9 – 69.4</b>	<b>62.8 – 84.2</b>	<b>73.4 – 88.5</b>

\* STD = sexually transmitted disease.

TABLE 21. Percentage of Schools in Which the Lead Health Education Teacher Received Staff Development During the Two Years Preceding the Survey on Specific Teaching Methods, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Teaching students with physical or cognitive disabilities	Teaching students of various cultural backgrounds	Teaching students with limited English proficiency	Using interactive teaching methods	Encouraging family or community involvement	Teaching skills for behavior change
<b>STATE SURVEYS</b>						
Alaska	48.0	63.7	40.2	46.2	51.9	49.6
Arizona	59.8	56.2	49.5	63.9	42.3	64.2
Arkansas	49.3	41.4	28.1	54.1	54.1	56.6
Connecticut	52.6	31.8	15.0	46.6	26.0	48.5
Delaware	40.1	40.6	22.1	60.0	30.0	61.1
Idaho	34.8	29.8	24.1	53.5	25.8	47.6
Iowa	44.8	34.9	18.3	52.3	35.1	57.6
Maine	33.4	14.1	8.7	45.4	22.2	39.5
Massachusetts	50.6	31.9	23.5	54.4	28.9	53.5
Michigan	38.8	26.5	10.1	51.0	33.8	55.9
Missouri	48.5	39.1	19.3	62.3	46.7	62.9
Montana	35.2	24.3	10.8	44.8	32.9	51.0
Nebraska	35.1	30.6	15.1	36.6	26.3	47.9
New Hampshire	54.4	17.0	8.5	55.2	36.5	60.4
New York	41.5	28.2	20.4	59.1	34.4	58.4
North Carolina	48.8	49.3	36.9	54.6	32.5	45.0
North Dakota	39.9	20.9	10.8	51.9	33.9	58.1
Oklahoma	47.8	60.6	32.8	52.8	53.5	55.8
Oregon	47.0	47.7	43.1	45.5	32.0	44.5
Pennsylvania	50.9	42.5	37.6	50.5	31.3	57.1
South Carolina	39.3	43.2	23.9	51.3	41.7	49.5
Tennessee	49.4	39.8	23.7	59.0	44.8	57.6
Utah	43.7	48.8	46.8	52.3	29.2	46.4
Washington	38.0	37.5	21.3	42.7	32.0	41.4
Wisconsin	32.8	28.3	18.6	40.7	30.8	45.4
<b>State Median</b>	<b>44.8</b>	<b>37.5</b>	<b>22.1</b>	<b>52.3</b>	<b>32.9</b>	<b>53.5</b>
<b>State Range</b>	<b>32.8 – 59.8</b>	<b>14.1 – 63.7</b>	<b>8.5 – 49.5</b>	<b>36.6 – 63.9</b>	<b>22.2 – 54.1</b>	<b>39.5 – 64.2</b>
<b>LOCAL SURVEYS</b>						
Chicago	60.3	45.9	32.6	65.0	49.2	58.9
Dallas	75.6	73.1	87.9	77.2	42.9	67.9
District of Columbia	54.9	48.5	45.4	57.9	51.2	57.9
Los Angeles	67.5	73.6	84.3	63.9	38.1	59.3
Memphis	62.6	50.3	27.4	71.0	60.8	62.7
Miami	31.2	46.1	60.9	62.8	37.3	40.6
New Orleans	39.4	30.2	9.1	71.8	51.7	66.8
Orange County	43.8	70.9	73.7	60.9	35.7	56.2
Philadelphia	16.7	27.5	16.6	42.8	29.4	46.8
San Diego	47.7	59.9	34.2	75.5	50.2	72.9
<b>Local Median</b>	<b>51.3</b>	<b>49.4</b>	<b>39.8</b>	<b>64.5</b>	<b>46.1</b>	<b>59.1</b>
<b>Local Range</b>	<b>16.7 – 75.6</b>	<b>27.5 – 73.6</b>	<b>9.1 – 87.9</b>	<b>42.8 – 77.2</b>	<b>29.4 – 60.8</b>	<b>40.6 – 72.9</b>

TABLE 22. Percentage of Schools in Which the Lead Health Education Teacher Wanted to Receive Staff Development on Specific Teaching Methods, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Teaching students with physical or cognitive disabilities	Teaching students of various cultural backgrounds	Teaching students with limited English proficiency	Using interactive teaching methods	Encouraging family or community involvement	Teaching skills for behavior change
<b>STATE SURVEYS</b>						
Alaska	57.9	53.4	48.2	58.7	66.5	78.1
Arizona	61.0	62.6	58.7	62.7	67.1	76.4
Arkansas	57.9	55.6	54.1	55.3	65.9	69.3
Connecticut	66.9	61.7	55.4	68.8	66.9	76.9
Delaware	61.6	61.6	56.8	66.2	69.1	75.2
Idaho	51.4	52.9	47.6	58.6	58.6	66.4
Iowa	52.3	43.8	35.0	48.5	54.1	65.0
Maine	50.0	40.1	35.8	56.9	61.0	68.6
Massachusetts	64.0	59.4	53.4	66.8	70.4	76.5
Michigan	57.3	48.4	41.0	51.7	58.8	66.6
Missouri	56.8	49.6	42.5	56.9	64.2	72.5
Montana	53.7	39.6	29.8	55.1	61.2	70.6
Nebraska	50.0	40.9	39.5	45.6	51.4	59.2
New Hampshire	58.8	45.9	29.8	62.2	67.6	79.5
New York	69.7	62.2	53.9	70.9	74.8	76.8
North Carolina	68.1	63.9	69.4	59.3	61.5	72.8
North Dakota	49.7	37.4	23.4	53.4	60.5	69.5
Oklahoma	64.6	57.5	51.2	59.8	72.3	74.7
Oregon	51.9	49.4	54.7	52.5	58.4	66.0
Pennsylvania	69.2	55.5	50.7	64.8	68.5	75.2
South Carolina	64.6	61.8	58.8	55.9	65.3	70.1
Tennessee	66.1	56.6	55.4	64.2	69.0	74.4
Utah	56.1	57.3	55.4	57.6	60.3	67.4
Washington	51.0	54.4	47.9	55.3	58.4	71.1
Wisconsin	51.6	52.3	39.9	59.2	63.1	71.2
<b>State Median</b>	<b>57.9</b>	<b>54.4</b>	<b>50.7</b>	<b>58.6</b>	<b>64.2</b>	<b>71.2</b>
<b>State Range</b>	<b>49.7 – 69.7</b>	<b>37.4 – 63.9</b>	<b>23.4 – 69.4</b>	<b>45.6 – 70.9</b>	<b>51.4 – 74.8</b>	<b>59.2 – 79.5</b>
<b>LOCAL SURVEYS</b>						
Chicago	76.8	66.7	63.9	69.8	72.9	78.6
Dallas	62.4	74.9	63.2	67.4	79.7	80.2
District of Columbia	79.8	82.9	77.4	93.0	82.9	89.9
Los Angeles	71.2	65.4	68.4	74.3	79.9	79.6
Memphis	87.3	73.8	82.5	85.0	83.2	81.3
Miami	68.5	68.9	66.6	67.1	73.0	84.6
New Orleans	90.9	87.4	81.0	84.8	90.7	90.7
Orange County	61.6	51.2	48.1	61.2	69.0	61.1
Philadelphia	78.7	74.0	70.5	78.9	82.7	87.2
San Diego	70.7	75.8	70.7	71.1	68.4	73.4
<b>Local Median</b>	<b>74.0</b>	<b>73.9</b>	<b>69.5</b>	<b>72.7</b>	<b>79.8</b>	<b>80.8</b>
<b>Local Range</b>	<b>61.6 – 90.9</b>	<b>51.2 – 87.4</b>	<b>48.1 – 82.5</b>	<b>61.2 – 93.0</b>	<b>68.4 – 90.7</b>	<b>61.1 – 90.7</b>

TABLE 23. Percentage of Schools That Required Physical Education in Any of Grades 6–12, and Among Those Schools, the Percentage That Required Students to Take Only One Course or Two or More Courses and the Percentage That Required Students Who Fail a Required Physical Education Course to Repeat It; and the Percentage of Schools in Which a Newly Hired Physical Education Teacher Is Required to Be Certified in Physical Education,\* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2004

Site	Required physical education				Newly hired physical education teacher required to be certified in physical education
	Required physical education	Required only one physical education course	Required two or more physical education courses	Required students who failed a required physical education course to repeat it	
<b>STATE SURVEYS</b>					
Alaska	88.8	21.2	76.2	78.2	45.8
Arizona	71.2	41.5	57.9	45.9	70.5
Arkansas	97.6	39.1	60.4	79.1	99.5
Connecticut	98.7	16.8	83.2	47.2	99.2
Delaware	100.0	16.6	81.4	48.8	98.2
Idaho	94.5	20.6	78.4	55.3	91.3
Iowa	98.5	8.8	90.8	65.4	98.1
Maine	99.5	20.6	79.4	63.5	96.4
Massachusetts	96.2	16.0	83.6	40.3	97.1
Michigan	90.4	38.4	61.6	56.2	95.1
Minnesota	98.8	9.5	90.0	67.1	98.7
Missouri	99.7	25.2	74.0	59.9	97.3
Montana	99.3	6.8	92.2	66.4	92.7
Nebraska	99.7	14.4	85.6	68.8	95.8
New Hampshire	99.5	14.9	83.7	42.3	96.7
New York	99.7	7.0	93.0	61.6	98.7
North Carolina	97.9	43.0	55.5	47.2	97.8
North Dakota	99.5	9.3	90.1	76.9	95.5
Oklahoma	41.1	53.5	44.2	9.7	94.0
Oregon	98.6	17.4	80.9	52.1	86.8
Pennsylvania	98.8	8.5	91.2	57.4	98.4
South Carolina	95.9	46.3	51.3	54.8	99.6
Tennessee	87.3	45.1	49.4	47.3	97.8
Texas**	98.1	18.1	81.7	70.7	97.7
Utah	99.5	9.6	90.4	60.4	100.0
Washington	95.9	5.9	94.1	58.1	88.6
Wisconsin	97.3	5.3	93.9	64.9	94.5
<b>State Median</b>	<b>98.5</b>	<b>16.8</b>	<b>81.7</b>	<b>58.1</b>	<b>97.1</b>
<b>State Range</b>	<b>41.1 – 100.0</b>	<b>5.3 – 53.5</b>	<b>44.2 – 94.1</b>	<b>9.7 – 79.1</b>	<b>45.8 – 100.0</b>
<b>LOCAL SURVEYS</b>					
Chicago	98.5	45.0	47.3	20.5	98.4
Dallas	96.0	15.3	84.7	55.7	98.0
District of Columbia	93.0	51.9	44.4	92.0	100.0
Los Angeles	100.0	3.2	96.8	43.4	98.9
Memphis	83.2	52.3	47.7	60.0	91.0
Miami	58.9	60.1	36.4	83.0	99.0
New Orleans	93.8	9.9	86.7	69.9	93.8
Orange County	53.5	42.2	57.8	73.3	85.0
Philadelphia	91.9	40.6	55.7	33.3	96.0
San Bernardino	93.8	6.7	93.3	57.1	100.0
San Diego	97.6	5.0	95.0	45.1	95.2
<b>Local Median</b>	<b>93.8</b>	<b>40.6</b>	<b>57.8</b>	<b>57.1</b>	<b>98.0</b>
<b>Local Range</b>	<b>53.5 – 100.0</b>	<b>3.2 – 60.1</b>	<b>36.4 – 96.8</b>	<b>20.5 – 92.0</b>	<b>85.0 – 100.0</b>

\* Certification, licensure, or endorsement by the state.

\*\* Survey did not include schools from two of the state's largest school districts.

TABLE 24. Among Schools That Required a Physical Education Course in Any of Grades 6–12, the Percentage That Taught a Required Physical Education Course in Each Grade, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2004

Site	6th grade	7th grade	8th grade	9th grade	10th grade	11th grade	12th grade
<b>STATE SURVEYS</b>							
Alaska	75.6	80.7	80.3	86.4	75.0	64.9	63.3
Arizona	93.0	91.6	88.5	86.4	61.3	54.8	56.6
Arkansas	88.9	87.1	73.2	86.9	65.3	53.2	53.2
Connecticut	100.0	100.0	100.0	98.0	93.9	80.6	72.9
Delaware	100.0	100.0	96.8	91.3	92.0	50.3	44.8
Idaho	94.4	95.6	90.5	54.0	70.8	45.2	36.6
Iowa	98.3	98.8	98.8	97.8	97.2	96.6	96.6
Maine	98.7	98.0	98.1	94.7	75.9	30.2	26.6
Massachusetts	99.3	98.9	99.2	97.4	94.1	69.5	59.9
Michigan	91.9	90.3	81.9	93.3	35.5	27.5	26.1
Minnesota	96.7	97.1	96.4	92.5	89.0	14.2	12.7
Missouri	97.6	98.8	97.9	93.3	51.9	34.8	34.6
Montana	98.3	99.5	99.5	99.5	96.1	12.1	12.1
Nebraska	98.6	99.1	99.5	88.1	54.8	26.4	25.4
New Hampshire	98.2	99.2	98.5	89.2	57.5	37.4	27.0
New York	98.1	99.0	99.5	100.0	100.0	100.0	100.0
North Carolina	97.4	96.9	95.8	99.3	17.1	12.0	11.3
North Dakota	99.2	100.0	100.0	96.6	74.3	30.0	27.3
Oklahoma	99.1	68.1	62.6	21.1	25.9	24.0	20.1
Oregon	97.5	95.9	95.3	91.9	56.9	36.8	22.9
Pennsylvania	98.5	98.2	97.5	96.1	95.9	92.5	86.0
South Carolina	98.0	98.3	98.4	97.1	48.1	46.8	46.8
Tennessee	95.7	95.4	94.2	95.2	33.3	24.2	24.2
Texas*	96.8	93.9	77.0	97.7	96.8	67.5	66.9
Utah	100.0	98.2	95.5	93.0	94.7	70.1	50.1
Washington	96.1	97.9	95.6	92.3	83.4	52.3	51.7
Wisconsin	100.0	100.0	98.8	96.6	95.9	83.1	42.4
<b>State Median</b>	<b>98.1</b>	<b>98.2</b>	<b>96.8</b>	<b>93.3</b>	<b>75.0</b>	<b>46.8</b>	<b>42.4</b>
<b>State Range</b>	<b>75.6 – 100.0</b>	<b>68.1 – 100.0</b>	<b>62.6 – 100.0</b>	<b>21.1 – 100.0</b>	<b>17.1 – 100.0</b>	<b>12.0 – 100.0</b>	<b>11.3 – 100.0</b>
<b>LOCAL SURVEYS</b>							
Chicago	98.5	98.5	98.5	100.0	92.9	24.7	18.7
Dallas	100.0	95.7	100.0	100.0	100.0	90.5	90.0
District of Columbia	80.0	71.4	71.4	96.0	85.7	57.1	42.9
Los Angeles	98.1	98.1	98.1	97.6	97.6	28.6	25.4
Memphis	82.4	100.0	100.0	100.0	33.3	27.8	27.8
Miami	79.2	52.2	43.2	78.5	83.8	41.2	41.2
New Orleans	100.0	93.7	93.7	87.6	100.0	46.2	38.5
Orange County	100.0	100.0	100.0	87.4	72.7	70.0	77.8
Philadelphia	97.2	100.0	100.0	79.8	83.3	77.8	65.7
San Bernardino	100.0	100.0	100.0	100.0	85.7	42.9	33.3
San Diego	95.6	100.0	100.0	100.0	94.4	27.3	27.3
<b>Local Median</b>	<b>98.1</b>	<b>98.5</b>	<b>100.0</b>	<b>97.6</b>	<b>85.7</b>	<b>42.9</b>	<b>38.5</b>
<b>Local Range</b>	<b>79.2 – 100.0</b>	<b>52.2 – 100.0</b>	<b>43.2 – 100.0</b>	<b>78.5 – 100.0</b>	<b>33.3 – 100.0</b>	<b>24.7 – 90.5</b>	<b>18.7 – 90.0</b>

\* Survey did not include schools from two of the state's largest school districts.

TABLE 25. Among Schools That Required a Physical Education Course for Students in Any of Grades 6–12, the Percentage That Allowed Students to Be Exempted from Taking a Required Physical Education Course and the Percentage That Did Not Allow Students to Be Exempted from Taking a Required Physical Education Course for Any Reason, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2004

Site	Reason for exemption from a required physical education course				Could not be exempted from a required physical education course for any reason
	Enrollment in other courses	Participation in school sports	Participation in other school activities*	Participation in community sports	
<b>STATE SURVEYS</b>					
Alaska	7.8	31.7	10.7	11.8	61.1
Arizona	15.6	8.8	19.7	2.3	68.5
Arkansas	3.5	18.7	10.1	1.1	78.0
Connecticut	2.7	1.6	1.2	1.3	93.9
Delaware	12.3	0.0	4.1	2.0	87.7
Idaho	4.8	5.7	5.6	1.7	85.0
Iowa	44.0	23.2	6.9	0.4	47.1
Maine	2.7	4.2	1.6	2.1	92.2
Massachusetts	7.7	4.6	5.3	3.2	84.6
Michigan	12.3	15.1	20.8	2.5	64.6
Minnesota	4.0	4.5	1.6	3.1	92.0
Missouri	2.2	1.5	2.4	0.2	94.6
Montana	3.1	1.1	1.4	1.0	95.9
Nebraska	7.5	4.9	2.6	0.4	86.0
New Hampshire	4.7	7.2	3.6	1.8	87.0
New York	1.4	8.4	1.7	2.7	88.6
North Carolina	6.3	0.8	5.2	0.8	89.3
North Dakota	2.6	0.5	0.0	0.0	97.4
Oklahoma	21.0	43.9	25.4	6.9	49.6
Oregon	13.1	9.1	9.5	10.3	76.9
Pennsylvania	6.4	2.8	4.4	0.3	89.3
South Carolina	8.7	1.4	43.8	0.0	52.5
Tennessee	8.6	3.6	27.9	0.0	66.4
Texas**	6.9	83.2	62.5	20.1	9.5
Utah	7.8	20.2	14.4	3.1	71.1
Washington	24.4	40.4	23.1	23.7	38.6
Wisconsin	2.0	0.5	0.3	0.8	96.4
<b>State Median</b>	<b>6.9</b>	<b>4.9</b>	<b>5.3</b>	<b>1.8</b>	<b>85.0</b>
<b>State Range</b>	<b>1.4 – 44.0</b>	<b>0.0 – 83.2</b>	<b>0.0 – 62.5</b>	<b>0.0 – 23.7</b>	<b>9.5 – 97.4</b>
<b>LOCAL SURVEYS</b>					
Chicago	1.5	4.4	13.6	0.8	84.2
Dallas	7.0	52.7	80.1	0.0	19.9
District of Columbia	7.8	0.0	10.4	0.0	85.7
Los Angeles	3.2	37.3	48.5	5.8	45.8
Memphis	16.8	0.0	35.1	0.0	53.9
Miami	23.2	38.9	41.7	2.0	37.8
New Orleans	11.2	0.0	34.1	0.0	55.8
Orange County	24.7	62.0	73.3	0.0	10.7
Philadelphia	4.6	1.9	3.6	0.9	89.9
San Bernardino	0.0	20.0	66.7	0.0	33.3
San Diego	0.0	39.1	42.6	17.6	48.4
<b>Local Median</b>	<b>7.0</b>	<b>20.0</b>	<b>41.7</b>	<b>0.0</b>	<b>48.4</b>
<b>Local Range</b>	<b>0.0 – 24.7</b>	<b>0.0 – 62.0</b>	<b>3.6 – 80.1</b>	<b>0.0 – 17.6</b>	<b>10.7 – 89.9</b>

\* Such as ROTC, marching band, chorus, or cheerleading.

\*\* Survey did not include schools from two of the state's largest school districts.

TABLE 26. Percentage of Schools That Allowed Use of Activity or Athletic Facilities\* or Offered Opportunities for Students to Participate in Intramural Activities or Physical Activity Clubs and, Among Schools That Offered Opportunities for Students to Participate in Intramural Activities or Physical Activity Clubs, the Percentage That Provided Transportation Home,\*\* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2004

Site	Allowed use of activity or athletic facilities	Offered intramural activities or physical activity clubs	Provided transportation home
<b>STATE SURVEYS</b>			
Alaska	83.0	74.6	9.8
Arizona	64.2	69.3	51.2
Arkansas	81.2	29.7	16.4
Connecticut	88.6	77.6	54.3
Delaware	91.7	76.4	61.6
Idaho	95.8	53.2	14.3
Iowa	92.1	41.7	28.8
Maine	94.3	81.4	54.1
Massachusetts	89.1	82.7	43.5
Michigan	88.3	73.6	15.4
Minnesota	89.5	60.4	47.5
Missouri	88.5	61.6	30.1
Montana	92.9	58.9	13.3
Nebraska	94.0	45.8	17.1
New Hampshire	94.9	79.8	36.8
New York	92.2	88.8	68.3
North Carolina	90.4	53.9	20.0
North Dakota	88.7	47.1	20.4
Oklahoma	84.2	32.9	16.0
Oregon	91.0	60.1	24.6
Pennsylvania	87.6	79.4	48.4
South Carolina	80.0	57.6	14.8
Tennessee	83.2	52.9	12.4
Texas <sup>§</sup>	87.1	41.7	28.4
Utah	95.5	67.7	28.4
Washington	89.4	64.3	43.6
Wisconsin	91.3	67.0	27.2
<b>State Median</b>	<b>89.4</b>	<b>61.6</b>	<b>28.4</b>
<b>State Range</b>	<b>64.2 – 95.8</b>	<b>29.7 – 88.8</b>	<b>9.8 – 68.3</b>
<b>LOCAL SURVEYS</b>			
Chicago	49.2	91.0	12.2
Dallas	70.9	76.3	35.4
District of Columbia	66.2	74.5	8.5
Los Angeles	94.7	87.5	31.0
Memphis	61.5	66.9	30.4
Miami	61.9	87.3	50.3
New Orleans	52.7	75.9	36.3
Orange County	92.5	85.0	15.8
Philadelphia	66.2	86.7	17.9
San Bernardino	86.7	75.0	54.5
San Diego	76.3	78.4	57.3
<b>Local Median</b>	<b>66.2</b>	<b>78.4</b>	<b>31.0</b>
<b>Local Range</b>	<b>49.2 – 94.7</b>	<b>66.9 – 91.0</b>	<b>8.5 – 57.3</b>

\* For community-sponsored sports teams or physical activity programs outside of school hours or when school is not in session.

\*\* For students who participated in after-school intramural activities or physical activity clubs.

§ Survey did not include schools from two of the state's largest school districts.

TABLE 27a. Percentage of Schools That Implemented School-Based Asthma Management Activities, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2004

Site	Assured immediate access to medications*	Provided a full-time registered nurse**	Identified and tracked all students with asthma	Obtained and used an Asthma Action Plan <sup>§</sup>	Provided intensive case management for students with asthma <sup>§§</sup>
<b>STATE SURVEYS</b>					
Alaska	71.3	18.7	47.1	24.8	14.3
Arizona	80.4	46.0	78.7	51.5	35.2
Arkansas	90.7	53.5	82.2	62.3	32.6
Connecticut	89.8	89.2	96.4	74.6	44.2
Delaware	91.1	98.4	94.8	79.0	47.1
Idaho	89.9	14.3	58.6	29.3	27.4
Iowa	94.3	39.3	86.8	54.5	37.0
Maine	94.3	51.6	92.2	57.0	35.4
Massachusetts	92.9	94.8	96.0	74.4	51.0
Michigan	94.2	12.0	72.3	43.2	21.9
Minnesota	93.7	36.1	87.2	62.9	39.6
Missouri	93.5	73.2	94.3	67.3	40.8
Montana	89.9	13.0	71.9	38.6	24.8
Nebraska	92.8	26.4	91.2	71.1	33.8
New Hampshire	94.5	87.5	97.3	67.2	46.7
New York	94.8	92.3	97.1	62.2	41.4
North Carolina	94.7	9.4	85.8	76.4	44.9
North Dakota	90.3	5.2	58.3	32.4	25.1
Oklahoma	95.2	25.9	66.8	42.6	24.2
Oregon	90.5	6.3	75.8	49.3	31.6
Pennsylvania	89.1	81.0	94.5	63.7	33.9
South Carolina	91.4	60.0	83.7	68.1	47.7
Tennessee	94.8	24.7	78.1	60.0	25.0
Texas <sup>+</sup>	92.8	78.9	88.0	63.2	41.4
Utah	94.5	0.5	59.6	46.3	40.9
Washington	93.8	16.2	92.5	76.3	36.8
Wisconsin	94.9	16.7	81.3	51.4	30.3
<b>State Median</b>	<b>92.9</b>	<b>36.1</b>	<b>85.8</b>	<b>62.2</b>	<b>35.4</b>
<b>State Range</b>	<b>71.3 – 95.2</b>	<b>0.5 – 98.4</b>	<b>47.1 – 97.3</b>	<b>24.8 – 79.0</b>	<b>14.3 – 51.0</b>
<b>LOCAL SURVEYS</b>					
Chicago	91.6	10.7	88.3	67.1	58.6
Dallas	85.8	79.9	91.9	75.9	65.2
District of Columbia	81.5	58.8	87.6	46.6	41.0
Los Angeles	87.0	93.2	91.5	66.3	48.5
Memphis	90.8	9.2	65.9	31.0	11.0
Miami	83.0	16.3	26.8	20.8	35.0
New Orleans	87.8	45.2	94.1	84.7	63.5
Orange County	91.8	55.8	67.5	57.3	54.0
Philadelphia	89.4	83.0	99.2	83.9	56.2
San Bernardino	92.9	12.5	68.8	38.5	21.4
San Diego	95.1	85.7	100.0	67.6	68.6
<b>Local Median</b>	<b>89.4</b>	<b>55.8</b>	<b>88.3</b>	<b>66.3</b>	<b>54.0</b>
<b>Local Range</b>	<b>81.5 – 95.1</b>	<b>9.2 – 93.2</b>	<b>26.8 – 100.0</b>	<b>20.8 – 84.7</b>	<b>11.0 – 68.6</b>

\* As prescribed by a physician and approved by parents.

\*\* All day every day.

§ For all students with asthma.

§§ Who are absent 10 days or more per year.

+ Survey did not include schools from two of the state's largest school districts.

TABLE 27b. Percentage of Schools That Implemented School-Based Asthma Management Activities, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2004

Site	Educated school staff about asthma	Educated students with asthma about asthma management	Taught asthma awareness to all students*	Encouraged full participation in physical education and physical activity**	Provided modified physical education and physical activities <sup>§</sup>
<b>STATE SURVEYS</b>					
Alaska	34.3	26.4	18.6	76.2	54.8
Arizona	46.0	47.0	12.2	85.5	70.7
Arkansas	52.5	45.4	30.4	95.6	85.8
Connecticut	57.0	75.9	24.3	98.1	86.3
Delaware	64.1	72.0	35.3	98.4	89.8
Idaho	33.2	28.9	25.7	96.1	78.3
Iowa	58.4	46.3	28.6	97.0	82.8
Maine	53.7	63.7	23.8	98.4	83.9
Massachusetts	55.4	70.9	24.3	97.5	84.7
Michigan	50.1	24.4	21.4	95.2	79.4
Minnesota	59.5	56.5	26.9	94.8	88.0
Missouri	63.1	63.6	31.7	98.9	86.2
Montana	43.1	36.9	26.0	95.7	79.0
Nebraska	82.0	56.2	33.1	96.8	87.4
New Hampshire	66.2	75.1	23.4	99.4	87.5
New York	51.9	69.1	39.4	98.4	83.6
North Carolina	70.5	63.2	36.1	96.4	89.4
North Dakota	47.9	25.0	30.5	91.3	77.1
Oklahoma	54.7	33.6	13.9	88.5	67.6
Oregon	47.6	38.8	26.0	97.9	86.1
Pennsylvania	63.3	68.6	39.5	98.4	85.5
South Carolina	54.6	54.9	19.7	96.4	86.6
Tennessee	54.9	43.4	20.9	96.1	84.9
Texas <sup>§§</sup>	50.1	55.8	25.8	97.1	84.8
Utah	53.3	32.8	36.2	97.0	91.3
Washington	53.7	52.4	19.0	94.3	87.6
Wisconsin	49.0	43.2	21.9	95.0	81.8
<b>State Median</b>	<b>53.7</b>	<b>52.4</b>	<b>25.8</b>	<b>96.4</b>	<b>84.9</b>
<b>State Range</b>	<b>33.2 – 82.0</b>	<b>24.4 – 75.9</b>	<b>12.2 – 39.5</b>	<b>76.2 – 99.4</b>	<b>54.8 – 91.3</b>
<b>LOCAL SURVEYS</b>					
Chicago	80.4	68.1	42.2	96.1	85.6
Dallas	54.1	67.9	24.5	90.0	78.1
District of Columbia	30.0	41.5	25.3	74.4	62.4
Los Angeles	48.7	69.9	38.9	95.6	92.0
Memphis	50.8	37.8	18.5	94.4	71.6
Miami	30.1	17.8	22.4	85.2	67.9
New Orleans	69.9	73.0	39.4	97.0	88.0
Orange County	43.8	56.0	25.3	92.5	83.0
Philadelphia	75.0	83.0	52.8	96.8	86.7
San Bernardino	40.0	57.1	7.1	93.3	93.3
San Diego	83.4	75.7	7.5	100.0	87.8
<b>Local Median</b>	<b>50.8</b>	<b>67.9</b>	<b>25.3</b>	<b>94.4</b>	<b>85.6</b>
<b>Local Range</b>	<b>30.0 – 83.4</b>	<b>17.8 – 83.0</b>	<b>7.1 – 52.8</b>	<b>74.4 – 100.0</b>	<b>62.4 – 93.3</b>

\* In at least one grade.

\*\* When students with asthma are doing well.

§ As indicated by the student's Asthma Action Plan.

§§ Survey did not include schools from two of the state's largest school districts.

TABLE 28. The Percentage of Schools That Served Lunch to Students and, Among Those Schools, the Percentage in Which Students Usually Had  $\geq 20$  Minutes to Eat Lunch Once They Were Seated, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2004

Site	Served lunch to students	$\geq 20$ minutes to eat lunch
<b>STATE SURVEYS</b>		
Alaska	85.9	89.6
Arizona	87.2	83.9
Arkansas	100.0	89.1
Connecticut	100.0	76.3
Delaware	98.4	79.3
Idaho	100.0	83.9
Iowa	98.5	63.8
Maine	100.0	66.8
Massachusetts	99.8	72.1
Michigan	99.7	79.5
Minnesota	98.0	78.9
Missouri	100.0	65.2
Montana	98.5	80.7
Nebraska	99.3	88.4
New Hampshire	100.0	67.3
New York	99.7	91.8
North Carolina	99.7	74.1
North Dakota	97.5	73.5
Oklahoma	99.4	79.4
Oregon	96.9	86.7
Pennsylvania	99.4	85.6
South Carolina	100.0	79.5
Tennessee	100.0	88.0
Texas*	99.7	86.6
Utah	98.6	77.3
Washington	95.5	83.8
Wisconsin	97.9	78.5
<b>State Median</b>	<b>99.4</b>	<b>79.5</b>
<b>State Range</b>	<b>85.9 – 100.0</b>	<b>63.8 – 91.8</b>
<b>LOCAL SURVEYS</b>		
Chicago	100.0	57.0
Dallas	98.0	67.4
District of Columbia	97.2	89.6
Los Angeles	98.9	89.7
Memphis	100.0	81.8
Miami	98.9	88.3
New Orleans	100.0	76.0
Orange County	100.0	68.0
Philadelphia	100.0	91.9
San Bernardino	93.8	73.3
San Diego	100.0	83.3
<b>Local Median</b>	<b>100.0</b>	<b>81.8</b>
<b>Local Range</b>	<b>93.8 – 100.0</b>	<b>57.0 – 91.9</b>

\* Survey did not include schools from two of the state's largest school districts.

TABLE 29. Percentage of Schools That Allowed Students to Purchase Snack Foods or Beverages From Vending Machines or at the School Store, Canteen, or Snack Bar and, Among Those Schools, the Percentage Offering Less Nutritious and More Nutritious Foods or Beverages, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2004

Site	Allowed students to purchase snack foods or beverages	Less nutritious foods and beverages				More nutritious foods and beverages				
		Chocolate candy	Other kinds of candy	Salty snacks that are not low in fat*	Soft drinks, sports drinks, or fruit drinks**	Salty snacks that are low in fat <sup>§</sup>	Fruits or vegetables	Low-fat baked goods	100% fruit juice	Bottled water
<b>STATE SURVEYS</b>										
Alaska	59.8	76.5	77.0	81.6	94.3	79.6	21.5	55.6	81.6	83.9
Arizona	73.8	63.0	64.7	74.8	88.0	77.6	43.4	60.5	71.1	90.8
Arkansas	93.8	69.7	73.7	74.3	96.6	72.5	28.4	49.7	71.9	90.3
Connecticut	81.1	38.3	41.2	72.7	88.9	82.8	57.8	63.1	85.6	93.2
Delaware	80.5	35.6	39.9	74.5	89.2	76.8	45.0	68.4	95.5	100.0
Idaho	92.7	72.4	73.4	73.9	97.3	73.2	32.9	52.9	77.0	96.2
Iowa	89.5	66.8	67.7	71.3	97.0	71.8	43.0	60.1	87.4	94.8
Maine	91.7	40.0	45.1	61.4	78.9	79.6	50.0	68.9	94.9	100.0
Massachusetts	85.4	35.3	40.5	73.9	86.4	79.8	51.0	58.0	86.1	94.0
Michigan	92.1	72.3	76.0	86.2	96.8	85.8	55.0	66.2	85.9	98.9
Minnesota	91.1	78.7	80.7	83.9	95.9	88.5	46.7	69.0	87.0	96.4
Missouri	90.2	68.6	71.1	75.9	97.2	76.1	32.6	56.8	76.3	90.5
Montana	88.8	60.1	61.9	53.5	94.5	57.5	27.4	46.0	85.9	90.4
Nebraska	84.1	61.9	63.3	61.2	99.5	61.7	24.8	53.3	82.3	93.5
New Hampshire	91.4	34.5	39.7	68.7	85.0	80.4	54.4	61.6	83.4	98.6
New York	90.1	41.5	43.4	70.3	84.9	75.1	44.5	58.0	83.6	93.3
North Carolina	88.4	54.9	60.0	81.8	92.4	81.1	40.4	68.2	84.2	94.8
North Dakota	77.0	61.7	62.9	56.5	96.7	55.6	21.9	41.0	83.5	98.7
Oklahoma	92.6	88.9	90.4	88.3	98.8	85.9	18.5	60.4	56.6	90.9
Oregon	87.6	70.9	77.6	85.2	95.4	82.8	48.0	62.9	78.3	94.9
Pennsylvania	84.0	60.0	68.0	83.3	90.5	86.3	45.7	64.7	87.4	94.3
South Carolina	92.1	74.1	76.5	87.0	95.8	83.8	26.9	67.1	76.5	92.1
Tennessee	90.1	73.8	78.1	80.1	96.8	78.7	27.7	59.9	75.7	93.3
Texas <sup>§§</sup>	81.1	65.0	56.0	63.9	86.2	82.8	56.3	75.2	89.5	97.8
Utah	95.0	93.4	94.4	89.4	97.6	92.5	50.5	80.2	87.0	94.1
Washington	90.7	71.6	74.7	80.3	96.6	83.9	45.6	64.1	86.1	95.6
Wisconsin	88.4	57.9	62.6	69.0	94.0	75.2	48.6	60.6	85.7	94.8
<b>State Median</b>	<b>89.5</b>	<b>65.0</b>	<b>67.7</b>	<b>74.5</b>	<b>95.4</b>	<b>79.6</b>	<b>44.5</b>	<b>60.6</b>	<b>84.2</b>	<b>94.3</b>
<b>State Range</b>	<b>59.8 – 95.0</b>	<b>34.5 – 93.4</b>	<b>39.7 – 94.4</b>	<b>53.5 – 89.4</b>	<b>78.9 – 99.5</b>	<b>55.6 – 92.5</b>	<b>18.5 – 57.8</b>	<b>41.0 – 80.2</b>	<b>56.6 – 95.5</b>	<b>83.9 – 100.0</b>
<b>LOCAL SURVEYS</b>										
Chicago	33.2	31.7	33.7	53.0	79.5	50.8	34.1	48.0	81.3	83.4
Dallas	95.8	82.4	75.9	89.2	100.0	82.5	45.8	66.4	89.1	97.8
District of Columbia	62.4	33.4	47.9	52.4	84.7	52.4	16.0	28.5	78.3	89.2
Los Angeles	96.1	73.5	82.5	82.6	53.4	88.1	54.0	60.0	86.6	95.6
Memphis	72.5	61.6	65.2	62.8	97.3	60.0	20.3	37.6	90.1	79.6
Miami	83.4	70.2	70.2	78.9	95.0	77.2	47.0	66.9	82.5	84.9
New Orleans	81.5	70.0	77.4	81.1	92.6	66.5	14.7	65.2	61.7	100.0
Orange County	86.3	58.0	62.3	85.8	87.2	83.5	61.4	66.5	84.3	100.0
Philadelphia	71.3	35.2	47.3	81.8	83.1	76.2	45.9	55.7	90.1	88.6
San Bernardino	81.3	27.3	41.7	91.7	100.0	83.3	66.7	75.0	83.3	100.0
San Diego	92.8	47.9	48.1	89.6	92.1	84.6	77.1	65.5	71.9	97.4
<b>Local Median</b>	<b>81.5</b>	<b>58.0</b>	<b>62.3</b>	<b>81.8</b>	<b>92.1</b>	<b>77.2</b>	<b>45.9</b>	<b>65.2</b>	<b>83.3</b>	<b>95.6</b>
<b>Local Range</b>	<b>33.2 – 96.1</b>	<b>27.3 – 82.4</b>	<b>33.7 – 82.5</b>	<b>52.4 – 91.7</b>	<b>53.4 – 100.0</b>	<b>50.8 – 88.1</b>	<b>14.7 – 77.1</b>	<b>28.5 – 75.0</b>	<b>61.7 – 90.1</b>	<b>79.6 – 100.0</b>

\* Such as regular potato chips.  
 \*\* That are not 100% juice.  
 § Such as pretzels, baked chips, or other low-fat chips.  
 §§ Survey did not include schools from two of the state's largest school districts.

TABLE 30. Among Schools That Allowed Students to Purchase Snack Foods or Beverages From Vending Machines or at the School Store, Canteen, or Snack Bar, the Percentage That Sold Items at Specific Times and the Percentage of Schools That Had a Policy That Fruits or Vegetables Will Be Offered at School Settings,\* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2004

Site	Before classes begin in the morning	During any school hours when meals are not being served	During school lunch periods	Offered fruits or vegetables
<b>STATE SURVEYS</b>				
Alaska	53.1	44.7	55.5	10.9
Arizona	60.3	41.0	67.6	15.6
Arkansas	57.2	34.0	84.7	6.6
Connecticut	40.6	32.7	67.0	9.2
Delaware	35.0	20.8	53.6	16.3
Idaho	82.8	62.0	82.6	7.5
Iowa	74.7	59.0	40.1	6.4
Maine	54.2	26.1	36.7	9.4
Massachusetts	41.2	24.1	73.1	10.5
Michigan	65.2	33.7	81.3	6.9
Minnesota	78.5	45.9	53.6	7.5
Missouri	68.7	38.0	69.2	5.5
Montana	73.8	47.8	66.0	10.0
Nebraska	67.8	52.2	22.2	7.2
New Hampshire	48.5	41.8	55.0	5.6
New York	37.2	31.2	57.4	17.4
North Carolina	34.8	34.7	46.7	11.5
North Dakota	75.3	51.7	42.2	8.5
Oklahoma	72.0	45.2	85.0	6.9
Oregon	67.1	51.2	72.2	8.4
Pennsylvania	47.7	23.3	70.3	11.0
South Carolina	38.8	28.7	75.1	11.1
Tennessee	40.9	52.9	35.3	10.4
Texas**	46.5	47.9	44.1	23.5
Utah	80.3	61.2	90.1	13.7
Washington	70.4	52.7	82.2	9.7
Wisconsin	68.0	45.3	49.3	10.7
<b>State Median</b>	<b>60.3</b>	<b>44.7</b>	<b>66.0</b>	<b>9.7</b>
<b>State Range</b>	<b>34.8 – 82.8</b>	<b>20.8 – 62.0</b>	<b>22.2 – 90.1</b>	<b>5.5 – 23.5</b>
<b>LOCAL SURVEYS</b>				
Chicago	33.2	15.9	66.7	34.1
Dallas	60.3	49.8	80.0	27.4
District of Columbia	18.1	13.6	79.3	28.6
Los Angeles	56.8	17.6	72.6	20.8
Memphis	23.7	7.4	48.7	11.3
Miami	63.8	44.8	55.8	21.1
New Orleans	40.3	22.1	70.3	40.9
Orange County	61.2	52.5	69.3	15.0
Philadelphia	28.1	11.0	81.8	39.0
San Bernardino	83.3	58.3	66.7	28.6
San Diego	43.1	22.8	89.5	17.0
<b>Local Median</b>	<b>43.1</b>	<b>22.1</b>	<b>70.3</b>	<b>27.4</b>
<b>Local Range</b>	<b>18.1 – 83.3</b>	<b>7.4 – 58.3</b>	<b>48.7 – 89.5</b>	<b>11.3 – 40.9</b>

\* Such as student parties, after-school programs, staff meetings, parents' meetings, or concession stands.

\*\* Survey did not include schools from two of the state's largest school districts.

TABLE 31. Percentage of Schools That Had a Policy Prohibiting Tobacco Use and, Among Those Schools, the Percentage That Had an "Ideal" Tobacco Use Prevention Policy,\* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2004

Site	Had a policy prohibiting tobacco use	Had an "ideal" tobacco use prevention policy
<b>STATE SURVEYS</b>		
Alaska	97.0	32.4
Arizona	96.8	63.1
Arkansas	99.5	61.8
Connecticut	96.2	50.9
Delaware	98.2	55.7
Idaho	100.0	37.4
Iowa	99.2	19.6
Maine	100.0	53.6
Massachusetts	99.3	58.5
Michigan	97.8	40.0
Minnesota	98.6	48.8
Missouri	97.4	27.6
Montana	99.4	30.6
Nebraska	100.0	21.2
New Hampshire	98.6	44.3
New York	98.0	61.8
North Carolina	97.5	32.3
North Dakota	98.6	25.3
Oklahoma	99.7	25.6
Oregon	98.4	50.0
Pennsylvania	99.4	46.1
South Carolina	99.6	46.2
Tennessee	98.3	30.1
Texas**	99.7	66.5
Utah	100.0	50.4
Washington	99.6	57.4
Wisconsin	97.0	54.6
<b>State Median</b>	<b>98.6</b>	<b>46.2</b>
<b>State Range</b>	<b>96.2 – 100.0</b>	<b>19.6 – 66.5</b>
<b>LOCAL SURVEYS</b>		
Chicago	85.4	36.0
Dallas	98.0	70.9
District of Columbia	93.3	39.7
Los Angeles	100.0	73.9
Memphis	89.2	56.4
Miami	95.8	50.5
New Orleans	100.0	28.0
Orange County	100.0	62.4
Philadelphia	90.3	37.0
San Bernardino	100.0	71.4
San Diego	100.0	66.5
<b>Local Median</b>	<b>98.0</b>	<b>56.4</b>
<b>Local Range</b>	<b>85.4 – 100.0</b>	<b>28.0 – 73.9</b>

\* An "ideal" tobacco use prevention policy, as described in CDC's *Guidelines for School Health Programs to Prevent Tobacco Use and Addiction*, prohibits all tobacco use by students, faculty, staff, and visitors during school and nonschool hours in school buildings; on school grounds; in school buses or other vehicles used to transport students; and at off-campus, school-sponsored events.

\*\* Survey did not include schools from two of the state's largest school districts.

TABLE 32a. Among Schools with a Policy Prohibiting Tobacco Use, the Percentage of Schools That Sometimes, Almost Always, or Always Took Specific Actions When Students Were Caught Smoking Cigarettes, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2004

Site	Informed parents or guardians	Referred to a school counselor	Referred to a school administrator	Encouraged to participate in an assistance, education, or cessation program	Required to participate in an assistance, education, or cessation program
<b>STATE SURVEYS</b>					
Alaska	98.0	55.2	94.9	56.2	33.8
Arizona	99.0	66.8	98.6	61.2	32.4
Arkansas	99.5	62.6	100.0	38.1	17.3
Connecticut	100.0	82.7	98.7	69.3	29.5
Delaware	100.0	73.3	100.0	59.8	24.3
Idaho	100.0	85.9	100.0	68.0	59.3
Iowa	99.6	78.0	99.6	62.6	32.4
Maine	100.0	88.6	99.5	82.2	51.9
Massachusetts	97.8	82.9	98.4	66.1	32.7
Michigan	100.0	71.9	100.0	66.2	37.5
Minnesota	98.5	75.5	97.2	65.0	46.1
Missouri	100.0	61.1	100.0	37.7	16.8
Montana	99.7	79.6	99.3	58.5	48.0
Nebraska	99.6	75.4	99.6	57.2	29.3
New Hampshire	100.0	84.5	100.0	73.8	37.1
New York	99.4	87.6	97.9	70.1	32.7
North Carolina	100.0	73.8	100.0	52.3	46.1
North Dakota	97.9	78.7	97.5	60.3	27.4
Oklahoma	98.6	66.4	98.8	39.5	15.2
Oregon	98.3	76.9	97.3	73.4	52.9
Pennsylvania	98.7	73.5	97.6	64.3	32.8
South Carolina	99.4	69.8	99.0	48.3	25.8
Tennessee	100.0	65.0	99.7	45.9	33.9
Texas*	100.0	76.4	99.7	47.0	25.3
Utah	100.0	74.6	100.0	80.0	68.4
Washington	98.8	72.7	98.2	77.9	59.6
Wisconsin	98.8	80.8	98.9	65.6	30.7
<b>State Median</b>	<b>99.6</b>	<b>75.4</b>	<b>99.3</b>	<b>62.6</b>	<b>32.7</b>
<b>State Range</b>	<b>97.8 – 100.0</b>	<b>55.2 – 88.6</b>	<b>94.9 – 100.0</b>	<b>37.7 – 82.2</b>	<b>15.2 – 68.4</b>
<b>LOCAL SURVEYS</b>					
Chicago	97.6	79.7	95.7	37.0	20.9
Dallas	100.0	73.7	100.0	57.2	42.9
District of Columbia	100.0	90.0	100.0	41.0	17.3
Los Angeles	100.0	93.5	86.5	73.8	77.3
Memphis	100.0	70.6	98.0	56.7	41.7
Miami	97.7	95.4	96.6	66.8	35.0
New Orleans	96.9	75.3	100.0	60.9	39.6
Orange County	100.0	95.5	100.0	85.3	54.6
Philadelphia	97.3	76.2	95.4	51.2	22.9
San Bernardino	100.0	100.0	100.0	60.0	92.9
San Diego	100.0	97.7	100.0	88.2	78.1
<b>Local Median</b>	<b>100.0</b>	<b>90.0</b>	<b>100.0</b>	<b>60.0</b>	<b>41.7</b>
<b>Local Range</b>	<b>96.9 – 100.0</b>	<b>70.6 – 100.0</b>	<b>86.5 – 100.0</b>	<b>37.0 – 88.2</b>	<b>17.3 – 92.9</b>

\* Survey did not include schools from two of the state's largest school districts.

TABLE 32b. Among Schools with a Policy Prohibiting Tobacco Use, the Percentage of Schools That Sometimes, Almost Always, or Always Took Specific Actions When Students Were Caught Smoking Cigarettes, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2004

Site	Referred to legal authorities	Placed in detention	Given in-school suspension	Suspended from school
<b>STATE SURVEYS</b>				
Alaska	50.9	54.1	59.0	74.3
Arizona	50.9	56.6	71.2	82.1
Arkansas	33.9	57.5	78.8	73.4
Connecticut	21.2	64.0	76.0	82.1
Delaware	12.6	60.5	77.0	78.4
Idaho	91.1	58.0	73.0	80.0
Iowa	73.8	55.5	73.9	68.1
Maine	81.3	36.4	51.3	91.3
Massachusetts	19.2	62.5	64.9	78.6
Michigan	69.0	33.7	49.4	92.3
Minnesota	76.8	45.8	64.1	80.0
Missouri	36.8	59.1	80.3	73.2
Montana	76.9	65.8	73.5	73.7
Nebraska	50.5	63.8	80.2	78.0
New Hampshire	83.7	39.5	60.4	82.2
New York	20.1	72.3	81.7	68.1
North Carolina	20.4	61.5	74.3	79.3
North Dakota	62.5	56.9	70.2	72.1
Oklahoma	39.1	70.9	77.1	73.6
Oregon	66.0	57.4	63.6	84.9
Pennsylvania	74.0	44.9	66.3	72.8
South Carolina	12.3	53.5	67.8	90.6
Tennessee	81.6	47.2	62.5	71.5
Texas*	75.5	62.1	91.2	53.1
Utah	98.1	46.8	56.1	85.5
Washington	47.2	53.1	60.5	82.9
Wisconsin	88.1	56.3	70.8	75.5
<b>State Median</b>	<b>62.5</b>	<b>56.9</b>	<b>70.8</b>	<b>78.4</b>
<b>State Range</b>	<b>12.3 – 98.1</b>	<b>33.7 – 72.3</b>	<b>49.4 – 91.2</b>	<b>53.1 – 92.3</b>
<b>LOCAL SURVEYS</b>				
Chicago	27.2	71.9	69.2	81.1
Dallas	54.9	79.5	87.8	85.8
District of Columbia	24.4	59.0	71.0	87.5
Los Angeles	45.0	67.5	57.8	84.6
Memphis	51.3	30.5	44.1	89.9
Miami	30.5	65.5	85.3	75.2
New Orleans	31.3	70.1	66.4	87.3
Orange County	61.8	57.7	79.5	76.3
Philadelphia	21.3	82.2	70.7	82.6
San Bernardino	66.7	46.7	73.3	93.3
San Diego	60.8	50.4	55.2	90.4
<b>Local Median</b>	<b>45.0</b>	<b>65.5</b>	<b>70.7</b>	<b>85.8</b>
<b>Local Range</b>	<b>21.3 – 66.7</b>	<b>30.5 – 82.2</b>	<b>44.1 – 87.8</b>	<b>75.2 – 93.3</b>

\* Survey did not include schools from two of the state's largest school districts.

TABLE 33. Among Schools with a Policy Prohibiting Tobacco Use, the Percentage of Schools That Had Procedures to Inform Specific Groups About the Tobacco Prevention Policy That Prohibits Their Use of Tobacco and to Inform Parents About the Policy That Prohibits Tobacco Use by Students, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2004

Site	Groups informed about policy prohibiting their use of tobacco			Informed parents of policy prohibiting tobacco use by students
	Students	Faculty and Staff	Visitors	
<b>STATE SURVEYS</b>				
Alaska	95.9	91.2	76.5	95.9
Arizona	99.4	98.3	89.8	97.3
Arkansas	99.5	98.1	91.6	98.6
Connecticut	98.8	95.5	78.5	97.9
Delaware	100.0	96.4	78.9	93.2
Idaho	99.5	93.9	72.9	97.6
Iowa	98.9	94.8	80.4	98.5
Maine	100.0	98.4	82.2	99.5
Massachusetts	99.8	96.7	84.0	99.3
Michigan	98.3	95.9	81.1	96.7
Minnesota	100.0	98.0	87.9	98.9
Missouri	98.8	95.4	81.4	98.5
Montana	99.0	95.3	79.0	98.0
Nebraska	100.0	94.3	72.1	98.2
New Hampshire	99.4	98.3	83.0	99.4
New York	98.4	96.7	91.6	97.3
North Carolina	100.0	99.6	83.2	99.1
North Dakota	99.5	97.5	80.2	97.3
Oklahoma	100.0	96.0	86.5	97.9
Oregon	98.4	96.4	83.8	95.8
Pennsylvania	99.4	98.1	89.9	98.7
South Carolina	99.2	97.1	82.8	98.8
Tennessee	100.0	96.5	87.9	98.9
Texas*	99.2	96.3	91.2	99.7
Utah	99.5	91.9	67.2	99.0
Washington	99.7	95.5	88.6	98.7
Wisconsin	99.3	95.5	82.2	96.3
<b>State Median</b>	<b>99.4</b>	<b>96.4</b>	<b>82.8</b>	<b>98.5</b>
<b>State Range</b>	<b>95.9 – 100.0</b>	<b>91.2 – 99.6</b>	<b>67.2 – 91.6</b>	<b>93.2 – 99.7</b>
<b>LOCAL SURVEYS</b>				
Chicago	93.4	92.0	81.5	92.5
Dallas	100.0	100.0	95.8	97.8
District of Columbia	97.0	96.9	93.6	97.0
Los Angeles	100.0	97.8	95.6	100.0
Memphis	100.0	97.8	85.2	100.0
Miami	100.0	98.9	91.8	96.7
New Orleans	100.0	96.9	90.3	90.7
Orange County	100.0	97.9	95.0	100.0
Philadelphia	98.2	94.5	84.1	97.2
San Bernardino	100.0	93.8	100.0	100.0
San Diego	100.0	95.2	90.5	100.0
<b>Local Median</b>	<b>100.0</b>	<b>96.9</b>	<b>91.8</b>	<b>97.8</b>
<b>Local Range</b>	<b>93.4 – 100.0</b>	<b>92.0 – 100.0</b>	<b>81.5 – 100.0</b>	<b>90.7 – 100.0</b>

\* Survey did not include schools from two of the state's largest school districts.

TABLE 34. Percentage of Schools That Prohibited Tobacco Advertising in Specific Locations, Through Sponsorship of School Events, on Student Apparel or Merchandise, and All Tobacco Advertising, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2004

Site	In the school building	On school grounds*	On school buses or other vehicles**	In school publications	Through sponsorship of school events	On tobacco brand-name apparel or merchandise <sup>§</sup>	All tobacco advertising <sup>§§</sup>
<b>STATE SURVEYS</b>							
Alaska	96.1	92.8	92.4	92.8	92.3	83.9	74.1
Arizona	96.4	97.0	95.0	94.8	94.8	93.5	83.4
Arkansas	94.9	95.4	93.0	92.5	86.4	95.7	79.6
Connecticut	93.7	92.4	93.2	93.7	93.9	74.9	68.0
Delaware	92.9	89.3	87.7	89.3	89.3	82.8	68.5
Idaho	96.0	95.9	94.5	95.4	91.4	97.0	85.3
Iowa	93.1	91.5	92.4	91.7	87.6	95.2	79.3
Maine	96.4	95.3	94.8	93.8	93.3	93.8	84.0
Massachusetts	94.9	94.1	93.6	93.1	92.4	71.8	62.8
Michigan	94.6	94.9	93.9	94.1	92.6	92.3	83.1
Minnesota	92.6	92.3	91.4	91.8	91.0	94.4	80.3
Missouri	94.8	93.6	93.4	94.1	90.4	93.6	80.1
Montana	94.8	94.1	93.1	91.8	91.2	96.8	83.2
Nebraska	93.2	92.5	92.8	92.8	92.2	98.9	86.4
New Hampshire	95.3	94.7	95.3	95.3	94.8	94.8	88.3
New York	95.8	95.1	95.2	93.2	92.0	72.9	65.2
North Carolina	90.9	90.5	90.5	88.7	93.3	79.5	67.9
North Dakota	94.0	92.6	94.0	93.6	88.3	98.5	82.3
Oklahoma	96.9	96.1	96.3	94.9	94.9	95.4	85.4
Oregon	96.2	96.2	95.9	95.9	96.3	94.3	88.0
Pennsylvania	94.9	94.4	92.0	92.0	93.0	86.6	74.4
South Carolina	95.1	95.1	94.7	93.9	93.2	87.1	75.8
Tennessee	94.5	94.0	94.4	93.2	93.7	93.2	82.8
Texas <sup>+</sup>	94.9	95.4	94.9	94.0	92.2	98.4	85.1
Utah	94.2	94.2	95.2	93.8	95.1	99.5	89.4
Washington	96.9	96.9	95.7	95.7	93.3	93.9	83.8
Wisconsin	94.2	93.5	93.6	92.2	94.2	93.4	81.9
<b>State Median</b>	<b>94.9</b>	<b>94.2</b>	<b>93.9</b>	<b>93.6</b>	<b>92.6</b>	<b>93.8</b>	<b>82.3</b>
<b>State Range</b>	<b>90.9 – 96.9</b>	<b>89.3 – 97.0</b>	<b>87.7 – 96.3</b>	<b>88.7 – 95.9</b>	<b>86.4 – 96.3</b>	<b>71.8 – 99.5</b>	<b>62.8 – 89.4</b>
<b>LOCAL SURVEYS</b>							
Chicago	93.4	91.0	90.5	90.2	88.0	89.3	72.4
Dallas	87.9	87.9	87.9	87.9	96.0	94.1	79.6
District of Columbia	91.0	87.6	91.0	88.2	82.5	82.0	60.6
Los Angeles	100.0	98.9	99.0	99.0	98.0	90.5	86.2
Memphis	94.6	92.7	92.6	92.7	94.4	92.7	82.9
Miami	94.7	92.4	93.6	89.2	92.6	81.9	68.9
New Orleans	91.1	91.1	88.0	85.1	84.5	100.0	72.0
Orange County	95.3	95.3	91.3	91.3	89.5	90.5	78.6
Philadelphia	97.5	94.4	93.5	92.7	91.9	83.5	69.0
San Bernardino	93.8	100.0	93.8	100.0	93.8	100.0	87.5
San Diego	97.6	97.6	97.6	95.1	90.6	100.0	85.7
<b>Local Median</b>	<b>94.6</b>	<b>92.7</b>	<b>92.6</b>	<b>91.3</b>	<b>91.9</b>	<b>90.5</b>	<b>78.6</b>
<b>Local Range</b>	<b>87.9 – 100.0</b>	<b>87.6 – 100.0</b>	<b>87.9 – 99.0</b>	<b>85.1 – 100.0</b>	<b>82.5 – 98.0</b>	<b>81.9 – 100.0</b>	<b>60.6 – 87.5</b>

\* Including on the outside of the building, on playing fields, or other areas of the campus.  
 \*\* Used to transport students.  
 § With tobacco company names, logos, or cartoon characters.  
 §§ Prohibiting all tobacco advertising in school buildings, on school grounds, on school buses or other vehicles, in school publications, and through sponsorship of school events, and prohibiting students from wearing tobacco brand-name apparel or carrying merchandise with tobacco company names, logos, or cartoon characters.  
 + Survey did not include schools from two of the state's largest school districts.

TABLE 35. Percentage of Schools That Provide Referrals to Tobacco Cessation Programs for Specific Groups and Post Signs Marking a Tobacco-Free School Zone,\* Selected U.S. Sites: School Health Profiles, Principal Surveys, 2004

Site	Provide referrals		Post signs marking a tobacco-free school zone
	Faculty and Staff	Students	
<b>STATE SURVEYS</b>			
Alaska	12.6	35.9	65.9
Arizona	18.7	39.8	81.7
Arkansas	8.3	21.1	85.3
Connecticut	19.6	42.7	41.6
Delaware	23.9	45.2	53.6
Idaho	18.8	64.9	61.0
Iowa	21.0	52.4	51.6
Maine	46.1	78.3	66.7
Massachusetts	37.5	59.6	65.9
Michigan	19.0	49.7	59.2
Minnesota	18.2	60.7	74.9
Missouri	13.2	23.4	51.6
Montana	20.9	53.3	68.8
Nebraska	17.5	35.7	56.0
New Hampshire	28.5	51.8	83.9
New York	24.8	50.1	62.5
North Carolina	25.1	52.1	56.5
North Dakota	19.8	44.8	58.5
Oklahoma	8.9	21.0	64.8
Oregon	23.7	58.9	69.8
Pennsylvania	16.2	53.3	59.5
South Carolina	14.9	34.9	46.9
Tennessee	7.5	29.0	60.6
Texas**	10.4	27.4	82.2
Utah	15.7	78.6	44.4
Washington	27.4	78.7	83.9
Wisconsin	20.0	47.0	68.0
<b>State Median</b>	<b>19.0</b>	<b>49.7</b>	<b>62.5</b>
<b>State Range</b>	<b>7.5 – 46.1</b>	<b>21.0 – 78.7</b>	<b>41.6 – 85.3</b>
<b>LOCAL SURVEYS</b>			
Chicago	11.4	18.8	33.1
Dallas	24.6	30.6	83.8
District of Columbia	2.8	9.5	73.0
Los Angeles	53.1	84.6	69.6
Memphis	11.3	35.7	34.5
Miami	27.6	52.3	90.5
New Orleans	18.4	30.4	54.9
Orange County	23.3	75.5	88.5
Philadelphia	12.0	30.9	29.9
San Bernardino	50.0	100.0	71.4
San Diego	42.4	90.5	73.7
<b>Local Median</b>	<b>23.3</b>	<b>35.7</b>	<b>71.4</b>
<b>Local Range</b>	<b>2.8 – 53.1</b>	<b>9.5 – 100.0</b>	<b>29.9 – 90.5</b>

\* A specified distance from school grounds where tobacco use by students, faculty and staff, and visitors is not allowed.

\*\* Survey did not include schools from two of the state's largest school districts.

TABLE 36. Percentage of Schools That Implemented Specific Safety and Security Measures, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2004

Site	Required visitors to report to main office or reception area	Maintained a closed campus	Used staff or adult volunteers to monitor halls	Routinely conducted bag, desk, or locker checks	Prohibited backpacks or book bags at school	Required school uniforms	Required student identification badges	Used metal detectors	Had uniformed police, undercover police, or security guards
<b>STATE SURVEYS</b>									
Alaska	89.6	43.9	75.9	27.6	13.6	4.3	1.0	2.2	12.6
Arizona	100.0	87.8	85.9	32.3	15.6	24.3	11.7	3.8	45.6
Arkansas	100.0	95.0	92.9	69.2	10.8	6.5	8.6	19.7	44.8
Connecticut	99.7	94.5	85.8	20.0	40.6	4.7	4.9	2.4	48.1
Delaware	100.0	95.0	90.8	23.4	38.0	12.5	10.5	1.7	48.1
Idaho	99.0	55.8	89.6	53.2	23.7	0.0	0.5	1.0	50.2
Iowa	98.9	72.6	85.3	38.3	37.6	0.0	0.4	0.7	15.4
Maine	98.0	82.0	72.6	17.1	30.6	0.0	0.0	0.5	30.0
Massachusetts	99.5	90.2	80.6	27.1	28.6	5.5	10.2	4.2	39.2
Michigan	100.0	86.7	88.5	49.5	44.3	8.0	8.8	3.6	32.6
Minnesota	98.8	67.3	83.5	47.3	35.6	0.9	2.4	0.8	37.8
Missouri	100.0	95.1	90.0	57.8	32.2	7.2	12.9	7.1	39.1
Montana	99.3	43.8	88.8	49.6	26.4	0.0	0.3	1.3	26.8
Nebraska	99.0	73.1	90.4	47.6	32.0	0.7	1.5	0.7	19.6
New Hampshire	100.0	83.8	75.4	20.3	24.8	0.0	2.3	1.8	36.0
New York	100.0	83.0	86.7	38.9	38.1	4.9	7.0	8.1	56.3
North Carolina	100.0	94.3	90.4	54.9	22.8	7.0	10.5	25.8	83.1
North Dakota	97.0	56.8	82.3	36.2	19.1	0.0	0.5	1.5	10.5
Oklahoma	100.0	73.8	92.8	66.9	15.0	3.6	4.6	9.7	29.6
Oregon	98.8	63.9	88.6	37.1	25.2	0.8	1.3	1.9	37.6
Pennsylvania	100.0	93.2	86.7	46.9	46.5	20.0	8.4	16.7	49.1
South Carolina	99.7	95.8	93.0	65.5	13.4	6.8	37.7	27.8	86.5
Tennessee	100.0	96.5	92.8	70.5	23.2	15.2	7.9	21.1	61.2
Texas*	100.0	84.9	94.3	65.1	24.5	13.6	15.0	12.1	59.2
Utah	98.5	59.1	91.6	46.0	22.5	2.9	1.5	0.0	63.4
Washington	99.2	73.9	85.6	29.4	15.6	2.2	3.9	3.7	41.7
Wisconsin	99.3	74.1	82.7	51.5	49.4	1.5	1.6	3.7	32.9
<b>State Median</b>	<b>99.7</b>	<b>83.0</b>	<b>88.5</b>	<b>46.9</b>	<b>25.2</b>	<b>4.3</b>	<b>4.6</b>	<b>3.6</b>	<b>39.2</b>
<b>State Range</b>	<b>89.6 – 100.0</b>	<b>43.8 – 96.5</b>	<b>72.6 – 94.3</b>	<b>17.1 – 70.5</b>	<b>10.8 – 49.4</b>	<b>0.0 – 24.3</b>	<b>0.0 – 37.7</b>	<b>0.0 – 27.8</b>	<b>10.5 – 86.5</b>
<b>LOCAL SURVEYS</b>									
Chicago	100.0	97.1	93.0	67.2	30.0	78.1	21.5	55.3	91.4
Dallas	100.0	92.2	94.0	89.8	18.0	14.3	61.5	90.0	91.9
District of Columbia	100.0	94.4	93.8	94.2	33.1	40.3	74.2	90.5	100.0
Los Angeles	100.0	96.6	96.9	92.5	2.9	37.0	10.3	92.4	79.8
Memphis	98.1	98.1	92.4	96.3	25.7	98.1	11.6	98.1	90.8
Miami	100.0	96.7	91.2	67.2	8.6	50.7	39.1	43.0	85.0
New Orleans	100.0	94.1	94.1	73.0	21.1	90.6	63.3	81.7	84.6
Orange County	100.0	97.3	93.1	35.3	10.3	5.5	41.0	7.5	94.5
Philadelphia	100.0	96.8	92.9	53.5	17.8	94.6	25.8	43.0	82.7
San Bernardino	100.0	75.0	87.5	37.5	6.3	0.0	0.0	12.5	93.8
San Diego	100.0	100.0	92.8	7.3	0.0	19.3	0.0	0.0	68.7
<b>Local Median</b>	<b>100.0</b>	<b>96.7</b>	<b>93.0</b>	<b>67.2</b>	<b>17.8</b>	<b>40.3</b>	<b>25.8</b>	<b>55.3</b>	<b>90.8</b>
<b>Local Range</b>	<b>98.1 – 100.0</b>	<b>75.0 – 100.0</b>	<b>87.5 – 96.9</b>	<b>7.3 – 96.3</b>	<b>0.0 – 33.1</b>	<b>0.0 – 98.1</b>	<b>0.0 – 74.2</b>	<b>0.0 – 98.1</b>	<b>68.7 – 100.0</b>

\* Survey did not include schools from two of the state's largest school districts.

TABLE 37. Percentage of Schools That Had or Participated in Specific Violence Prevention Programs and That Had a Written Plan for Responding to Violence at the School, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2004

Site	Violence prevention programs				Written plan to respond to violence
	Peer mediation program	Safe-passage to school program	Program to prevent gang violence	Program to prevent bullying	
<b>STATE SURVEYS</b>					
Alaska	28.4	5.6	14.6	45.9	91.7
Arizona	43.8	12.1	38.6	59.8	93.9
Arkansas	29.7	4.1	23.6	73.4	97.5
Connecticut	67.8	5.9	23.4	76.3	96.9
Delaware	53.8	6.9	15.2	48.9	94.3
Idaho	44.9	6.6	29.3	56.6	95.5
Iowa	30.7	2.8	10.2	46.8	93.8
Maine	43.6	1.6	7.2	49.0	95.8
Massachusetts	61.3	6.7	18.9	60.2	95.8
Michigan	41.9	6.9	22.5	59.0	96.0
Minnesota	46.7	5.2	20.8	62.4	95.4
Missouri	46.6	6.8	20.6	47.5	97.4
Montana	32.1	3.0	20.1	55.7	89.5
Nebraska	20.5	2.8	16.3	51.7	96.1
New Hampshire	60.0	5.2	12.3	72.9	96.3
New York	59.7	7.6	26.6	65.5	97.4
North Carolina	57.4	10.4	25.5	46.5	97.5
North Dakota	17.8	2.5	17.3	57.4	86.9
Oklahoma	22.3	11.2	26.9	69.9	95.8
Oregon	37.7	5.8	26.8	57.6	94.4
Pennsylvania	59.0	11.0	19.0	66.5	99.0
South Carolina	49.3	8.9	37.7	49.9	97.1
Tennessee	44.4	11.6	36.5	61.1	98.1
Texas*	46.6	9.8	39.8	58.2	96.6
Utah	43.9	21.0	48.2	65.9	98.0
Washington	39.0	3.8	27.9	67.6	93.8
Wisconsin	48.8	3.4	21.5	53.2	94.7
<b>State Median</b>	<b>44.4</b>	<b>6.6</b>	<b>22.5</b>	<b>58.2</b>	<b>95.8</b>
<b>State Range</b>	<b>17.8 – 67.8</b>	<b>1.6 – 21.0</b>	<b>7.2 – 48.2</b>	<b>45.9 – 76.3</b>	<b>86.9 – 99.0</b>
<b>LOCAL SURVEYS</b>					
Chicago	43.4	57.1	64.2	62.2	97.8
Dallas	62.1	20.1	54.2	48.4	98.0
District of Columbia	89.9	28.5	84.8	52.8	89.9
Los Angeles	64.5	29.6	70.0	64.6	97.7
Memphis	64.1	13.7	65.7	62.9	96.4
Miami	90.3	26.7	66.9	70.8	100.0
New Orleans	63.6	26.1	62.8	59.7	93.8
Orange County	97.3	32.9	79.0	85.8	100.0
Philadelphia	66.5	37.0	39.5	73.6	99.2
San Bernardino	56.3	26.7	81.3	60.0	93.8
San Diego	43.7	7.4	41.6	54.0	94.9
<b>Local Median</b>	<b>64.1</b>	<b>26.7</b>	<b>65.7</b>	<b>62.2</b>	<b>97.7</b>
<b>Local Range</b>	<b>43.4 – 97.3</b>	<b>7.4 – 57.1</b>	<b>39.5 – 84.8</b>	<b>48.4 – 85.8</b>	<b>89.9 – 100.0</b>

\* Survey did not include schools from two of the state's largest school districts.

TABLE 38. Percentage of Schools With a Written Policy That Protects the Rights of Students or Staff with HIV\* Infection or AIDS\*\* and, Among Those Schools, Percentage Whose Policy Addressed Specific Issues, Selected U.S. Sites: School Health Profiles, Principal Surveys, 2004

Site	Topic addressed by a written policy								
	Had a written policy	Attendance of students with HIV infection	Procedures to protect HIV-infected students and staff from discrimination	Maintenance of confidentiality of HIV-infected students and staff	Worksite safety	Confidential counseling for HIV-infected students	Communication of the policy to students, school staff, and parents	Adequate training about HIV infection for school staff	Procedures for implementing the policy
<b>STATE SURVEYS</b>									
Alaska	47.5	89.2	97.0	98.5	94.9	64.5	82.8	78.1	89.4
Arizona	52.8	93.3	98.7	99.4	98.3	75.0	85.8	86.4	93.4
Arkansas	44.9	86.0	96.6	98.9	96.6	86.5	87.3	82.2	88.6
Connecticut	70.2	89.5	96.2	99.4	98.3	82.3	80.5	86.7	86.5
Delaware	51.3	96.7	100.0	100.0	96.6	81.5	85.6	65.2	89.0
Idaho	67.9	96.2	93.8	96.9	95.5	79.0	82.2	69.6	86.6
Iowa	57.5	92.0	91.3	95.1	98.7	67.2	87.9	87.6	88.5
Maine	75.4	90.3	97.1	97.9	97.9	75.2	81.7	85.9	86.8
Massachusetts	71.5	91.2	96.8	98.5	97.5	79.3	83.8	76.2	84.9
Michigan	52.7	86.5	93.0	96.8	97.6	64.1	76.8	82.3	87.7
Minnesota	55.5	86.9	97.3	98.6	97.9	77.2	87.3	88.4	91.9
Missouri	66.0	92.6	97.7	98.6	99.0	80.4	88.8	86.2	94.7
Montana	57.1	91.9	94.4	97.5	96.3	74.1	88.0	84.6	92.5
Nebraska	58.6	90.7	94.4	97.5	98.1	77.3	87.9	82.6	92.6
New Hampshire	84.2	94.3	99.1	99.3	99.3	70.8	82.2	85.8	90.8
New York	72.7	92.9	97.7	100.0	97.3	75.3	83.9	90.7	92.8
North Carolina	51.5	88.0	94.2	99.0	99.4	84.3	83.3	89.4	92.9
North Dakota	52.3	94.5	97.3	97.3	99.1	74.0	92.2	86.4	92.3
Oklahoma	68.7	89.3	98.7	98.3	97.4	82.5	94.7	96.5	97.8
Oregon	81.5	95.1	97.9	99.6	98.9	76.7	86.4	90.1	92.0
Pennsylvania	67.1	92.2	97.9	98.7	96.1	74.1	85.0	77.9	91.9
South Carolina	67.4	92.1	97.0	98.7	100.0	78.5	83.9	95.8	93.4
Tennessee	66.7	92.3	98.5	99.5	99.5	81.4	89.5	89.3	96.3
Texas <sup>§</sup>	52.1	91.2	97.9	100.0	97.8	81.2	91.3	83.2	86.7
Utah	59.4	95.0	97.4	100.0	98.3	81.5	84.5	81.2	92.2
Washington	68.6	91.3	94.8	98.1	98.1	81.1	85.8	90.7	93.5
Wisconsin	54.4	86.5	92.8	96.7	97.3	71.0	77.1	82.5	85.1
<b>State Median</b>	<b>59.4</b>	<b>91.9</b>	<b>97.1</b>	<b>98.6</b>	<b>97.9</b>	<b>77.3</b>	<b>85.6</b>	<b>85.9</b>	<b>91.9</b>
<b>State Range</b>	<b>44.9 – 84.2</b>	<b>86.0 – 96.7</b>	<b>91.3 – 100.0</b>	<b>95.1 – 100.0</b>	<b>94.9 – 100.0</b>	<b>64.1 – 86.5</b>	<b>76.8 – 94.7</b>	<b>65.2 – 96.5</b>	<b>84.9 – 97.8</b>
<b>LOCAL SURVEYS</b>									
Chicago	54.2	95.3	99.2	99.2	100.0	85.5	93.4	78.0	92.3
Dallas	60.0	96.4	93.1	100.0	100.0	86.1	89.7	79.5	89.7
District of Columbia	100.0	94.2	97.1	100.0	97.1	100.0	96.5	90.7	86.7
Los Angeles	63.5	89.0	100.0	100.0	100.0	86.9	89.2	89.3	89.0
Memphis	64.2	84.7	100.0	100.0	100.0	84.5	87.4	81.8	87.9
Miami	79.2	88.8	100.0	100.0	100.0	91.6	93.1	90.2	93.0
New Orleans	69.4	91.5	100.0	100.0	95.6	91.1	95.5	82.7	90.9
Orange County	65.3	91.5	95.1	100.0	95.1	90.1	85.2	90.1	95.1
Philadelphia	56.1	93.9	96.9	98.4	100.0	91.0	88.0	78.8	88.0
San Bernardino	100.0	100.0	100.0	100.0	100.0	100.0	85.7	92.9	100.0
San Diego	83.4	96.9	100.0	100.0	100.0	87.8	87.6	90.9	96.9
<b>Local Median</b>	<b>65.3</b>	<b>93.9</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>90.1</b>	<b>89.2</b>	<b>89.3</b>	<b>90.9</b>
<b>Local Range</b>	<b>54.2 – 100.0</b>	<b>84.7 – 100.0</b>	<b>93.1 – 100.0</b>	<b>98.4 – 100.0</b>	<b>95.1 – 100.0</b>	<b>84.5 – 100.0</b>	<b>85.2 – 96.5</b>	<b>78.0 – 92.9</b>	<b>86.7 – 100.0</b>

\* HIV = human immunodeficiency virus.

\*\* AIDS = acquired immunodeficiency syndrome.

§ Survey did not include schools from two of the state's largest school districts.

TABLE 39. Percentage of Schools That Had a School Health Committee or Advisory Group\* and Engaged Parents and Families in Specific Health Education Activities During the 2003–2004 School Year, Selected U.S. Sites: School Health Profiles, Principal and Lead Health Education Teacher Surveys, 2004

Site	School health committee or advisory group	Provided families with information on the health education program	Met with a parents' organization to discuss the health education program	Invited family members to attend a health education class
<b>STATE SURVEYS</b>				
Alaska	47.5	55.6	20.0	36.7
Arizona	36.8	54.4	19.5	29.8
Arkansas	30.5	58.6	15.1	37.0
Connecticut	48.7	79.1	29.1	31.9
Delaware	50.7	77.7	24.1	47.6
Idaho	31.8	63.4	11.8	40.6
Iowa	43.7	61.4	7.8	29.2
Maine	65.0	75.3	14.5	34.4
Massachusetts	61.6	81.4	33.6	29.5
Michigan	72.7	77.7	26.0	48.8
Minnesota	49.9	NA**	NA	NA
Missouri	60.7	67.8	16.8	30.0
Montana	31.9	57.1	11.3	34.1
Nebraska	28.5	50.4	4.3	27.9
New Hampshire	36.9	77.5	20.6	29.1
New York	58.7	80.0	28.5	29.0
North Carolina	66.6	74.9	25.5	45.6
North Dakota	36.5	52.7	12.9	26.4
Oklahoma	20.5	37.2	13.9	18.6
Oregon	41.7	77.6	12.0	40.0
Pennsylvania	45.1	70.9	16.3	35.5
South Carolina	69.6	65.8	25.1	43.1
Tennessee	46.3	54.5	24.7	33.8
Texas <sup>§</sup>	67.4	NA	NA	NA
Utah	50.7	74.9	18.3	54.6
Washington	55.3	72.5	19.9	34.6
Wisconsin	56.7	75.8	13.6	41.0
<b>State Median</b>	<b>48.7</b>	<b>70.9</b>	<b>18.3</b>	<b>34.4</b>
<b>State Range</b>	<b>20.5 – 72.7</b>	<b>37.2 – 81.4</b>	<b>4.3 – 33.6</b>	<b>18.6 – 54.6</b>
<b>LOCAL SURVEYS</b>				
Chicago	40.9	58.5	32.9	31.0
Dallas	62.6	69.2	37.5	43.5
District of Columbia	75.8	83.8	51.8	87.5
Los Angeles	63.9	81.0	32.1	59.3
Memphis	69.0	89.8	56.7	71.6
Miami	72.3	65.5	26.8	30.3
New Orleans	73.0	75.0	37.8	64.6
Orange County	75.5	80.0	20.5	27.2
Philadelphia	75.0	64.4	24.5	41.3
San Bernardino	57.1	NA	NA	NA
San Diego	85.3	87.7	18.7	34.0
<b>Local Median</b>	<b>72.3</b>	<b>77.5</b>	<b>32.5</b>	<b>42.4</b>
<b>Local Range</b>	<b>40.9 – 85.3</b>	<b>58.5 – 89.8</b>	<b>18.7 – 56.7</b>	<b>27.2 – 87.5</b>

\* That develops policies, coordinates activities, or seeks student and family involvement in programs that address health issues.

\*\* NA = data not available.

§ Survey did not include schools from two of the state's largest school districts.

TABLE 40. Among Schools That Required a Health Education Course in Any of Grades 6–12, the Percentage of Schools That Asked Students to Participate in Health-Related Community Activities as Part of a Required Health Education Course During the 2003–2004 School Year, Selected U.S. Sites: School Health Profiles, Lead Health Education Teacher Surveys, 2004

Site	Performed volunteer work*	Participated in or attended a health fair	Gathered information about health services**	Compared prices of health products	Identified potential injury sites <sup>§</sup>	Identified and analyzed advertising <sup>§§</sup>	Advocated for a health-related issue	Completed homework with family members
<b>STATE SURVEYS</b>								
Alaska	15.8	37.3	57.1	13.4	49.4	48.9	46.1	66.6
Arizona	24.2	26.7	48.9	26.7	49.0	66.3	48.3	62.3
Arkansas	16.2	21.2	55.5	36.7	66.3	66.0	47.6	73.3
Connecticut	15.9	28.6	56.4	19.0	48.9	65.1	51.2	76.9
Delaware	13.5	36.2	71.1	30.0	66.6	82.4	66.2	80.2
Idaho	12.7	27.1	55.7	35.2	62.0	71.9	51.3	78.1
Iowa	16.8	13.8	47.0	23.1	45.6	62.2	45.6	80.9
Maine	10.0	21.3	70.2	27.7	54.4	68.1	49.0	75.1
Massachusetts	14.8	19.7	54.4	23.9	54.6	73.6	57.9	76.7
Michigan	14.4	14.3	47.9	20.9	47.7	70.0	50.8	84.9
Missouri	13.9	36.1	53.8	23.6	67.3	72.2	41.7	78.3
Montana	14.0	22.8	49.5	22.4	56.5	59.6	53.5	73.0
Nebraska	9.5	23.0	46.4	17.9	53.0	57.6	45.0	81.3
New Hampshire	12.7	13.3	50.7	22.5	46.7	73.7	52.3	76.5
New York	18.1	29.5	71.6	31.2	65.5	78.2	70.0	80.5
North Carolina	16.0	30.7	60.9	26.6	71.2	74.5	56.8	71.8
North Dakota	12.0	34.5	57.4	21.4	55.1	66.7	48.8	77.9
Oklahoma	10.9	35.9	45.5	16.5	63.4	49.2	35.7	66.3
Oregon	23.6	19.0	60.0	27.4	52.4	71.7	53.2	87.1
Pennsylvania	20.9	29.9	57.5	25.9	63.3	76.7	47.4	77.4
South Carolina	19.2	29.1	63.8	26.3	59.1	69.3	52.1	74.1
Tennessee	19.6	37.9	65.3	34.6	67.6	70.9	50.7	74.4
Utah	19.2	22.4	45.8	29.8	58.4	72.7	50.3	87.6
Washington	21.4	17.6	46.2	18.4	36.5	62.0	44.2	76.4
Wisconsin	18.9	22.8	57.0	25.2	52.8	70.9	50.9	79.3
<b>State Median</b>	<b>15.9</b>	<b>26.7</b>	<b>55.7</b>	<b>25.2</b>	<b>55.1</b>	<b>70.0</b>	<b>50.7</b>	<b>76.9</b>
<b>State Range</b>	<b>9.5 – 24.2</b>	<b>13.3 – 37.9</b>	<b>45.5 – 71.6</b>	<b>13.4 – 36.7</b>	<b>36.5 – 71.2</b>	<b>48.9 – 82.4</b>	<b>35.7 – 70.0</b>	<b>62.3 – 87.6</b>
<b>LOCAL SURVEYS</b>								
Chicago	19.1	36.2	50.2	31.8	63.4	56.9	46.0	79.7
Dallas	30.6	47.8	74.8	41.7	83.9	87.4	65.1	80.1
District of Columbia	46.4	78.6	78.3	47.1	62.7	71.9	56.6	89.1
Los Angeles	26.2	44.6	66.0	43.6	58.9	84.8	58.8	88.1
Memphis	39.4	54.9	88.1	57.4	75.8	78.9	76.0	94.0
Miami	29.5	34.0	68.1	29.4	62.7	67.9	49.8	81.7
New Orleans	34.7	44.9	71.3	44.7	69.1	68.6	51.8	72.4
Orange County	11.7	44.4	71.6	40.1	65.4	79.6	55.6	92.0
Philadelphia	27.1	43.5	66.0	47.0	67.4	70.7	53.6	86.8
San Diego	NA <sup>†</sup>	NA	NA	NA	NA	NA	NA	NA
<b>Local Median</b>	<b>29.5</b>	<b>44.6</b>	<b>71.3</b>	<b>43.6</b>	<b>65.4</b>	<b>71.9</b>	<b>55.6</b>	<b>86.8</b>
<b>Local Range</b>	<b>11.7 – 46.4</b>	<b>34.0 – 78.6</b>	<b>50.2 – 88.1</b>	<b>29.4 – 57.4</b>	<b>58.9 – 83.9</b>	<b>56.9 – 87.4</b>	<b>46.0 – 76.0</b>	<b>72.4 – 94.0</b>

\* At a hospital, a local health department, or any other community organization that addresses health issues.  
 \*\* That are available in the community.  
 § At school, home, or in the community.  
 §§ In the community designed to influence health behaviors or health risk behaviors.  
 † NA = data not available.







