Philadelphia, Pennsylvania – 2003 (N=300)

Figure A. Age of GISP participants, in years, 2003



Figure B. Race/ethnicity of GISP participants, 2003



Figure C. Percentage of GISP participants identifying as men who have sex with men, 1988–2003



Figure D. Reason for visit among GISP participants, 2003



Philadelphia, Pennsylvania – 2003 (N=300)

Figure E. Previous episode of gonorrhea among GISP participants, 1991–2003



*Data first collected in 1991.
**Data first collected in 1992.
Note: Data points not shown when >30% data missing.

Figure F. Drugs used to treat gonorrhea among GISP participants, 2003



Figure G. Drugs used to treat *Chlamydia trachomatis* infection among GISP participants, 2003



Figure H. Resistance to penicillin and tetracycline among GISP isolates, 2003



Philadelphia, Pennsylvania – 2003 (N=300)

Figure I. Decreased susceptibility to ceftriaxone among GISP isolates, 1988–2003



Figure J. Decreased susceptibility to cefixime among GISP isolates, 1992–2003



Figure K. Intermediate resistance and resistance to ciprofloxacin among GISP isolates, 1990–2003



Note: Susceptibility to ciprofloxacin first measured in 1990.

Figure L. Decreased susceptibility to azithromycin among GISP isolates, 1992–2003



Note: Susceptibility to azithromycin first measured in 1992.
Note: Decreased susceptibility to azithromycin is defined here as ≥ 1.0 μg/ml.
No NCCLS criteria currently exist.