

Portland, Oregon – 2003 (N=132)

Figure A. Age of GISP participants, in years, 2003




Figure B. Race/ethnicity of GISP participants, 2003




Figure C. Percentage of GISP participants identifying as men who have sex with men, 1988-2003






Figure D. Reason for visit among GISP participants, 2003



Portland, Oregon – 2003 (N=132)

Figure E. Previous episode of gonorrhoea among GISP participants, 1991 – 2003



*Data first collected in 1991.

**Data first collected in 1992.

Note: Data points not shown when >30% data missing.

Figure F. Drugs used to treat gonorrhoea among GISP participants, 2003




Figure G. Drugs used to treat *Chlamydia trachomatis* infection among GISP participants, 2003





Figure H. Resistance to penicillin and tetracycline among GISP isolates, 2003



Portland, Oregon – 2003 (N=132)

Figure I. Decreased susceptibility to ceftriaxone among GISP isolates, 1988–2003

No isolates with decreased susceptibility to ceftriaxone have been identified at this clinic.

Figure J. Decreased susceptibility to cefixime among GISP isolates, 1992–2003

No isolates with decreased susceptibility to cefixime have been identified at this clinic.

Figure K. Intermediate resistance and resistance to ciprofloxacin among GISP isolates, 1990–2003




Figure L. Decreased susceptibility to azithromycin among GISP isolates, 1992–2003

No isolates with decreased susceptibility to azithromycin have been identified at this clinic.

Note: Susceptibility to ciprofloxacin first measured in 1990.