

Portland, Oregon – 2004 (N=191)

Figure A. Age of GISP participants, in years, 2004




Figure B. Race/ethnicity of GISP participants, 2004




Figure C. Percentage of GISP participants identifying as men who have sex with men, 1988–2004






Figure D. Reason for visit among GISP participants, 2004



Portland, Oregon – 2004 (N=191)

Figure E. Previous episode of gonorrhea among GISP participants, 1991–2004



*Data first collected in 1991.

**Data first collected in 1992.

Note: Data points not shown when >30% data missing.

Figure F. Drugs used to treat gonorrhea among GISP participants, 2004




Figure G. Drugs used to treat *Chlamydia trachomatis* infection among GISP participants, 2004





Figure H. Resistance to penicillin and tetracycline among GISP isolates, 2004



Portland, Oregon – 2004 (N=191)

Figure I. Decreased susceptibility to ceftriaxone among GISP isolates, 1988 – 2004

No isolates with decreased susceptibility to ceftriaxone have been identified at this clinic.

Figure J. Decreased susceptibility to cefixime among GISP isolates, 1992 – 2004

No isolates with decreased susceptibility to cefixime have been identified at this clinic.

Figure K. Intermediate resistance and resistance to ciprofloxacin among GISP isolates, 1990 – 2004




Figure L. Decreased susceptibility to azithromycin among GISP isolates, 1992 – 2004

No isolates with decreased susceptibility to azithromycin have been identified at this clinic.

Note: Susceptibility to ciprofloxacin first measured in 1990.