

Adolescent Behaviors and Experiences Survey														
Dietary Behaviors	American Indian or Alaska Native*		Asian*		Black or African American*		Hispanic or Latino*		Native Hawaiian or Other Pacific Islander*		White*		Multiple Race*	
	Percentage	Confidence Interval	Percentage	Confidence Interval	Percentage	Confidence Interval	Percentage	Confidence Interval	Percentage	Confidence Interval	Percentage	Confidence Interval	Percentage	Confidence Interval
Did not drink fruit juice (100% fruit juices one or more times during the 7 days before the survey)	27.6	22.0 - 34.0	44.9	40.3 - 49.7	30.0	26.4 - 33.9	37.1	34.6 - 39.7	N/A	N/A	43.6	41.1 - 46.2	35.1	28.7 - 42.1
Did not eat fruit (one or more times during the 7 days before the survey)	2.6	1.5 - 4.5	9.5	6.4 - 13.8	20.2	17.0 - 23.8	12.3	10.5 - 14.3	N/A	N/A	15.1	11.2 - 20.2	11.8	8.4 - 16.2
Did not eat green salad (one or more times during the 7 days before the survey)	46.7	29.6 - 64.6	42.8	31.2 - 55.3	57.3	52.8 - 61.7	51.3	48.0 - 54.6	N/A	N/A	44.8	40.5 - 49.2	48.1	40.6 - 55.6
Did not eat potatoes (one or more times during the 7 days before the survey)	36.3	29.6 - 43.5	43.6	35.5 - 52.1	46.2	41.6 - 50.8	38.2	35.2 - 41.3	N/A	N/A	33.6	31.3 - 35.9	39.4	32.4 - 46.9
Did not eat carrots (one or more times during the 7 days before the survey)	57.0	47.1 - 66.3	49.1	43.7 - 54.4	73.6	69.8 - 77.1	56.0	51.3 - 60.6	N/A	N/A	57.3	53.4 - 61.2	53.0	46.5 - 59.4
Did not eat other vegetables (one or more times during the 7 days before the survey)	18.8	13.1 - 26.3	8.4	4.9 - 13.8	27.2	23.8 - 30.9	19.0	15.7 - 22.8	N/A	N/A	17.6	14.0 - 22.1	13.2	9.4 - 18.3
Did not eat fruit or drink 100% fruit juices (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)	2.0	1.3 - 3.0	6.4	3.8 - 10.5	10.7	8.2 - 14.0	6.9	5.3 - 8.8	N/A	N/A	10.3	7.4 - 14.0	5.7	3.5 - 9.3
Ate fruit or drank 100% fruit juices one or more times per day (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)	62.7	51.1 - 73.0	49.2	43.3 - 55.1	53.1	48.3 - 57.8	49.1	44.5 - 53.8	N/A	N/A	44.4	40.5 - 48.4	49.7	43.2 - 56.2
Ate fruit or drank 100% fruit juices two or more times per day (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)	34.8	27.2 - 43.2	24.1	18.6 - 30.7	28.4	24.5 - 32.6	23.2	19.4 - 27.5	N/A	N/A	19.1	16.8 - 21.8	22.1	17.5 - 27.5
Did not eat vegetables (green salad, potatoes [excluding french fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	6.3	3.5 - 10.9	5.0	3.1 - 7.9	13.2	9.7 - 17.7	8.3	6.6 - 10.2	N/A	N/A	7.1	5.4 - 9.3	6.1	4.2 - 8.9
Ate vegetables one or more times per day (green salad, potatoes [excluding french fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	48.4	34.6 - 62.4	65.1	55.2 - 73.7	44.0	39.8 - 48.2	51.7	47.7 - 55.6	N/A	N/A	57.9	54.4 - 61.4	60.3	54.2 - 66.1
Ate vegetables two or more times per day (green salad, potatoes [excluding french fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	22.6	16.3 - 30.3	35.0	27.3 - 43.7	19.4	16.3 - 22.9	19.9	17.3 - 22.9	N/A	N/A	21.6	19.3 - 24.1	24.7	19.6 - 30.7
Ate vegetables three or more times per day (green salad, potatoes [excluding french fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	19.7	13.4 - 28.1	17.0	12.0 - 23.6	11.3	9.3 - 13.8	11.3	9.4 - 13.5	N/A	N/A	10.5	8.9 - 12.3	13.0	9.2 - 18.0
Drank a can, bottle, or glass of soda or pop one or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	18.6	13.3 - 25.4	3.1	1.7 - 5.5	15.5	12.9 - 18.3	11.7	9.8 - 13.9	N/A	N/A	20.1	14.6 - 27.0	9.9	6.0 - 15.7
Did not drink a can, bottle, or glass of soda or pop (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, one or more times during the 7 days before the survey)	39.6	28.6 - 51.7	46.8	35.7 - 58.1	37.8	34.0 - 41.8	35.0	31.3 - 38.8	N/A	N/A	33.1	28.9 - 37.6	33.4	28.0 - 39.4
Drank a can, bottle, or glass of soda or pop two or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	14.3	11.2 - 18.1	1.7	1.0 - 2.9	10.2	7.9 - 13.2	6.4	5.3 - 7.7	N/A	N/A	12.2	8.0 - 18.0	5.4	3.0 - 9.6
Did not drink milk (during the 7 days before the survey)	32.3	22.9 - 43.4	36.8	31.0 - 43.0	50.7	45.8 - 55.7	29.9	26.4 - 33.8	N/A	N/A	32.8	29.8 - 36.0	37.4	30.3 - 45.1
Drank one or more glasses per day of milk (counting the milk they drank in a glass or cup, from a carton, or with cereal and counting the half pint of milk served at school as equal to one glass, during the 7 days before the survey)	18.7	12.4 - 27.0	30.9	23.8 - 39.1	14.2	11.8 - 17.0	24.2	21.7 - 26.9	N/A	N/A	25.2	22.6 - 27.9	16.8	11.2 - 24.6
Drank three or more glasses per day of milk (counting the milk they drank in a glass or cup, from a carton, or with cereal and counting the half pint of milk served at school as equal to one glass, during the 7 days before the survey)	3.0	1.2 - 7.4	2.8	1.2 - 6.2	3.5	2.3 - 5.4	3.7	2.7 - 5.0	N/A	N/A	5.9	4.7 - 7.4	4.0	2.2 - 7.3
Did not eat breakfast (during the 7 days before the survey)	11.8	5.4 - 23.8	15.9	12.4 - 20.1	17.0	13.6 - 21.0	11.2	9.4 - 13.3	N/A	N/A	16.4	14.4 - 18.7	13.3	9.4 - 18.6
Ate breakfast on all 7 days (during the 7 days before the survey)	22.3	16.9 - 28.8	34.4	23.5 - 47.3	25.1	21.2 - 29.5	33.1	28.5 - 38.1	N/A	N/A	27.9	25.3 - 30.6	20.7	16.6 - 25.4
Did not drink a can, bottle, or glass of a sports drink (such as Gatorade or PowerAde, not counting low-calorie sports drinks such as Propel or G2, during the 7 days before the survey)	31.4	26.8 - 36.3	79.0	73.4 - 83.6	51.9	47.2 - 56.5	56.8	52.6 - 60.9	N/A	N/A	51.8	48.0 - 55.5	57.8	48.9 - 66.2
Drank a can, bottle, or glass of a sports drink one or more times per day (such as Gatorade or PowerAde, not counting low calorie sports drinks such as Propel or G2, during the 7 days before the survey)	19.2	11.0 - 31.4	2.8	1.2 - 6.5	14.2	11.4 - 17.4	7.4	5.9 - 9.3	N/A	N/A	11.2	9.2 - 13.5	7.0	3.9 - 12.4
Drank a can, bottle, or glass of a sports drink two or more times per day (such as Gatorade or PowerAde, not counting low calorie sports drinks such as Propel or G2, during the 7 days before the survey)	8.3	4.4 - 15.0	0.6	0.2 - 1.6	9.0	6.7 - 12.0	4.2	3.1 - 5.7	N/A	N/A	5.7	4.3 - 7.5	4.0	2.4 - 6.7
Did not drink a bottle or glass of plain water (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)	7.1	4.0 - 12.2	4.3	1.9 - 9.6	7.8	5.2 - 11.4	2.6	2.0 - 3.5	N/A	N/A	3.7	2.6 - 5.1	2.9	1.5 - 5.5
Drank a bottle or glass of plain water one or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)	68.3	57.7 - 77.2	80.6	72.5 - 86.7	69.4	63.6 - 74.6	74.9	72.6 - 77.2	N/A	N/A	77.5	73.3 - 81.2	75.0	65.6 - 82.5
Drank a bottle or glass of plain water two or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)	63.5	50.7 - 74.6	75.6	67.9 - 81.9	63.7	58.0 - 69.1	69.7	66.3 - 73.0	N/A	N/A	68.2	63.6 - 72.4	69.3	60.3 - 77.1
Drank a bottle or glass of plain water three or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)	55.4	44.0 - 66.3	65.2	59.8 - 70.2	52.3	46.7 - 57.8	55.2	51.3 - 59.0	N/A	N/A	51.6	47.2 - 56.1	57.7	49.6 - 65.4

*Non-Hispanic

N/A: Less than 30 respondents for the subgroup