

Maternal/Pediatric Respiratory Syncytial Virus (RSV) Work Group

Sarah S. Long, MD Chair, Maternal/Pediatric RSV Work Group

ACIP Meeting

June 28, 2024

Two products are recommended to protect infants and young children from RSV lower respiratory tract disease

- RSV prevention in infants during their first RSV season (i.e., aged <8 months)
 - To protect infants in their first season, either maternal RSV vaccination (Abrysvo, Pfizer), or use of nirsevimab (Beyfortus, Sanofi and AstraZeneca) in the infant is recommended to prevent RSV lower respiratory tract disease
 - Administration of both products is not needed for most infants
- Children ages 8–19 months who are at increased risk of severe RSV disease and entering their second RSV season are recommended to receive one dose of nirsevimab

Timing of RSV vaccine and nirsevimab




Figure represents recommended timing of immunization product deployment for most of the continental U.S. In jurisdictions with seasonality that differs from most of the continental United States (e.g., Alaska, jurisdictions with tropical climates), providers should follow state, local, or territorial guidance on timing of administration

Timing of administration of maternal Pfizer RSV vaccine (Abrysvo) and nirsevimab (Beyfortus)

Maternal RSV vaccine

- For pregnant people at 32–36 weeks' gestation, vaccinate using seasonal administration (meaning **September–January** in most of the United States)

• Nirsevimab¹

- For infants <8 months of age, immunize shortly before season onset (e.g., October)
- For infants born during **October–March**, immunize within 1 week of birth

https://www.cdc.gov/mmwr/volumes/72/wr/mm7234a4.htm; https://www.cdc.gov/mmwr/volumes/72/wr/mm7241e1.htm

¹In jurisdictions with seasonality that differs from most of the continental United States (e.g., Alaska, jurisdictions with tropical climates), providers should follow state, local, or territorial guidance on timing of administration

Vaccine administration errors reported in young children and pregnant people



COCA Now

CDC Clinician Outreach and Communication Activity

January 22, 2024

Information on Respiratory Syncytial Virus (RSV) Vaccine Administration Errors in Young Children and Pregnant People

Information on Respiratory Syncytial Virus (RSV) Vaccine Administration Errors in Young Children and Pregnant People

Administration of the GSK Respiratory Syncytial Virus Vaccine to Pregnant Persons in Error - PubMed (nih.gov)

Incorrect Administration of Adult RSV Vaccines to Young Children - PubMed (nih.gov)

- Multiple products introduced in the same season created confusion among healthcare workers as to which groups were eligible for which immunizations
- Rare reports of administration errors were received by the Vaccine Adverse Events Reporting System (VAERS)
 - Reports of pregnant persons receiving Arexvy (GSK's vaccine for older adults)
 - Reports of children receiving RSV vaccine instead of monoclonal antibody products (e.g., nirsevimab)
 - Wrong dose (nirsevimab) administered
- CDC made additional resources available and increased education efforts with healthcare providers

Work group members (external)

ACIP Members

Sarah Long (chair)

Oliver Brooks

Camille Kotton

Denise Jamieson

<u>Liaisons</u>

James McAuley (IDSA) Brenna L. Hughes (ACOG) Nicole Chaisson (AAFP) Sean O'Leary (AAP) Jennifer Schuster (PIDS) Molly Howell (AIM) Dana DeShon (NAPNAP)

Ex Officio Members

Lucia Lee (FDA-CBER) Judy Beeler (FDA-CBER) Yodit Belew (FDA-CDER) Yugenia Hong-Nguyen (FDA-CDER) Sonnie Kim (NIH-NIAID) April Killikelly (Public Health Agency of Canada) Winnie Siu (Public Health Agency of Canada) Jessica Lee (CMS/CMCS) Terry Dalle-Tezze (HRSA) Matthew Clark (IHS)

Consultants

Cody Meissner (Dartmouth Geisel School of Medicine) Helen Chu (University of Washington) Natasha Halasa (Vanderbilt University) Daniel Feikin (World Health Organization) Kevin Ault (Western Michigan University) Pablo Sanchez (Nationwide Children's Hospital)

GRADE/EtR Consultants

Doug Campos-Outcalt Rebecca Morgan

Work group members (CDC)

<u>CDC</u>

Katherine Fleming-Dutra (co-lead)AmJefferson Jones (co-lead)ChrDanielle MouliaTamMeredith McMorrowAngMila PrillMicNatalie ThornburgAmAron HallNodIsmael Ortega-SanchezClaiMelissa CoughlinFiorJamison PikePraLauren RoperAm

Amber Winn Chris Taylor Tami Skoff Angie Campbell Michael Melgar Amanda Payne Noelle Molinari Claire Midgley Fiona Havers Pragna Patel Amadea Britton

Ruth Link-Gelles Megan Wallace Monica Godfrey Karen Broder Naomi Tepper Heidi Moline Hannah Rosenblum **Derrell Powers Raigan Wheeler** Sally Ezra Manisha Patel

Demorah Hayes Elizabeth Greene Monica Patton Jarrett Gartin Dennis Wang Jordan Singleton Trang Nguyen Wisard Fatimah Dawood John Su Michael McNeil

CDC ACIP Staff

A. Patricia Wodi

Christine Olson

Andrew Leidner

Anne Hause

David Shay

Sarah Meyer

Julianne Gee

Agustin Lopez

Lakshmi Panagiotakopoulos

Kristen Folsom

Melinda Wharton Stephanie Thomas Jessica MacNeil

Agenda

- Implementation and uptake of nirsevimab and maternal RSV vaccine Dr. Shannon Stokley (CDC/NCIRD)
- Maternal RSV vaccine safety surveillance Dr. Pedro Moro (CDC/NCEZID)
- Summary of effectiveness of nirsevimab in infants Dr. Amanda Payne (CDC/NCIRD)
- Work Group considerations Dr. Jefferson Jones (CDC/NCIRD)

Thank you

For more information, contact CDC 1-800-CDC-INFO (232-4636) TTY: 1-888-232-6348 cdc.gov

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the U.S. Centers for Disease Control and Prevention.

