



# Effectiveness of COVID-19 vaccines

**Ruth Link-Gelles, PhD, MPH**

CDR, US Public Health Service

Vaccine Effectiveness Program Lead

Coronavirus and Other Respiratory Viruses Division

Centers for Disease Control and Prevention

October 23, 2024

# Agenda – COVID-19 vaccine effectiveness (VE)

- **VE data to inform need for:**
  - Additional doses in immunocompromised
    - What's known about VE in immunocompromised vs. non-immunocompromised adults, including waning by time since dose
  - Additional dose for adults  $\geq 65$  years
    - What's known about waning of a single dose in healthy adults  $\geq 65$  years, including against more severe outcomes
    - Benefits of an additional dose in past seasons

# Some key issues in COVID-19 vaccine effectiveness (VE)

## 1. Time since dose impacts protection

- How much and over what time period?

## 2. SARS-CoV-2 variants change over time


- Variant/vaccine match may impact effectiveness

## 3. Surges in disease, seroprevalence, and time since last SARS-CoV-2 infection impact measured VE

Difficult to disentangle time since dose (true “waning”) vs. impact of changes in variants vs. impact of time since prior infection

## Context for interpreting COVID-19 VE across age groups: SARS-CoV-2 seroprevalence before 2023-2024 respiratory virus season

- High rates of SARS-CoV-2 infection-induced immunity by July – August 2023.\*



VE findings should be interpreted as the added benefit provided by COVID-19 vaccination in a population with a high prevalence of vaccine- and infection-induced immunity.

\* Internal CDC data. Data on persons aged ≥16 years is from a longitudinal, national cohort of >35,000 blood donors.

Methods and prior data available at: <https://covid.cdc.gov/covid-data-tracker/#nationwide-blood-donor-seroprevalence-2022>

# Measuring COVID-19 VE

Measure	Definition	Example vaccinated group	Example comparison group
<b>Absolute VE</b>	Compares frequency of health outcomes in vaccinated and unvaccinated people	Received <b>original monovalent COVID-19 vaccine</b>	Received no COVID-19 vaccines ever
<b>Relative VE</b>	Compares frequency of health outcomes in people who received one type of vaccine to people who received a different vaccine	Received <b>bivalent COVID-19 vaccine</b>	Eligible for, but did not receive, bivalent COVID-19 vaccine, but received <b>original monovalent COVID-19 vaccine</b>
<b>VE of 2023-2024 COVID-19 vaccines</b>	Compares people who received <b>2023-2024</b> COVID-19 vaccine to people who did not, regardless of past vaccination	Received <b>updated (2023-24) dose</b>	Eligible for, but did not receive, an <b>updated (2023-24) dose</b> , regardless of past vaccination history

**Methods**

# VISION Multi-Site Network of Electronic Health Records


>300 emergency rooms and urgent cares and >200 hospitals

- **Design:** Test-negative design
- **Population:** Adults visiting a participating emergency department or urgent care (ED/UC) or hospitalized with COVID-19-like illness (CLI) with a SARS-CoV-2 NAAT test result within 10 days before or 72 hours after encounter
  - **Cases:** CLI with *positive* NAAT for SARS-CoV-2 and no positive NAAT for RSV or influenza
  - **Controls:** CLI with *negative* NAAT for SARS-CoV-2 and no positive NAAT for influenza
- **Vaccination data:** Documented by electronic health records and state and city registries



# IVY Network — 26 hospitals, 20 U.S. States

- **Design:** Test-negative, case-control design
- **Population:** Adults aged  $\geq 18$  years hospitalized with COVID-like illness (CLI)\* and SARS-CoV-2 test results within 10 days of illness onset and 3 days of admission
  - **Cases:** CLI and test *positive* for SARS-CoV-2 by NAAT or antigen
    - Co-infections with influenza and RSV are excluded
  - **Controls:** CLI and test *negative* for SARS-CoV-2 and influenza by RT-PCR
- **Vaccination data:** Electronic medical records (EMR), state and city registries, and plausible self-report
- **Specimens:** Nasal swabs obtained on all patients for central RT-PCR testing and whole genome sequencing




\*CLI is defined as presence of any one of the following: fever, cough, shortness of breath, chest imaging consistent with pneumonia, or hypoxemia



# Medicare data

- **Design:** Retrospective cohort
- **Data source:** Medicare fee-for-service claims data\*
- **Population:** Persons aged  $\geq 65$
- **Censoring events:**
  - COVID-19-related thromboembolic event
  - Death
  - Disenrollment in Medicare Parts A/B
  - Enrollment in Medicare Part C
  - Nursing home stay lasting  $\geq 100$  days
  - Admission to hospice facility
  - Dialysis encounter
  - Receipt of multiple bivalent booster doses
  - Bivalent booster dose < 60 days from the last COVID-19 vaccine dose
  - End of study period
- **VE =  $(1 - \text{adjusted hazard ratio}^{**}) \times 100\%$**

where adjusted hazard ratio =  $\frac{\text{rate of RSV hospitalization}_{\text{vaccinated}}}{\text{rate of RSV hospitalization}_{\text{unvaccinated}}}$




\*Data sources included Medicare Enrollment Database (EDB) and Common Medicare Environment (CME), Common Working File (CWF) and Shared System Data (SSD) Medicare Parts A/B claims data, Minimum Data Set (MDS), and CDC/ATSDR Social Vulnerability Index (SVI). Events identified from Medicare claims data using International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10 CM) and common procedural terminology (CPT) codes.

\*\*Hazard ratios adjusted for race, age group, bivalent COVID-19 vaccination status, 2022/2023 influenza vaccination status.


**COVID-19 VE data to inform need for  
additional doses in persons with  
immunocompromise**

## VISION: Original monovalent vs. unvaccinated mRNA VE for hospitalization by number of doses and time since last dose receipt for adults ≥50 years, Dec 2021–Mar 2022, by immunocompromised status




VE of original mRNA doses vs. unvaccinated persons during Delta and early Omicron was higher among non-immunocompromised adults. An additional dose restored/increased protection in immunocompromise and non-immunocompromised, with some waning apparent.

## VISION: Original monovalent vs. unvaccinated mRNA VE for hospitalization by number of doses and time since last dose receipt for adults ≥50 years, Dec 2021–Mar 2022, by immunocompromised status




VE of original mRNA doses vs. unvaccinated persons during Delta and early Omicron was higher among non-immunocompromised adults. An additional dose restored/increased protection in immunocompromise and non-immunocompromised, with some waning apparent.

IVY: Bivalent vs. unvaccinated mRNA VE against **hospitalization** by time since last dose receipt for adults  $\geq 18$  years, **September 2022–August 2023**, by immunocompromised status




A dose of bivalent vaccine added protection among those >1 year from their last original monovalent dose during Omicron predominance.

IVY: *Bivalent vs. unvaccinated* mRNA VE against **hospitalization** by time since last dose receipt for adults  $\geq 18$  years, **September 2022–August 2023**, by immunocompromised status



A dose of bivalent vaccine added protection among those >1 year from their last original monovalent dose during Omicron predominance.


IVY: *Bivalent vs. unvaccinated* mRNA VE against **hospitalization** by time since last dose receipt for adults  $\geq 18$  years, **September 2022–August 2023**, by immunocompromised status



A dose of bivalent vaccine added protection among those >1 year from their last original monovalent dose during Omicron predominance.

# VISION: VE of 2023-2024 vs. no 2023-2024 COVID-19 vaccine against *hospitalization* among adults aged $\geq 18$ years, by immunocompromise status

September 2023 – August 2024



During the 2023-2024 season, VE appeared somewhat lower in immunocompromised individuals, though waning patterns were similar to those in non-immunocompromised individuals.




# Conclusions: VE among those with immunocompromising conditions

- COVID-19 vaccines provided protection for both persons with and without immunocompromise.
- Patterns of COVID-19 VE in immunocompromised were different season-to-season, with generally lower VE compared to non-immunocompromised, but inconsistent waning patterns.
  - During 2023-2024, VE against hospitalization in immunocompromised waned to 0 by ~4-6 months.
- **This inconsistency is likely multifactorial, including:**
  - Heterogeneity among those classified as immunocompromised
  - Variation in underlying immunity and response to prior infection
  - Differing health behaviors (e.g., masking, social distancing) over time and by immunocompromise status

**COVID-19 VE data to inform need for an additional COVID-19 vaccine dose for adults aged  $\geq 65$  years**

# COVID-19 VE against **hospitalization** wanes over time, but is more sustained against **critical illness**, though some waning is evident

Data from VISION and IVY showing VE by vaccine formulation of most recent dose.



Recipients of bivalent and 2023-2024 doses included in this analysis received a single dose of the most recent formulation.

Sources: DeCuir, et al., MMWR 2023/Link-Gelles, ACIP Slides, April 20, 2022; CDC unpublished data updated from Link-Gelles, ACIP Slides, June 23, 2023; CDC unpublished data updated from Link-Gelles, ACIP Slides, June 27, 2024

\* For original monovalent doses, VE for hospitalization was for 180-364 days from last dose combined. For 2023-2024 doses, VE for hospitalization was for  $\geq 180$  days combined, with a median of 228 days (IQR: 202-259).

# IVY: Original monovalent vs. unvaccinated VE against hospitalization among immunocompetent adults ≥18 years, by Omicron sublineage period

December 2021-August 2022

SARS-CoV-2 variant period/ mRNA dosage pattern/ days since dose	Median (IQR) days from last dose	Vaccinated case-patients no./total no. (%)	Vaccinated control-patients, no./total no. (%)	Adjusted VE (95% CI)
<b>BA.1/BA.2 period</b>				
<b>2 original monovalent doses</b>				
14-150 days	111 (87-130)	62/771 (8)	79/514 (15)	<b>63 (46-75)</b>
>150 days	290 (241-351)	471/1,180 (40)	404/839 (48)	<b>34 (20-46)</b>
<b>3 original monovalent doses</b>				
7-120 days	80 (55-100)	167/876 (19)	393/828 (47)	<b>79 (74-84)</b>
>120 days	180 (154-208)	265/974 (27)	301/736 (41)	<b>41 (23-55)</b>
<b>BA.4/BA.5 period</b>				
<b>2 original monovalent doses</b>				
14-150 days	102 (77-123)	3/189 (2)	13/168 (8)	<b>83 (35-96)</b>
>150 days	430 (329-471)	128/314 (41)	168/323 (52)	<b>37 (12-55)</b>
<b>3 original monovalent doses</b>				
7-120 days	74 (33-110)	13/199 (7)	24/179 (13)	<b>60 (12-81)</b>
>120 days	237 (204-269)	219/405 (54)	208/363 (57)	<b>29 (3-48)</b>

VE of an additional original monovalent dose provided increased protection during Omicron with similar waning patterns for 2 and 3 doses.


# VISION: Original monovalent vs. unvaccinated VE against hospitalization among immunocompetent adults ≥65 years

June-August 2022

SARS-CoV-2 variant period/ mRNA dosage pattern/ days since dose	Total	SARS-CoV-2- test-positive, N (%)	Median interval since last dose, days (IQR)	Adjusted VE (95% CI)
<b>BA.4/BA.5 period</b>				
Unvaccinated (ref)	2,971	743 (25)	--	
<b>2 original monovalent doses</b>				
14-149 days	77	14 (18)	80 (57-105)	--
≥150 days	2828	556 (20)	473 (422-503)	<b>31 (21-39)</b>
<b>3 original monovalent doses</b>				
7-119 days	289	26 (9)	72 (42-98)	<b>73 (55-84)</b>
≥120 days	4838	913 (19)	240 (211-266)	<b>38 (30-46)</b>
<b>4 original monovalent doses</b>				
7-59 days	765	89 (12)	38 (23-50)	<b>66 (53-75)</b>
≥60 days	1549	210 (14)	88 (75-105)	<b>57 (44-66)</b>

VE of an additional original monovalent dose provided increased protection during Omicron with similar waning patterns for 2 and 3 doses.


## IVY: Bivalent vs. unvaccinated VE against COVID-19-associated *severe in-hospital outcomes* among adults aged $\geq 18$ years without immunocompromising conditions



VE of COVID-19 vaccines was most durable against the most severe outcomes.

# Medicare: VE of *bivalent* vs. original monovalent COVID-19 vaccine against *COVID-19-related thromboembolic events* among immunocompetent Medicare beneficiaries aged ≥65 years

September 2022 – March 2023



\*A COVID-19–related thromboembolic event was defined as presence of an *International Classification of Disease, 10<sup>th</sup> Edition* (ICD-10) or common procedural terminology (CPT) code indicating ischemic stroke, venous thromboembolism, or myocardial infarction from 7 days before through 30 days after a medical claim indicating a COVID–19 diagnosis.

VE estimates adjusted for age, sex, race and ethnicity, geographic region, and calendar time. Updated from: Link-Gelles et al., MMWR,

<https://www.cdc.gov/mmwr/volumes/72/wr/mm7221a3.htm>

TE = thromboembolic event

# VISION: VE of 2 vs. 1 bivalent doses against *hospitalization* among immunocompetent adults aged ≥65 years

September 2022 – September 2023

mRNA Dosage Pattern	Total encounters	SARS-CoV-2- test-positive, N (%)	Median interval since last dose, days (IQR)	Adjusted VE (95% CI)
<b>Hospitalization, 2 vs. 1 bivalent doses (at least 4 months apart)</b>				
1 bivalent dose (ref)	21,825	1,894 (8)	--	Ref
2 bivalent doses	599	50 (8)	50 (26-76)	20 (-9 to 41)


Vaccine Effectiveness (%)

VE of an additional bivalent dose at least 4 months after the original bivalent dose appeared to provide some protection, though the confidence interval was wide and crossed 0.



# VISION: VE of 2023-2024 COVID-19 vaccine against ED/UC encounters among immunocompetent adults aged ≥18 years, by age group

September 2023 – August 2024




<https://www.cdc.gov/mmwr/volumes/73/wr/mm7308a5.htm> (Results updated with additional data since publication.) VE was calculated as  $(1 - \text{odds ratio}) \times 100\%$ , estimated using a test-negative case-control design, with the odds ratio adjusted for age, sex, race and ethnicity, geographic region, and calendar time.

# VISION: VE of 2023-2024 COVID-19 vaccine against hospitalization among immunocompetent adults aged ≥18 years, by age group

September 2023 – August 2024


Age group/2023-2024 COVID-19 vaccination status/days since dose	Total encounters	SARS-CoV-2-test-positive, N (%)	Median interval since last dose among vaccinated, days (IQR)	Adjusted VE (95% CI)
<b>≥18 years</b>				
No 2023-2024 COVID-19 dose (ref)	83,596	8,025 (10)	728 (499-911)	Ref
<b>2023-2024 COVID-19 dose, 7-299 days earlier</b>	21,468	1,664 (8)	120 (62-189)	<b>30 (25 to 34)</b>
<b>2023-2024 COVID-19 dose, 7-59 days earlier</b>	5,095	382 (8)	34 (21-47)	<b>50 (44 to 55)</b>
<b>2023-2024 COVID-19 dose, 60-119 days earlier</b>	5,623	431 (8)	88 (74-104)	<b>38 (31 to 44)</b>
<b>2023-2024 COVID-19 dose, 120-179 days earlier</b>	4,754	268 (6)	148 (134-164)	<b>21 (10 to 31)</b>
<b>2023-2024 COVID-19 dose, 180-299 days earlier</b>	5,996	583 (10)	227 (202-257)	<b>-8 (-19 to 3)</b>
<b>18-64 years</b>				
No 2023-2024 COVID-19 dose (ref)	33,335	2,076 (6)	783 (600-934)	Ref
<b>2023-2024 COVID-19 dose, 7-299 days earlier</b>	3,694	201 (5)	116 (61-182)	<b>16 (1 to 28)</b>
<b>2023-2024 COVID-19 dose, 7-59 days earlier</b>	903	57 (6)	33 (21-45)	<b>26 (3 to 44)</b>
<b>2023-2024 COVID-19 dose, 60-119 days earlier</b>	999	47 (5)	89 (75-104)	<b>34 (11 to 51)</b>
<b>2023-2024 COVID-19 dose, 120-179 days earlier</b>	834	26 (3)	148 (135-164)	<b>28 (-8 to 51)*</b>
<b>2023-2024 COVID-19 dose, 180-299 days earlier</b>	958	71 (7)	226 (199-254)	<b>-35 (-76 to -4)*</b>
<b>≥65 years</b>				
No 2023-2024 COVID-19 dose (ref)	50,261	5,949 (12)	692 (460-890)	Ref
<b>2023-2024 COVID-19 dose, 7-299 days earlier</b>	17,774	1,463 (8)	121 (63-190)	<b>31 (27 to 36)</b>
<b>2023-2024 COVID-19 dose, 7-59 days earlier</b>	4,192	325 (8)	34 (21-47)	<b>53 (47 to 59)</b>
<b>2023-2024 COVID-19 dose, 60-119 days earlier</b>	4,624	384 (8)	88 (74-104)	<b>38 (30 to 44)</b>
<b>2023-2024 COVID-19 dose, 120-179 days earlier</b>	3,920	242 (6)	148 (134-164)	<b>19 (7 to 30)</b>
<b>2023-2024 COVID-19 dose, 180-299 days earlier</b>	5,038	512 (10)	228 (202-257)	<b>-4 (-16 to 7)</b>



\*Some estimates are imprecise, which might be due to a relatively small number of persons in each level of vaccination or case status. This imprecision indicates that the actual VE could be substantially different from the point estimate shown, and estimates should therefore be interpreted with caution. Additional data accrual could increase precision and allow more precise interpretation. <https://www.cdc.gov/mmwr/volumes/73/lwr/mm7308a5.htm> (Results updated with additional data since publication.) VE was calculated as  $(1 - \text{odds ratio}) \times 100\%$ , estimated using a test-negative case-control design, adjusted for age, sex, race and ethnicity, geographic region, and calendar time.

# VISION: VE of 2023-2024 COVID-19 vaccine against *hospitalization and critical illness* among adults aged $\geq 18$ years


September 2023 – August 2024



Additional methods, including definition of immunocompromised available: <https://www.cdc.gov/mmwr/volumes/73/wr/mm7308a5.htm> (Results updated with additional data since publication.) VE was calculated as  $(1 - \text{odds ratio}) \times 100\%$ , estimated using a test-negative case-control design, with odds ratios adjusted for age, sex, race and ethnicity, geographic region, and calendar time.

\* Some estimates are imprecise, which might be due to a relatively small number of persons in each level of vaccination or case status. This imprecision indicates that the actual VE could be substantially different from the point estimate shown, and estimates should therefore be interpreted with caution. Additional data accrual could increase precision and allow more precise interpretation.

## Effectiveness of 2023-2024 COVID-19 vaccines against critical outcomes in immunocompetent adults ≥65 years, Medicare and VISION data



\*A COVID-19–related thromboembolic event was defined as presence of an *International Classification of Disease, 10<sup>th</sup> Edition* (ICD-10) or common procedural terminology (CPT) code indicating ischemic stroke, venous thromboembolism, or myocardial infarction from 7 days before through 30 days after a medical claim indicating a COVID-19 diagnosis.

# Conclusions: COVID-19 VE in adults $\geq 65$ years

- VE findings should be interpreted as the added benefit provided by COVID-19 vaccination in a population with a high prevalence of vaccine- and infection-induced immunity at the start of the 2023-2024 respiratory virus season.
- 2023-2024 COVID-19 vaccination provided increased protection against COVID-19-associated ED/UC visits and hospitalizations compared to no 2023-2024 vaccine dose.
  - Protection waned to 0 against COVID-19-associated ED/UC visits and hospitalization by ~4-6 months.
- Waning patterns of 2023-2024 COVID-19 vaccines appeared similar to previous COVID-19 vaccine formulations; most durable protection appeared to be for critical illness
  - VE against critical illness remained above 40% at 5 months after vaccination among those  $\geq 65$  years
- As with previous COVID-19 vaccine formulations, effectiveness was similar across age groups
- Data from prior seasons show that an additional dose of the same formula appeared to provide additional protection.

# Acknowledgements

## CDC

Amadea Britton  
Allison Ciesla  
Fatimah Dawood  
Jennifer DeCuir  
Monica Dickerson  
Katherine E. Fleming-Dutra  
Sascha Ellington  
Shikha Garg  
Amber Kautz  
Nathaniel M. Lewis  
Kevin Ma  
Josephine Mak  
Joe Miller  
Morgan Najdowski  
Lakshmi Panagiotakopoulos  
Amanda B. Payne  
Zach Smith  
Diya Surie  
Caitlin Ray  
Lauren Roper  
Ryan Wiegand

## VISION Collaborators

### **Westat**

Sarah Ball  
Angela Cheung  
Margaret Dunne  
Patrick Mitchell  
Sarah Reese  
Elizabeth Rowley  
Janet Watts  
Zack Weber

### **Intermountain Health**

Kristin Dascomb

### **Kaiser Permanente Center for Health Research**

Stephanie A. Irving

### **Kaiser Permanente Northern California**

Nicola P. Klein

### **Regenstrief**

Shaun J. Grannis

### **University of Colorado**

Toan C. Ong

### **HealthPartners**

Malini B. DeSilva

### **Columbia University**

Karthik Natarajan

## IVY Collaborators

Cristie Columbus  
Laurence W. Busse  
Steven Y. Chang  
Abhijit Duggal  
Matthew C. Exline  
Manjusha Gaglani  
Kevin W. Gibbs  
Adit A. Ginde  
David N. Hager  
Estelle S. Harris  
Cassandra Johnson  
Nicholas J. Johnson  
Akram Khan  
Jennie H. Kwon  
Adam S. Lauring  
Christopher Mallow  
Emily Martin  
Amira Mohamed

Nicholas M. Mohr  
Jarrod M. Mosier  
Ithan D. Peltan  
Matthew Prekker  
Basmah Safdar  
Wesley H. Self  
Nathan I. Shapiro  
Jay S. Steingrub  
Ivana A. Vaughn  
Jennifer G. Wilson  
Yuwei Zhu

## CMS Collaborators

### **CMS**

Alia Bayatti

### **Acumen**

Heng-Ming Sung  
Ivy Zhang  
Carla Gomez Victor  
Yenlin Lai  
Bradley Lufkin  
Yoganand Chillarige

**+ many more!**