

**Table C2\_1**  
**Child Current Asthma Prevalence Rate (Percent)**  
**by Sex and State or Territory: BRFSS 2006**

State	Male Sample Size	Male Prevalence	Male Standard Error	Male 95% CI *	Female Sample Size	Female Prevalence	Female Standard Error	Female 95% CI *
Total **	31,968	10.2	0.37	(9.5 - 11.0)	30,280	7.7	0.34	(7.1 - 8.4)
AK	408	7.7	1.72	(4.9 - 11.8)	384	6.1	1.62	(3.6 - 10.1)
AZ	730	9.7	1.93	(6.5 - 14.2)	670	10.6	2.67	(6.4 - 17.1)
CA	876	10.1	1.23	(7.9 - 12.8)	859	6.5	1.00	(4.8 - 8.7)
DC	426	14.9	2.24	(11.1 - 19.9)	461	13.7	2.33	(9.8 - 19.0)
GA	1,237	11.4	1.28	(9.1 - 14.1)	1,115	8.7	1.15	(6.7 - 11.3)
HI	1,122	15.5	1.39	(12.9 - 18.4)	1,043	9.3	1.16	(7.3 - 11.8)
ID	994	7.3	0.93	(5.7 - 9.4)	870	5.2	0.89	(3.7 - 7.3)
IN	1,014	9.8	1.14	(7.8 - 12.3)	1,023	6.9	0.94	(5.3 - 9.0)
IA	835	7.6	1.09	(5.7 - 10.0)	812	5.5	0.94	(3.9 - 7.7)
KS	1,316	10.1	1.01	(8.3 - 12.3)	1,202	8.2	0.86	(6.7 - 10.0)
KY	768	11.2	1.53	(8.5 - 14.6)	758	7.3	1.28	(5.2 - 10.3)
ME	582	10.4	1.47	(7.8 - 13.6)	540	6.9	1.15	(5.0 - 9.5)
MD	1,372	10.7	1.12	(8.7 - 13.2)	1,277	7.5	0.97	(5.8 - 9.6)
MI	912	8.9	1.19	(6.8 - 11.5)	856	8.1	1.24	(6.0 - 10.9)
MN	701	9.1	1.23	(7.0 - 11.8)	627	4.8	0.83	(3.4 - 6.7)
MS	986	11.9	1.21	(9.7 - 14.5)	954	8.5	1.12	(6.6 - 11.0)
MO	797	8.8	1.34	(6.5 - 11.8)	767	10.9	1.88	(7.7 - 15.1)
MT	830	8.1	1.27	(5.9 - 10.9)	830	5.1	0.89	(3.7 - 7.2)
NE	1,268	7.1	0.96	(5.4 - 9.3)	1,134	6.2	0.88	(4.7 - 8.2)
NV	605	8.5	1.58	(5.9 - 12.1)	530	5.6	1.75	(3.0 - 10.2)
NH	907	9.9	1.23	(7.7 - 12.6)	895	8.4	1.13	(6.4 - 10.9)
NJ	1,965	11.6	0.92	(10.0 - 13.6)	1,786	9.2	0.99	(7.4 - 11.3)
NY	847	10.9	1.41	(8.4 - 14.0)	788	10.6	1.40	(8.2 - 13.7)
OH	824	9.1	1.77	(6.1 - 13.2)	809	7.2	1.69	(4.5 - 11.3)
OK	1,100	9.8	1.02	(8.0 - 12.0)	1,004	7.4	0.89	(5.9 - 9.3)
PA	1,779	9.5	1.22	(7.4 - 12.2)	1,774	7.3	1.21	(5.2 - 10.0)

Notes:  
\* CI denotes confidence interval.  
\*\* Total includes data from 33 states plus the District of Columbia and excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

**Table C2\_1**  
**Child Current Asthma Prevalence Rate (Percent)**  
**by Sex and State or Territory: BRFSS 2006**

State	Male Sample Size	Male Prevalence	Male Standard Error	Male 95% CI *		Female Sample Size	Female Prevalence	Female Standard Error	Female 95% CI *
RI	659	13.3	1.53	(10.6 - 16.6)		640	8.0	1.19	(6.0 - 10.7)
TX	1,132	11.4	1.57	(8.6 - 14.8)		1,056	7.4	1.33	(5.2 - 10.5)
UT	1,171	7.5	0.96	(5.9 - 9.6)		1,106	5.5	0.82	(4.1 - 7.4)
VT	954	10.4	1.13	(8.4 - 12.8)		945	8.2	1.07	(6.3 - 10.6)
VA	848	11.2	1.57	(8.5 - 14.7)		789	9.5	1.47	(7.0 - 12.8)
WV	514	9.6	1.49	(7.0 - 12.9)		507	7.1	1.27	(5.0 - 10.1)
WI	718	9.3	1.32	(7.0 - 12.2)		737	5.6	0.98	(4.0 - 7.9)
WY	771	10.5	1.28	(8.3 - 13.3)		732	6.6	1.09	(4.8 - 9.1)
Territories									
PR	756	20.4	1.67	(17.3 - 23.9)		720	11.9	1.42	(9.4 - 15.0)

Notes:  
\* CI denotes confidence interval.  
\*\* Total includes data from 33 states plus the District of Columbia and excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.