

Table L2_1
Child Lifetime Asthma Prevalence Rate (Percent)
by Sex and State or Territory: BRFSS 2006

State	Male Sample Size	Male Prevalence	Male Standard Error	Male 95% CI *		Female Sample Size	Female Prevalence	Female Standard Error	Female 95% CI *
Total **	32,117	15.0	0.44	(14.1 - 15.9)		30,385	10.7	0.38	(10.0 - 11.5)
AK	410	12.1	2.05	(8.6 - 16.7)		386	7.3	1.69	(4.6 - 11.4)
AZ	733	14.3	2.29	(10.3 - 19.4)		671	14.4	2.85	(9.7 - 21.0)
CA	880	15.2	1.48	(12.6 - 18.4)		860	9.7	1.17	(7.6 - 12.3)
DC	431	21.7	2.59	(17.1 - 27.2)		462	16.7	2.46	(12.4 - 22.1)
GA	1,242	17.0	1.43	(14.4 - 20.0)		1,120	11.6	1.24	(9.4 - 14.2)
HI	1,136	24.3	1.66	(21.2 - 27.7)		1,049	16.8	1.61	(13.8 - 20.2)
ID	1,001	10.8	1.12	(8.8 - 13.2)		875	8.1	1.08	(6.2 - 10.5)
IN	1,017	14.0	1.29	(11.6 - 16.7)		1,030	10.6	1.15	(8.5 - 13.1)
IA	839	10.9	1.25	(8.7 - 13.6)		814	7.2	1.06	(5.4 - 9.6)
KS	1,318	13.3	1.12	(11.3 - 15.7)		1,206	10.4	0.97	(8.7 - 12.5)
KY	770	16.2	1.75	(13.1 - 20.0)		762	10.1	1.45	(7.6 - 13.3)
ME	588	15.8	1.71	(12.8 - 19.5)		542	8.9	1.29	(6.7 - 11.8)
MD	1,380	15.2	1.30	(12.9 - 18.0)		1,282	11.0	1.19	(8.9 - 13.5)
MI	913	14.4	1.58	(11.6 - 17.8)		858	10.9	1.35	(8.5 - 13.8)
MN	702	12.3	1.41	(9.8 - 15.4)		628	6.6	0.96	(4.9 - 8.7)
MS	993	17.3	1.47	(14.6 - 20.4)		958	11.1	1.23	(8.9 - 13.8)
MO	800	13.4	1.69	(10.4 - 17.1)		767	14.1	2.01	(10.6 - 18.5)
MT	834	12.1	1.47	(9.5 - 15.3)		831	7.6	1.05	(5.8 - 9.9)
NE	1,273	11.3	1.16	(9.2 - 13.7)		1,141	9.0	1.08	(7.1 - 11.3)
NV	607	13.7	1.93	(10.3 - 17.9)		530	9.7	2.25	(6.1 - 15.1)
NH	911	15.3	1.47	(12.6 - 18.4)		898	11.0	1.24	(8.8 - 13.7)
NJ	1,972	17.5	1.10	(15.5 - 19.8)		1,792	13.2	1.11	(11.2 - 15.5)
NY	852	16.6	1.67	(13.5 - 20.1)		790	13.4	1.55	(10.7 - 16.8)
OH	828	10.7	1.85	(7.6 - 14.9)		813	8.8	1.78	(5.8 - 12.9)
OK	1,105	15.1	1.26	(12.8 - 17.8)		1,005	10.9	1.06	(9.0 - 13.2)
PA	1,783	14.3	1.53	(11.5 - 17.5)		1,783	9.8	1.34	(7.4 - 12.7)

Notes:
* CI denotes confidence interval.
** Total includes data from 33 states plus the District of Columbia and excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Table L2_1
Child Lifetime Asthma Prevalence Rate (Percent)
by Sex and State or Territory: BRFSS 2006

State	Male Sample Size	Male Prevalence	Male Standard Error	Male 95% CI *		Female Sample Size	Female Prevalence	Female Standard Error	Female 95% CI *
RI	667	21.0	1.83	(17.6 - 24.8)		641	12.0	1.61	(9.1 - 15.5)
TX	1,134	16.6	1.81	(13.3 - 20.4)		1,059	10.3	1.49	(7.8 - 13.7)
UT	1,174	11.3	1.13	(9.3 - 13.7)		1,111	6.9	0.88	(5.3 - 8.8)
VT	958	14.2	1.26	(11.9 - 16.9)		947	11.5	1.21	(9.3 - 14.1)
VA	853	15.6	1.75	(12.5 - 19.4)		793	13.1	1.60	(10.2 - 16.5)
WV	518	12.7	1.67	(9.7 - 16.3)		509	9.9	1.49	(7.3 - 13.2)
WI	719	12.4	1.54	(9.6 - 15.7)		739	7.9	1.23	(5.8 - 10.7)
WY	776	13.2	1.39	(10.7 - 16.2)		733	8.5	1.19	(6.5 - 11.2)
Territories									
PR	758	34.4	1.96	(30.7 - 38.4)		720	24.1	1.80	(20.7 - 27.8)

Notes:
* CI denotes confidence interval.
** Total includes data from 33 states plus the District of Columbia and excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.