

Table C2_1
Child Current Asthma Prevalence
by Sex and State or Territory: BRFSS 2017

State	Male Unweighted Number	Male Prevalence	Male Standard Error	Male 95% CI *		Female Unweighted Number	Female Prevalence	Female Standard Error	Female 95% CI *
Total **	27,030	8.8	0.35	(8.2–9.5)		24,712	7.1	0.34	(6.4–7.8)
CA	913	6.8	1.00	(5.1–9.1)		827	5.6	0.96	(4.0–7.8)
CT	977	14.4	1.56	(11.6–17.8)		920	11.5	1.83	(8.4–15.6)
DC	406	12.8	1.96	(9.5–17.2)		365	10.5	1.86	(7.3–14.7)
FL	2,193	8.5	1.19	(6.5–11.2)		1,980	6.2	1.00	(4.5–8.5)
GA	580	10.4	1.56	(7.7–13.9)		504	7.8	1.53	(5.3–11.4)
HI	974	11.0	1.43	(8.5–14.1)		858	8.9	1.38	(6.5–12.0)
IL	586	5.6	1.03	(3.9–8.0)		522	5.8	1.17	(3.9–8.6)
IN	1,419	8.6	0.84	(7.1–10.4)		1,319	5.0	0.64	(3.9–6.4)
KS	2,435	10.0	0.92	(8.4–12.0)		2,278	8.6	0.86	(7.0–10.5)
ME	924	9.3	1.37	(6.9–12.3)		834	6.7	1.13	(4.8–9.3)
MA	276	17.3	3.94	(10.9–26.4)		232	15.6	4.79	(8.3–27.4)
MI	1,179	9.6	1.03	(7.8–11.8)		1,131	7.8	0.98	(6.1–10.0)
MN	1,768	5.1	0.65	(4.0–6.5)		1,612	3.9	0.58	(2.9–5.2)
MS	482	11.2	1.85	(8.0–15.3)		462	11.6	2.38	(7.6–17.1)
MO	733	10.7	1.50	(8.1–14.0)		657	8.8	1.57	(6.2–12.4)
MT	647	9.4	1.79	(6.5–13.6)		612	7.8	1.68	(5.1–11.8)
NE	1,829	7.3	1.07	(5.5–9.7)		1,643	5.0	0.80	(3.7–6.8)
NV	373	7.3	1.97	(4.3–12.2)		365	8.1	2.35	(4.6–14.1)
NJ	1,173	9.5	1.25	(7.3–12.3)		1,091	7.0	1.10	(5.1–9.4)
NM	689	9.8	2.03	(6.5–14.6)		669	9.1	1.61	(6.4–12.8)
NY	604	8.4	1.55	(5.9–12.0)		637	9.3	1.36	(6.9–12.3)
OH	1,145	9.7	1.23	(7.5–12.4)		1,047	7.8	1.14	(5.8–10.3)
OR	545	8.2	1.35	(5.9–11.3)		488	3.5	0.83	(2.1–5.5)
PA	684	12.6	1.59	(9.8–16.1)		632	5.0	1.03	(3.3–7.4)
RI	497	12.4	1.98	(9.0–16.8)		461	8.2	1.75	(5.4–12.4)
UT	1,814	7.4	0.83	(5.9–9.2)		1,542	5.0	0.69	(3.8–6.6)
VT	640	8.5	1.30	(6.3–11.5)		508	6.0	1.10	(4.2–8.6)
WI	545	7.1	1.69	(4.4–11.2)		516	7.9	1.65	(5.2–11.8)
Territory									
PR	514	17.9	2.59	(13.3–23.5)		484	13.4	2.62	(9.1–19.4)

Notes:

* CI denotes confidence interval.

** Total includes data from 27 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.