

Table L1
Child Lifetime Asthma Prevalence
and Weighted Number by State or Territory: BRFSS 2017

State	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)	 	Weighted Number	95% CI * (Number)
Total **	52,683	12.1	0.29	(11.5–12.7)		5,281,110	(5,025,515–5,536,704)
CA	1,753	10.8	0.90	(9.2–12.7)		918,537	(767,527–1,069,548)
CT	1,954	18.9	1.38	(16.4–21.8)		134,457	(113,649–155,265)
DC	795	16.6	1.58	(13.7–19.9)		20,367	(16,485–24,249)
FL	4,285	11.8	0.99	(10.0–13.9)		470,243	(391,025–549,461)
GA	1,116	13.5	1.24	(11.2–16.1)		317,837	(259,380–376,295)
HI	1,861	16.3	1.14	(14.1–18.6)		48,832	(41,868–55,795)
IL	1,117	11.0	1.12	(9.0–13.4)		306,617	(244,290–368,945)
IN	2,759	10.5	0.64	(9.4–11.8)		159,232	(140,456–178,009)
KS	4,775	13.4	0.69	(12.1–14.8)		90,674	(81,122–100,225)
ME	1,766	11.2	1.02	(9.3–13.3)		27,680	(22,755–32,605)
MA	535	21.0	3.18	(15.4–27.9)		263,720	(176,611–350,830)
MI	2,340	11.9	0.79	(10.5–13.6)		250,402	(217,483–283,321)
MN	3,454	6.5	0.50	(5.6–7.6)		81,656	(69,354–93,959)
MS	955	15.3	1.59	(12.4–18.7)		105,418	(83,276–127,560)
MO	1,416	13.6	1.21	(11.4–16.1)		180,737	(148,435–213,039)
MT	1,285	12.0	1.51	(9.3–15.3)		26,582	(19,770–33,395)
NE	3,510	9.4	0.78	(8.0–11.1)		43,507	(36,334–50,680)
NV	750	10.9	1.69	(8.0–14.6)		70,091	(48,371–91,812)
NJ	2,322	13.0	1.00	(11.2–15.1)		239,678	(203,120–276,235)
NM	1,385	14.3	1.52	(11.5–17.5)		67,250	(52,646–81,854)
NY	1,260	13.0	1.19	(10.8–15.5)		515,593	(421,391–609,796)
OH	2,232	11.8	0.94	(10.1–13.8)		291,096	(244,673–337,518)
OR	1,050	9.2	0.98	(7.5–11.3)		76,771	(61,065–92,476)
PA	1,341	12.9	1.12	(10.9–15.3)		328,894	(273,073–384,714)
RI	973	15.5	1.54	(12.7–18.8)		30,926	(24,931–36,922)
UT	3,441	8.7	0.61	(7.6–10.0)		78,635	(67,756–89,513)
VT	1,175	11.6	1.12	(9.6–14.0)		13,059	(10,636–15,482)
WI	1,078	9.9	1.25	(7.7–12.6)		122,617	(91,738–153,497)

Notes:

* CI denotes confidence interval.

** Total includes data from 27 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Table L1
Child Lifetime Asthma Prevalence
and Weighted Number by State or Territory: BRFSS 2017

State	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)	 	Weighted Number	95% CI * (Number)
Territory							
PR	1,011	24.4	2.06	(20.6–28.6)		175,785	(144,153–207,417)

Notes:

* CI denotes confidence interval.

** Total includes data from 27 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.