

**Table L2_1
Child Lifetime Asthma Prevalence
by Sex and State or Territory: BRFSS 2017**

State	Male Unweighted Number	Male Prevalence	Male Standard Error	Male 95% CI *		Female Unweighted Number	Female Prevalence	Female Standard Error	Female 95% CI *
Total **	27,128	13.9	0.43	(13.1–14.7)		24,763	10.3	0.40	(9.6–11.1)
CA	916	12.4	1.33	(10.0–15.3)		829	9.3	1.22	(7.2–12.0)
CT	980	21.5	1.81	(18.2–25.2)		922	16.8	2.19	(13.0–21.6)
DC	408	16.9	2.21	(13.0–21.7)		371	16.7	2.30	(12.7–21.7)
FL	2,201	13.6	1.47	(11.0–16.7)		1,983	10.0	1.36	(7.7–13.1)
GA	581	15.8	1.83	(12.5–19.7)		505	11.0	1.69	(8.1–14.8)
HI	979	18.2	1.63	(15.2–21.6)		860	14.4	1.63	(11.5–18.0)
IL	586	12.0	1.67	(9.1–15.7)		523	10.0	1.50	(7.4–13.4)
IN	1,420	13.2	0.99	(11.4–15.3)		1,319	7.7	0.78	(6.3–9.4)
KS	2,449	15.3	1.04	(13.4–17.4)		2,284	11.2	0.91	(9.6–13.1)
ME	926	12.8	1.53	(10.1–16.1)		834	9.5	1.31	(7.3–12.4)
MA	276	25.0	4.29	(17.5–34.3)		232	18.9	4.88	(11.1–30.3)
MI	1,186	13.6	1.17	(11.4–16.0)		1,133	10.2	1.08	(8.2–12.5)
MN	1,772	7.9	0.77	(6.5–9.6)		1,613	5.0	0.65	(3.8–6.4)
MS	484	15.8	2.09	(12.1–20.3)		463	14.8	2.46	(10.6–20.3)
MO	735	15.8	1.74	(12.6–19.5)		661	11.6	1.71	(8.7–15.4)
MT	650	14.2	2.43	(10.1–19.7)		612	10.0	1.86	(6.9–14.3)
NE	1,840	11.1	1.22	(8.9–13.7)		1,643	7.7	0.98	(6.0–9.9)
NV	374	11.0	2.20	(7.4–16.1)		366	10.7	2.63	(6.5–17.1)
NJ	1,178	13.9	1.40	(11.4–16.9)		1,094	12.2	1.47	(9.6–15.4)
NM	692	15.9	2.44	(11.7–21.3)		671	12.7	1.85	(9.5–16.8)
NY	605	13.5	1.86	(10.2–17.6)		638	12.3	1.55	(9.6–15.7)
OH	1,147	14.0	1.44	(11.4–17.0)		1,051	9.9	1.24	(7.7–12.6)
OR	548	13.7	1.68	(10.8–17.4)		491	4.9	0.98	(3.3–7.2)
PA	688	18.0	1.78	(14.8–21.7)		634	7.8	1.31	(5.6–10.8)
RI	501	19.4	2.26	(15.4–24.2)		461	11.7	2.01	(8.3–16.2)
UT	1,819	11.1	0.95	(9.4–13.1)		1,544	6.6	0.78	(5.3–8.3)
VT	641	13.9	1.67	(11.0–17.5)		510	9.3	1.50	(6.7–12.6)
WI	546	8.8	1.76	(5.9–12.9)		516	11.3	1.92	(8.0–15.6)
Territory									
PR	517	27.7	2.87	(22.4–33.7)		485	21.6	3.00	(16.3–28.0)

Notes:

* CI denotes confidence interval.

** Total includes data from 27 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.