

Table L3
Child Lifetime Asthma Prevalence
and Weighted Number by Age and State or Territory: BRFSS 2017

State	Age Group	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)	 	Weighted Number	95% CI * Weighted Number
Total **	0-4	10,526	5.7	0.42	(4.9–6.6)		564,409	(481,538–647,281)
Total **	5-9	11,209	13.3	0.67	(12.1–14.7)		1,499,801	(1,342,322–1,657,279)
Total **	10-14	13,422	15.2	0.64	(14.0–16.5)		1,559,706	(1,422,634–1,696,779)
Total **	15-17	11,260	15.9	0.74	(14.5–17.4)		1,138,730	(1,027,543–1,249,917)
CA	0-4	432	3.1	0.88	(1.8–5.4)		56,088	(24,963–87,214)
CA	5-9	396	12.8	1.95	(9.4–17.1)		292,824	(201,306–384,342)
CA	10-14	472	14.0	1.88	(10.7–18.1)		300,794	(217,656–383,933)
CA	15-17	319	14.0	2.36	(9.9–19.3)		211,155	(136,753–285,556)
CT	0-4	328	9.8	2.19	(6.3–15.0)		15,574	(8,441–22,707)
CT	5-9	380	23.9	3.27	(18.1–30.9)		45,536	(31,020–60,053)
CT	10-14	513	20.1	2.90	(15.0–26.4)		31,803	(21,314–42,291)
CT	15-17	483	21.2	2.65	(16.5–26.9)		24,874	(18,335–31,413)
DC	0-4	178	9.0	2.97	(4.6–16.7)		2,504	(790–4,219)
DC	5-9	186	18.4	3.07	(13.2–25.2)		7,167	(4,714–9,620)
DC	10-14	185	19.6	3.21	(14.0–26.6)		5,077	(3,345–6,810)
DC	15-17	143	28.3	4.93	(19.7–38.9)		4,863	(2,843–6,884)
FL	0-4	909	9.1	2.15	(5.6–14.3)		70,467	(35,856–105,077)
FL	5-9	868	13.6	2.26	(9.7–18.7)		137,234	(89,395–185,074)
FL	10-14	977	14.7	2.35	(10.6–19.9)		118,165	(78,726–157,603)
FL	15-17	760	14.7	2.39	(10.6–20.1)		106,308	(70,510–142,106)
GA	0-4	192	8.2	2.51	(4.4–14.6)		38,658	(14,631–62,684)
GA	5-9	225	15.9	2.92	(11.0–22.5)		92,167	(56,204–128,129)
GA	10-14	260	18.3	2.91	(13.3–24.7)		87,853	(57,811–117,895)
GA	15-17	216	15.1	2.86	(10.3–21.6)		56,232	(34,197–78,267)
HI	0-4	387	11.3	2.36	(7.4–16.8)		10,091	(5,643–14,539)
HI	5-9	431	18.1	2.24	(14.1–22.9)		15,205	(11,251–19,158)
HI	10-14	464	21.0	2.51	(16.5–26.3)		11,145	(8,198–14,093)
HI	15-17	362	20.5	2.71	(15.7–26.3)		8,948	(6,472–11,423)
IL	0-4	209	4.8	1.72	(2.3–9.5)		30,690	(8,533–52,848)
IL	5-9	245	11.0	2.26	(7.3–16.3)		85,753	(50,006–121,500)
IL	10-14	295	15.6	2.72	(11.0–21.7)		110,375	(69,467–151,282)

Notes:

* CI denotes confidence interval.

** Total includes data from 27 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

**Table L3
Child Lifetime Asthma Prevalence
and Weighted Number by Age and State or Territory: BRFSS 2017**

State	Age Group	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)	 	Weighted Number	95% CI * Weighted Number
IL	15-17	272	13.1	2.22	(9.3–18.1)		56,520	(37,259–75,781)
IN	0-4	539	4.0	0.91	(2.5–6.2)		16,546	(8,941–24,151)
IN	5-9	611	9.8	1.33	(7.5–12.8)		32,467	(23,465–41,470)
IN	10-14	696	14.6	1.51	(11.9–17.8)		51,451	(40,491–62,411)
IN	15-17	602	15.8	1.66	(12.8–19.3)		41,230	(32,181–50,279)
KS	0-4	1,056	6.0	1.17	(4.1–8.8)		9,813	(5,892–13,733)
KS	5-9	1,043	13.2	1.40	(10.7–16.2)		25,439	(19,745–31,133)
KS	10-14	1,234	16.7	1.54	(13.9–19.9)		25,559	(20,457–30,661)
KS	15-17	1,032	20.2	1.74	(17.0–23.9)		22,747	(18,444–27,049)
ME	0-4	325	2.1	0.83	(1.0–4.5)		1,357	(317–2,397)
ME	5-9	377	14.6	2.65	(10.2–20.6)		9,383	(5,806–12,961)
ME	10-14	526	14.0	2.00	(10.5–18.4)		8,318	(5,850–10,787)
ME	15-17	421	15.6	2.31	(11.6–20.7)		6,974	(4,838–9,111)
MA	0-4	98	4.4				10,211	
MA	5-9	90	36.3	7.81	(22.7–52.5)		133,043	(58,893–207,192)
MA	10-14	135	9.5	2.99	(5.1–17.2)		24,948	(9,571–40,325)
MA	15-17	116	23.1	7.08	(12.1–39.6)		47,348	(14,015–80,680)
MI	0-4	453	6.7	1.29	(4.5–9.7)		32,463	(19,889–45,037)
MI	5-9	496	11.7	1.65	(8.8–15.3)		65,698	(46,616–84,779)
MI	10-14	599	15.1	1.81	(11.9–19.0)		79,385	(59,334–99,436)
MI	15-17	597	15.6	1.73	(12.5–19.3)		54,141	(41,817–66,466)
MN	0-4	649	2.2	0.68	(1.2–4.0)		7,741	(3,028–12,453)
MN	5-9	713	4.6	0.87	(3.1–6.6)		16,291	(10,142–22,441)
MN	10-14	849	10.0	1.21	(7.9–12.7)		20,531	(15,511–25,552)
MN	15-17	708	12.6	1.67	(9.7–16.3)		27,108	(19,536–34,680)
MS	0-4	175	9.7	3.43	(4.7–18.8)		16,284	(4,167–28,400)
MS	5-9	183	16.7	3.67	(10.7–25.1)		28,012	(15,038–40,986)
MS	10-14	279	19.2	3.23	(13.6–26.3)		32,754	(20,931–44,577)
MS	15-17	218	19.5	3.15	(14.0–26.4)		21,197	(13,968–28,426)
MO	0-4	269	6.3	1.92	(3.5–11.3)		19,242	(7,478–31,006)
MO	5-9	304	14.0	2.66	(9.5–20.0)		48,365	(28,962–67,768)

Notes:

* CI denotes confidence interval.

** Total includes data from 27 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Table L3
Child Lifetime Asthma Prevalence
and Weighted Number by Age and State or Territory: BRFSS 2017

State	Age Group	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)	 	Weighted Number	95% CI * Weighted Number
MO	10-14	363	17.8	2.58	(13.3–23.4)		59,179	(40,814–77,545)
MO	15-17	330	21.0	3.15	(15.5–27.9)		44,319	(29,774–58,864)
MT	0-4	287	1.7				731	
MT	5-9	277	12.2	3.68	(6.6–21.4)		7,516	(2,672–12,360)
MT	10-14	328	15.6	2.97	(10.6–22.3)		8,271	(4,958–11,583)
MT	15-17	262	18.5	3.70	(12.4–26.9)		7,098	(4,010–10,186)
NE	0-4	752	4.9	1.42	(2.8–8.5)		5,925	(2,474–9,375)
NE	5-9	746	7.4	1.40	(5.1–10.7)		9,449	(5,852–13,046)
NE	10-14	980	12.4	1.56	(9.7–15.8)		13,260	(9,836–16,684)
NE	15-17	767	15.6	2.35	(11.5–20.8)		11,419	(7,706–15,131)
NV	0-4	155	5.8	2.67	(2.3–13.8)		10,575	(824–20,326)
NV	5-9	153	15.9	4.15	(9.3–25.8)		30,121	(13,504–46,738)
NV	10-14	215	9.9	2.66	(5.7–16.4)		13,729	(6,242–21,217)
NV	15-17	145	14.7	4.78	(7.6–26.7)		10,953	(3,337–18,569)
NJ	0-4	376	11.8	2.58	(7.6–17.9)		52,996	(28,968–77,025)
NJ	5-9	457	10.9	2.02	(7.5–15.6)		47,766	(29,724–65,808)
NJ	10-14	606	16.2	1.93	(12.7–20.3)		64,876	(48,922–80,831)
NJ	15-17	524	18.9	2.41	(14.6–24.1)		51,654	(37,865–65,443)
NM	0-4	262	3.5	1.84	(1.2–9.5)		3,625	(0–7,452)
NM	5-9	304	18.0	3.89	(11.6–26.9)		23,044	(11,961–34,128)
NM	10-14	350	18.4	3.03	(13.1–25.1)		18,679	(12,048–25,309)
NM	15-17	276	20.7	3.61	(14.5–28.7)		14,768	(9,119–20,416)
NY	0-4	269	6.2	1.94	(3.3–11.3)		59,070	(21,784–96,355)
NY	5-9	271	14.1	2.54	(9.8–19.9)		123,915	(77,864–169,966)
NY	10-14	299	17.9	2.90	(12.9–24.3)		174,614	(113,785–235,443)
NY	15-17	248	15.4	2.80	(10.7–21.7)		100,057	(62,004–138,110)
OH	0-4	381	5.9	1.60	(3.4–9.9)		34,153	(15,359–52,948)
OH	5-9	466	12.5	2.06	(9.0–17.1)		72,370	(47,446–97,294)
OH	10-14	572	14.4	2.03	(10.9–18.8)		84,828	(59,651–110,004)
OH	15-17	505	15.6	2.25	(11.7–20.5)		56,397	(39,122–73,672)
OR	0-4	222	2.7	1.25	(1.1–6.6)		5,906	(584–11,228)

Notes:

* CI denotes confidence interval.

** Total includes data from 27 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Table L3
Child Lifetime Asthma Prevalence
and Weighted Number by Age and State or Territory: BRFSS 2017

State	Age Group	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)	 	Weighted Number	95% CI * Weighted Number
OR	5-9	245	8.0	1.91	(4.9–12.6)		17,803	(9,288–26,318)
OR	10-14	251	13.8	2.31	(9.9–19.0)		25,901	(17,052–34,751)
OR	15-17	223	15.1	2.99	(10.2–22.0)		19,075	(11,481–26,669)
PA	0-4	263	5.9	1.55	(3.5–9.8)		32,684	(15,722–49,645)
PA	5-9	300	11.2	2.16	(7.6–16.2)		70,203	(43,025–97,382)
PA	10-14	332	15.2	2.40	(11.0–20.5)		103,633	(69,309–137,957)
PA	15-17	312	19.9	2.97	(14.7–26.3)		81,098	(55,172–107,023)
RI	0-4	184	4.0	1.70	(1.7–9.0)		2,170	(359–3,981)
RI	5-9	187	17.5	3.61	(11.5–25.7)		9,565	(5,347–13,783)
RI	10-14	271	23.4	3.20	(17.7–30.2)		10,003	(7,027–12,979)
RI	15-17	222	19.2	3.34	(13.5–26.6)		6,003	(3,843–8,163)
UT	0-4	731	3.5	0.85	(2.2–5.6)		6,142	(3,218–9,065)
UT	5-9	772	7.0	1.01	(5.3–9.3)		16,662	(11,841–21,482)
UT	10-14	801	12.6	1.51	(9.9–15.8)		30,061	(22,512–37,609)
UT	15-17	665	13.6	1.79	(10.5–17.5)		17,762	(12,852–22,672)
VT	0-4	227	3.8	1.15	(2.1–6.9)		1,123	(465–1,781)
VT	5-9	261	11.4	2.47	(7.4–17.2)		3,078	(1,750–4,407)
VT	10-14	300	17.1	2.79	(12.3–23.2)		4,799	(3,160–6,438)
VT	15-17	282	14.6	2.37	(10.5–19.9)		2,776	(1,871–3,680)
WI	0-4	218	4.3	2.31	(1.5–11.9)		11,582	(0–24,097)
WI	5-9	222	9.7	2.81	(5.4–16.8)		33,723	(13,628–53,818)
WI	10-14	270	13.7	2.82	(9.0–20.2)		39,713	(22,549–56,877)
WI	15-17	250	12.6	2.83	(8.0–19.3)		25,708	(13,971–37,444)
Territory								
PR	0-4	210	12.1	2.77	(7.6–18.7)		18,586	(9,965–27,207)
PR	5-9	225	31.4	4.75	(22.9–41.4)		49,203	(30,553–67,854)
PR	10-14	253	31.6	4.62	(23.3–41.2)		64,172	(42,755–85,589)
PR	15-17	222	26.1	3.58	(19.7–33.7)		31,066	(22,388–39,743)

Notes:

* CI denotes confidence interval.

** Total includes data from 27 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.