Group B Streptococcus Infections **Caused by Improper Sourcing** and Handling of Fish for Raw Consumption, Singapore, 2015–2016

Man L. Chau,¹ Swaine L. Chen,¹ Min Yap, Sri H.P. Hartantyo, Paul K.T. Chiew, Charlene J. Fernandez, Wai K. Wong, Rockey K. Fong, Wei L. Tan, Brian Z.Y. Tan, Youming Ng, Kyaw T. Aung, Kurosh S. Mehershahi, Christopher Goh, Joanne S.L. Kang, Timothy Barkham, Adeline O.K. Leong, Ramona A. Gutiérrez, Lee C. Ng

Medscape ACTIVITY



JOINTLY ACCREDITED PROVIDER ERPROFESSIONAL CONTINUING EDUCATIO

In support of improving patient care, this activity has been planned and implemented by Medscape, LLC and Emerging Infectious Diseases. Medscape, LLC is jointly accredited by the Accreditation Council for Continuing Medical Education (ACCME), the Accreditation Council for Pharmacy Education (ACPE), and the American Nurses Credentialing Center (ANCC), to provide continuing education for the healthcare team.

Medscape, LLC designates this Journal-based CME activity for a maximum of 1.00 AMA PRA Category 1 Credit(s)TM. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

All other clinicians completing this activity will be issued a certificate of participation. To participate in this journal CME activity: (1) review the learning objectives and author disclosures; (2) study the education content; (3) take the post-test with a 75% minimum passing score and complete the evaluation at http://www.medscape.org/journal/eid; and (4) view/print certificate. For CME questions, see page 2134.

Release date: November 15, 2017; Expiration date: November 15, 2018

Learning Objectives

Upon completion of this activity, participants will be able to:

- Evaluate clinical and epidemiological findings of a human group B Streptococcus outbreak in Singapore based on a microbial study of raw fish and human samples
- Determine the pathogenic potential of fish and human samples after a human group B Streptococcus outbreak in Singapore, based on a microbial study of these samples
- · Assess the public health implications of a human group B Streptococcus outbreak in Singapore, based on a microbial study of raw fish and human samples.

CME Editor

Thomas J. Gryczan, MS, Technical Writer/Editor, Emerging Infectious Diseases. Disclosure: Thomas J. Gryczan, MS, has disclosed no relevant financial relationships.

CME Author

Laurie Barclay, MD, freelance writer and reviewer, Medscape, LLC. Disclosure: Laurie Barclay, MD, has disclosed the following relevant financial relationships: owns stock, stock options, or bonds from Alnylam; Biogen; Pfizer.

Authors

Disclosures: Man Ling Chau, BSc; Swaine L. Chen, MD, PhD; Min Yap, BSc; Sri Harminda Pahm Hartantyo, PhD; Paul King Tiong Chiew, DVM, MVPM; Charlene Judith Fernandez, BS, DVM, MPH, CPH; Wai Kwan Wong, BSc; Rockey Kwonghung Fong, BVSc; Wei Ling Tan, BSc; Brian Zi Yan Tan, PhD; Youming Ng, BSc; Kyaw Thu Aung, MBBS, MSc, PGDip; Kurosh S. Mehershahi, PhD; Christopher Goh, MSc; Joanne Su Lin Kang, PhD; Timothy Barkham, MBBS, MSc, FRCPath; Adeline Oi Kheng Leong, MSc; Ramona Alikiiteaga Gutiérrez, DVM, PhD; and Lee Ching Ng, PhD, have disclosed no relevant financial relationships.

Author affiliations: National Environment Agency, Singapore (M.L. Chau, M. Yap, S.H.P. Hartantyo, Y. Ng, K.T. Aung, C. Goh, J.S.L. Kang, A.O.K. Leong, R.A. Gutiérrez, L.C. Ng); Genome Institute of Singapore, Singapore (S.L. Chen); National University of Singapore, Singapore (S.L. Chen, K.S. Mehershahi); Agri-Food and Veterinary Authority of Singapore, Singapore (P.K.T. Chiew,

C.J. Fernandez, W.K. Wong, R.K. Fong, W.L. Tan, B.Z.Y. Tan); Nanyang Technological University, Singapore (P.K.T. Chiew, L.C. Ng); Tan Tock Seng Hospital, Singapore (T. Barkham)

DOI: https://doi.org/10.3201/eid2312.170596

¹These authors contributed equally to this article.

We assessed microbial safety and quality of raw fish sold in Singapore during 2015–2016 to complement epidemiologic findings for an outbreak of infection with group B *Streptococcus* serotype III sequence type (ST) 283 associated with raw fish consumption. Fish-associated group B *Streptoccocus* ST283 strains included strains nearly identical (0–2 single-nucleotide polymorphisms) with the human outbreak strain, as well as strains in another distinct ST283 clade (57–71 single-nucleotide polymorphisms). Our investigations highlight the risk for contamination of freshwater fish (which are handled and distributed separately from saltwater fish sold as sashimi) and the need for improved hygienic handling of all fish for raw consumption. These results have led to updated policy and guidelines regarding the sale of ready-to-eat raw fish dishes in Singapore.

A major outbreak of group B *Streptococcus* (GBS) infection associated with consumption of a Chinese-style raw fish dish (*yusheng*) occurred in Singapore during 2015 and involved 238 persons during the first half of the year (1). The *yusheng* was typically made from sliced Asian bighead carp (*Hypophthalmichthys nobilis*) and snakehead (*Channa* spp.) and served as a side dish with porridge by food stalls within larger eating establishments. Persons with severe clinical cases had meningoencephalitis, bacteremia, and septic arthritis (2–4). GBS, or *Streptococcus agalactiae*, was identified as the causative agent (2,3).

GBS is found in $\approx 30\%$ of healthy adults (5) and is a member of the human commensal gastrointestinal and genitourinary flora (4). GBS is a common cause of neonatal sepsis, is acquired by newborns from the vaginal flora of the mother, and is an increasingly common pathogen among vulnerable populations (6). The incidence of invasive disease in adults, particularly older adults, has been increasing (7,8). GBS is also a fish and bovine pathogen (9). Although GBS has been shown to colonize the gastrointestinal tract of humans linked to fish consumption (9), foodborne transmission leading to invasive disease has not been reported. Local epidemiologic investigations conducted separately (2,3)identified a single strain of GBS serotype III sequence type (ST) 283 as the causative agent of the outbreak in Singapore during 2015. GBS ST283 had previously been isolated from tilapia in Thailand (10) and in adult human cases in Hong Kong (11). However, GBS ST283 has not been reported to colonize the human gastrointestinal tract, although to date only 1 study of fish mongers and fish handlers has specifically looked for colonization by this strain (12).

We investigated microbial safety and quality of fish sold in the Singapore market during and after the outbreak during 2015 to trace the source of GBS ST283 and provide risk assessment data to support outbreak control and prevention measures. Shortly after identification of GBS ST283 as the cause of the outbreak, these data supported implementing a ban on the sale of ready-to-eat (RTE) dishes containing raw freshwater fish, as well as imposing additional requirements for sale of RTE raw fish dishes made with saltwater fish (13). We report the results of our analysis, which might assist the review of guidelines for handling of fish meant for raw consumption in Singapore and other countries. This report offers unique food and environmental insights into the investigation of this outbreak and complements published epidemiologic findings (2,3).

Materials and Methods

Collection of Fish and Fish Tank Water Samples

We collected samples of fish commonly used for raw consumption (n = 997) and fish tank water for holding live freshwater fish (n = 102) along the supply chain in Singapore during August 2015–January 2016 (online Technical Appendix Table 1, https://wwwnc.cdc.gov/EID/ article/23/12/17-0596-Techapp1.pdf). We tested samples for GBS, *Aeromonas* spp., *Listeria monocytogenes*, *Salmonella* spp., *Vibrio cholerae*, and *V. parahaemolyticus*, and determined *Escherichia coli* counts, *Staphylococcus aureus* counts, and standard plate counts (SPCs) (online Technical Appendix). We characterized selected species to determine their virulence potential (online Technical Appendix).

Statistical Analysis

We evaluated significant differences (p<0.05) between bacterial counts (log₁₀ CFU/g) and presence of specific foodborne bacteria by using Kruskal-Wallis, Mann-Whitney, χ^2 , and Fisher exact tests as appropriate. We performed analysis by using SPSS version 24.0 software (IBM, Armonk, NY, USA).

Results

Raw Fish Samples from Food Stalls and Restaurants/Snack Bars

Although raw freshwater and saltwater fish were served as RTE food at food stalls, only raw saltwater fish were reportedly served at restaurants/snack bars. PCR positivity rates were 43.5% (20/46) for GBS and 23.9% (11/46) for GBS serotype III in sliced fish samples from food stalls. Fish sampled from restaurants/snack bars had significantly lower rates (p<0.05) of 9.2% (26/282) for GBS and 0.7% (2/282) for GBS serotype III (Table). Among the 20 GBS PCR-positive samples from food stalls, 5 yielded isolates; these isolates were of serotype II ST652, serotype III ST283, serotype III ST335, and serotype V ST1 (online Technical Appendix Table 3). The GBS ST283 isolated was from a RTE sliced fish sample sold as grass carp collected from a food stall linked to a human case, as described (*12*). We did not detect GBS ST283 in samples from restaurant/snack

bars; however, we did find a range of other GBS, including serotypes Ia ST7, Ia ST103, Ia ST485, III ST651, III ST861, V ST1, V ST24, VI ST167, and VII ST1.

We found *Salmonella* serogroup B ST29 (serovar Stanley) (n = 2); *V. parahaemolyticus* (negative for *tdh*, *trh1*, and *trh2* genes) (n = 1); and non–O1 *V. cholerae* (n = 1) in freshwater fish samples from food stalls. We also isolated *V. cholerae* from saltwater fish samples, 1 from a food stall and 1 from a restaurant. We detected *L. monocytogenes* in 5 samples from restaurants/snack bars.

SPCs of most RTE raw freshwater (71.4%, 5/7) and saltwater (85.7%, 18/21) fish samples from food stalls exceeded the regulatory limit for RTE food ($5 \log_{10} \text{CFU/g}$) in Singapore (14). We observed no difference in SPCs for fish slices intended for raw consumption and cooking purposes (Figure 1). We also found that 24.8% (70/282) of saltwater fish samples from restaurants/snack bars did not comply with regulatory limits for SPCs, *E. coli* counts (1.3 log₁₀ CFU/g), or both (14). These results showed the poor quality of RTE raw freshwater and saltwater fish sold at food stalls in comparison to those sold at restaurants and snack bars.

Comparison of Freshwater and Saltwater Fish Samples from Fresh Produce Markets

Fish sold at food stalls were typically procured from local fresh produce markets. For the 62 samples of whole fish and fish parts we collected from these markets, we detected GBS ST283 in 28.2% (11/39) of the freshwater fish (Table), which included fish sold as tilapia, Asian bighead carp, grass carp, snakehead-haruan, snakeheadtoman, and silver carp (online Technical Appendix Table 3). However, we did not detect GBS ST283 in saltwater fish. Other GBS strains detected among these fish include serotypes Ia ST7, Ia ST23, Ia ST24, and II ST28 (online Technical Appendix Table 3).

We detected *Aeromonas* spp. (48.4%, 30/62), *S. au*reus (27.4%, 17/62), non–O1 *V. cholerae* (12.9%, 8/62) and *V. parahaemolyticus* (negative for *tdh*, *trh1*, and *trh2* genes) (6.4%, 4/62) in fish samples from fresh produce markets. There was no difference in positivity rates of these organisms between freshwater and saltwater fish. We did not detect *L. monocytogenes* or *Salmonella* spp. in any fish samples collected from fresh produce markets.

Approximately 42% (15/36) of freshwater fish muscle samples had SPCs or *E. coli* counts, or both, exceeding regulatory limits for RTE food in Singapore (14). Positivity rates for GBS, GBS serotype III, and *E. coli*, as well as SPCs for saltwater fish, were significantly lower (p<0.05) (Figure 1; Table). *E. coli* and *S. aureus* counts for freshwater fish surfaces were significantly higher (p<0.05) than those for saltwater fish (Figure 2).

We collected 4 fish tank water samples from wet markets and supermarkets. One water sample and the live



Figure 1. SPCs for sliced fish samples collected from various eating establishments during investigation of group B *Streptococcus* infections, Singapore, 2015–2016. Dashed horizontal line indicates regulatory limit of Singapore for SPCs for ready-to-eat foods (<5 \log_{10} CFU/g) (*14*). Top and bottom of boxes in plots indicate 25th and 75th percentiles, horizontal lines indicate medians, and whiskers indicate minimum and maximum values. *p<0.05. †Food stalls housed within larger eating establishments that include hawker centers, coffee shops, and eating houses. Open circle indicates an outlier. NS, not significant (p>0.05); RTE, ready to eat; SG, Singapore government; SPCs, standard plate counts.

freshwater fish the tank contained were positive for GBS by PCR and non–O1 *V. cholerae* by culture; the associated fish was positive for GBS ST283 by culture. Two other fish tank water samples and the live fish the tanks contained were positive for *E. coli*, *S. aureus*, or both. The level of *E. coli* detected in each positive fish tank water sample was 1.3 log10 CFU/500 mL, which was greater than the l log₁₀ CFU/500 mL coliform (which includes *E. coli*) limit set by the British Columbia Centre for Disease Control (*15*).

Whole Freshwater Fish and Fish Tank Water from Ports

We tested for GBS only in whole fish and fish tank water samples collected from ports. We detected GBS ST283 in 1% (6/586) of freshwater fish samples; positive samples were from Asian bighead carps imported from and farmed in Malaysia. For 98 fish tank water samples collected from ports, 55.1% (54/98) were positive for GBS, and 6.1% (6/98) were positive for GBS ST283. Three of the GBS ST283–positive fish were kept in fish tank water that was also positive for GBS ST283.

Comparison of Saltwater Fish from Fresh Produce Markets and Sashimi Suppliers

Our data indicate the risk for contamination of fish sold at local fresh produce markets, although saltwater fish samples from fresh produce markets had lower rates of

Table. T Ositivity fates	Targeted bacteria, no, positive samples/no, tested (%)									
		GBS	GBS		no. poolar	o oumpioo		. (/0)		
		serotype	serotype	Aeromonas		S.				Salmonella
Characteristic	All GBS	III	III ST283	spp.†	E. coli	aureus	V. c.	V. p.	L. m.	spp.
Detection method	PCR	PCR	Culture, PCR	Culture	Culture	Culture	Culture	Culture	Culture	Culture
Ports										
Freshwater fish, n = 586	27/586 (4.6)	12/586 (2.0)	6/586 (1.0)	NT	NT	NT	NT	NT	NT	NT
Fresh produce markets	s‡									
Freshwater fish,	30/39	14/39	11/39	16/39	32/39	11/39	6/39	2/39	0/39	0/39
n = 39	(76.9) ^a	(35.9) ^b	(28.2)	(41.0)	(82.0) ^c	(28.2)	(15.4)	(5.1)		
Saltwater fish,	5/23	2/23	0/23	14/23	8/23	6/23	2/23	2/23	0/23	0/23
n = 23	(21.7) ^a	(8.7) ^b		(60.9)	(34.8) ^c	(26.1)	(8.7)	(8.7)		
Sashimi suppliers§										
Saltwater fish,	0/21	0/21	0/21	10/21	1/21	0/21	0/21	0/21	1/21	0/21
n = 21				(47.6)	(4.7) ^c				(4.7)	
Food stalls¶										
RTE freshwater	5/7	4/7	1/7	NT	0/7	0/7 (0)	0/7	0/7	0/7	1/7
fish, n = 7	(71.4)	(57.1)	(14.3)							(14.3)
Freshwater fish	8/18	4/18	0/18	NT	2/18	0/18	1/18	1/18	0/18	1/18
for cooking, n = 18	(44.4)	(22.2)			(11.1)		(5.6)	(5.6)		(5.6)
RTE saltwater fish,	7/21	3/21	0/21	NT	0/21	0/21	1/21	0/21	0/21	0/21
n = 21	(33.3) ^d	(14.3) ^e					(5.0)			
Restaurants, snack ba	rs									
RTE saltwater fish,	26/282	2/282	0/282	NT	0/282	0/282	1/282	0/282	5/282	0/282
n = 282	(9.2) ^d	(0.7) ^e					(0.4)		(1.8)	

*A sample was considered positive when a specific organism was detected in ≥1 subsamples (surface, muscle, or organs) of a fish sample. Superscript letters a-e indicate a significant difference (p<0.05) in positivity rates of targeted bacteria between fish types. E. coli, Escherichia coli; GBS, group B Streptococcus; L. m., Listeria monocytogenes; NT, not tested; RTE, ready to eat; S. aureus, Staphylococcus aureus; ST283, sequence type 283; V. c., Vibrio cholerae; V. p., V. parahaemolyticus.

†Aeromonas caviae, A. hydrophila, and A. sobria

‡Fish stalls at ports and wet markets, and fresh produce sections of supermarkets, excluding sashimi and sushi counters of supermarkets.

Companies that supplied sashimi-grade fish to restaurants and snack bars.

¶Within larger eating establishments that include hawker centers, coffee shops, and eating houses

contamination than freshwater fish samples. The SPCs and the positivity rates for E. coli in saltwater fish samples from sashimi suppliers were significantly lower (p<0.05) than those for saltwater fish samples from fresh produce markets (Figure 3; Table), which suggested that the microbial quality of fish could be managed by improvements in handling throughout distribution channels. None of the saltwater fish muscle samples from sashimi suppliers exceeded the Singapore SPC (5 log₁₀ CFU/g) and *E. coli* $(1.3 \log_{10} \text{CFU/g})$ limits for RTE food (14). We did not detect GBS, S. aureus, V. cholerae, and V. parahaemolyticus in any fish samples collected from sashimi suppliers. However, we detected L. monocytogenes in 1 salmon sample.

Characterization of GBS Isolates

We detected 6 GBS serotypes (Ia, II, III, V, VI, and VII) and 13 STs (1, 7, 23, 24, 28, 103, 167, 283, 335, 485, 651, 652, and 861) in fish (online Technical Appendix Table 3). Although most strains were within clonal complexes (1, 10, 17, 19, and 23) associated with human carriage and diseases (16), a total of 20 isolates from 7 sashimi samples (SGE-HI2015-IV45, SGEHI2015-IV72, SGEHI2015-IV74, SGE-HI2015-IV89, SGEHI2015-IV100, SGEHI2015-IV211,

and SGEHI2015-IV232) did not belong to these clonal complexes. These strains had few closely related strains in the public genomic databases, and the closely related strains are mostly from animals (online Technical Appendix Figure 2).

We found GBS ST283 only among freshwater fish and water for holding freshwater fish. Genomic analyses indicated that GBS ST283 isolated from fish clustered in 2 clades (Figure 4). The first clade included 12 isolates from 6 fish from a food stall, a fresh produce market and a port, and 4 fish tank water samples from a port. Genome sequencing showed that these 12 isolates were nearly identical (0-2 SNPs and 0, 1, and 12 indels all in homopolymeric runs of >4 nt) compared with the 2.1-Mbp genome of the reference human outbreak strain, SG-M1, isolated from a meningitis patient during the GBS outbreak in Singapore during 2015 (12,17). Isolates that clustered into the second clade were 20 isolates from 12 fish and 2 fish tank water samples and did not include any human isolates either from this outbreak or from previous reports of human GBS infecting isolates. Sequences of these isolates showed higher intraclade diversity (57-71 SNPs and 11-33 indels) when compared with the SG-M1 genome (Figure 4).



Figure 2. Escherichia coli (A and C) and Staphylococcus aureus (B and D) counts in fish samples (muscle and surface swabs) collected from fresh produce markets during investigation of group B Streptococcus infections, Singapore, 2015–2016. Solid horizontal lines indicate ICMSF limit for E. coli count in fresh fish intended for cooking (<2.7 log₁₀ CFU/g or <4.7 log10 CFU/100 cm2) (23). Dashed horizontal line indicates Singapore regulatory limit for E. coli count in ready-to-eat foods (<1.3 log₁₀ CFU/g) (14). Top and bottom of boxes in plots indicate 25th and 75th percentiles, horizontal lines indicate medians, and whiskers indicate minimum and maximum values. Open circles indicate outliers. *p<0.05. +Companies that supplied sashimi grade fish to restaurants and snack bars. ‡Fish stalls at ports and wet markets, as well as fresh produce sections of supermarkets, excluding sashimi and sushi counters of supermarkets. ICMSF, International Commission on Microbiological Specifications of Foods; NS, not significant (p>0.05); SG, Singapore government; SPCs, standard plate counts.

Characterization of *S. aureus, V. cholerae*, and *V. parahaemolyticus* Isolates

We characterized 18 *S. aureus* isolates from 17 fish. All except 1 were obtained from fish surfaces. We detected >1 enterotoxin gene in two thirds of these isolates and the *sec* gene in 55.6% (10/18) of the isolates. Other enterotoxin genes (*sea, seg, seh, sei,* and *sel*) were detected at much lower rates (5.6% [1/18] to 11.1% [2/18]). We detected 4 enterotoxin genes (*sec, seg, sei,* and *sel*) in a *S. aureus* isolate obtained from the surface of a wolf herring sample collected from a port. We did not detect virulence genes (*ctxA, ctxB,* and *tcpA*) in any of the 16 non–O1 *V. cholerae* isolates from 9 fish and 1 fish tank water samples and did not detect virulence genes (*tdh, trh1,* and *trh2*) in any of the 6 *V. parahaemolyticus* isolates from 5 fish samples.

Discussion

We found GBS ST283, the causative strain of a severe foodborne outbreak in Singapore, in the local freshwater fish supply chain that stretches from food stalls to local fresh produce markets and back to ports. Patients with GBS ST283 infections during this outbreak were more likely to show development of meningoencephalitis, septic arthritis, and spinal infection than were persons with non–GBS ST283 infections (12). Although this study suggested Malaysia as a source of the strain, the finding of the same ST in Hong Kong and Thailand (10,11) suggested that GBS ST283 is generally prevalent throughout the region.

Our analysis shows that there are at least 2 clades of GBS ST283 strains among fish in local markets. Fish and water strains from 1 clade were nearly identical to clinical strains from this outbreak (Figure 4). The small variability of 0–2 SNPs and 0–12 indels between fish and water strains and the reference human outbreak strain (SG-M1) is equivalent to variability observed in 131 clinical strains from the same outbreak reported elsewhere (0–5 SNPs from the SG-M1 reference) (*12*). Strains from a second clade of GBS ST283 had a difference of 57–71 SNPs and 11–33 indels when compared with the SG-M1 genome. Other GBS ST283 isolates, many collected in Hong Kong \leq 17 years before this outbreak (*11*) are also different from the SG-M1 strain (\leq 129 SNPs) (Figure 4). We found no human-infecting isolate from Singapore or elsewhere within the second fish-associated GBS ST283 clade.

A major issue is whether all GBS ST283 strains are capable of causing invasive human disease by the foodborne route. If strains from the fish-associated clade are not pathogenic to humans, they could be used as effective controls for identifying the genetic basis of pathogenicity



Figure 3. SPCs for fish samples (muscle) collected from fresh produce markets during investigation of group B *Streptococcus* infections, Singapore, 2015–2016. Solid horizontal line indicates ICMSF limit for SPCs in fresh fish intended for cooking (<7 log₁₀ CFU/g) (23). Dashed horizontal line indicates Singapore regulatory limit for SPCs for ready-to-eat foods (<5 log₁₀ CFU/g) (*14*). Top and bottom of boxes in plots indicate 25th and 75th percentiles, horizontal lines indicate medians, and whiskers indicate minimum and maximum values. Open circles indicate outliers. *p<0.05. †Companies that supplied sashimi grade fish to restaurants and snack bars. ‡Fish stalls at ports and wet markets, as well as fresh produce sections of supermarkets, excluding sashimi and sushi counters of supermarkets. ICMSF, International Commission on Microbiological Specifications of Foods; SG, Singapore government; SPCs, standard plate counts.

of the first clade and the cause of its emergence, which resulted in outbreak in Singapore in 2015. If these strains are pathogenic to humans, then broader tracking of the prevalence of GBS ST283 would be warranted.

In contrast to GBS strains that are known to cause disease outbreaks in fish (10,18), the live and whole fish from which GBS ST283 was recovered in this study did not have defects, such as corneal opacity and exophthalmia (18), which suggests that this ST might not be pathogenic for freshwater fish. The closest GBS fish pathogens with published genomes, GD201008–001 (19) and HN016 (20), are serotype Ia ST7 strains that are distant (>4,000 SNPs) from all ST283 strains that our group and others have identified (12).

Detection of 6 GBS serotypes and 13 STs showed the diversity of GBS strains in fish. Although the sample size in this study was small and our results might not represent the distribution of GBS in all fish species, our findings provide valuable data for characterizing the public health risk from consuming raw fish. No baseline information on GBS in fish was publicly available before this outbreak because fish were not a recognized source or a recognized route of transmission of GBS. Further work on GBS STs other than ST283 is underway to investigate the role of fish as a source of GBS disease in humans.

Several GBS strains from sashimi had relatively few closely related strains in the public genomic databases (online Technical Appendix Figure 2), which suggests that the GBS population associated with saltwater fish could be different from that associated with freshwater fish and humans. Another reason for this observation is that GBS from food and environmental sources are relatively undersampled in the genomic databases than those from humans.

We detected GBS serotypes Ia ST23 and Ia ST7, which are associated with human carriage (10), in fish samples. Although GBS ST7 has been described as a fish pathogen, the presence of GBS serotype Ia ST23 has not been reported in fish (10). GBS serotypes Ia ST23, and Ia ST7 and *E. coli*, which are all associated with human gut flora, suggest possible contamination of fish by effluent water.

The intentional introduction of animal feces into fish ponds as part of integrated farming (21,22) might further contribute to the complex flow of pathogens between animals and humans. Such findings point to areas for research to clarify the diversity and role of GBS strains in affecting animal and human health. For instance, GBS ST861, which was isolated from salmon in this study (online Technical Appendix Table 3), was also isolated from a clinical case in the same year in Singapore on the basis of metadata available in the PubMLST *S. agalactiae* database (http://pubmlst.org/sagalactiae/).

In addition to the finding of GBS ST283 in freshwater fish, detection of high SPCs and E. coli and S. aureus counts indicates the hazard of using such fish for preparing raw RTE dishes. Because E. coli is not part of the intestinal flora of cold-blooded animals (23), its presence suggests contamination from polluted water, unhygienic handling, or temperature abuse after harvesting. Similarly, because S. aureus is not part of usual fish flora, its presence on fish surfaces suggests possible transfer of human skin flora caused by unhygienic handling (24). We detected V. parahaemolyticus, an organism known to grow well in seawater but lyse rapidly in freshwater (24), in freshwater fish samples from fresh produce markets (5.1%, 2/39). This finding was not surprising because freshwater and saltwater fish are typically sold, handled, stored, and degutted within the same confined areas in fresh produce markets. Thus, despite lower SPCs and positive rates for E. coli in saltwater fish than in freshwater fish from fresh produce markets, saltwater fish procured from such environments are prone to cross-contamination.

Fish used by food stalls were generally obtained from such markets. Moreover, microbial counts for sliced fish samples from eating establishments indicated that most food stalls were not able to prepare RTE raw fish dishes of acceptable hygiene quality. Poor practices observed included

use of common chopping boards, knives, or slicers for preparing fish slices meant for raw consumption and cooking. If fish slices are contaminated, rinsing with water cannot improve their quality (online Technical Appendix).

In contrast to the quality of saltwater fish samples from fresh produce markets, all saltwater fish samples from sashimi suppliers complied with local SPCs (5 \log_{10} CFU/g) and *E. coli* (1.3 \log_{10} CFU/g) limits for readyto-eat food (*14*); all samples were negative for GBS, *S. aureus*, *Salmonella* spp., and *V. parahaemolyticus*. The compliance rate among restaurants/snack bars was higher because such premises are more likely to procure fish from



Figure 4. Phylogenetic analysis of group B *Streptococcus* (GBS) infections caused by improper sourcing and handling of raw fish for raw consumption, Singapore, 2015–2016. A) Maximum-likelihood single-nucleotide polymorphism (SNP)–based tree for GBS ST283 strains relative to the SG-M1 reference human outbreak strain. Year, location, and source (human or fish) for isolates are indicated. Twelve strains from 6 fish (SGEHI2015-NWC941, SGEHI2015–95, LG01, LG02, LG04, and LG06) and 4 fish tank water samples (LG07, LG08, LG09, and LG11) were nearly identical to the local reference outbreak strain SG-M1 (no SNP, 0 and 12 indels, respectively). Scale bar indicates distance over 4,093 total SNPs. B) Enlargement of blue subtree from bottom of tree in panel A showing fish GBS ST283 isolates that were different (57–71 SNPs and 11–33 indels) from the human outbreak strain. ST, sequence type.

sashimi suppliers that harvest fish from cleaner waters and adhere to stricter cold chain management practices. However, some saltwater fish samples from sashimi suppliers and restaurants were found to contain *Aeromonas* spp. (47.6%, 10/21) and *L. monocytogenes* (2.0%, 6/303), whose psychrotrophic nature has posed a challenge to the fish industry. *L. monocytogenes* is also a concern in chilled RTE food because of its ubiquity and persistence in food-processing environments (25).

Food and environmental findings of our study were consistent with epidemiologic findings for this outbreak (2,3). Multivariate analyses of a case–control study showed that persons who had consumed *yusheng* were more likely to acquire GBS ST283 infections than those who had not consumed *yusheng* (2). However, there was no strong association between GBS ST283 infections and consumption of sashimi and sushi (2).

Findings of this study have led to implementation of new policies in Singapore. These new policies included banning the use of freshwater fish in RTE dishes and requiring procurement of saltwater fish from suppliers for raw fish approved by the Agri-Food and Veterinary Authority of Singapore. Food stalls and food establishments providing catering services were required to stop selling RTE raw fish dishes until they complied with practices required for preparing RTE raw saltwater fish dishes.

The number of RTE fish samples collected from food stalls was limited because eating establishments were advised to stop the sale of RTE raw fish dishes containing Asian bighead carp and snakehead during July 24–December 5, 2015, while the outbreak investigation was underway (I). Sampling was not random because it was part of an outbreak investigation, but it was biased toward fish species and food stalls implicated in the outbreak. Thus, contamination rates might not reflect contamination rates of all fish species sold for raw consumption in the market. Similarly, testing of samples from ports and retail outlets was performed by using different protocols, which limited comparisons that could be made.

In conclusion, we detected GBS ST283, which caused a severe foodborne outbreak in Singapore in 2015, in freshwater fish, not only in food stalls and fish markets, but also in ports from which fish are imported. Comparison of human and fish isolates showed as few as 0-2 SNPs between human and fish isolates of GBS ST283 on a background of a diversity of GBS and other bacteria in freshwater fish. These data indicate the risk for contamination of raw freshwater fish and underscore the need for proper sourcing and handling of all fish for raw consumption. To control the outbreak, a ban on the sale of RTE raw freshwater fish dishes was implemented, and additional requirements were imposed for the sale of RTE raw fish dishes made with saltwater fish (*13*). Our

study complements the epidemiologic findings for this outbreak (2,3) and illustrates the need for public health authorities and industries to remain vigilant regarding emerging pathogens.

Acknowledgments

We thank the Agri-Food and Veterinary Authority of Singapore, the National Environment Agency, and the Ministry of Health of Singapore for providing operational support and Lyn Gilbert and Fanrong Kong for providing the GBS serotype VIII reference strain.

Analyses of fish and fish tank water samples was supported by the National Environment Agency and the Agri-Food and Veterinary Authority of Singapore. Whole-genome sequencing and analysis was supported by the Genome Institute of Singapore, the Agency for Science, Technology, and Research, and the Singapore Ministry of Health National Medical Research Council (NMRC/CIRG/1357/2013).

Ms. Chau is research group leader of the Applied Food Hygiene Epidemiology Group, Food and Environmental Hygiene Research Program, Environmental Health Institute, National Environment Agency, Singapore. Her research interests include microbiology, food hygiene and safety, and foodborne diseases.

References

- Joint news release between the Ministry of Health Singapore (MOH) and the Agri-Food and Veterinary Authority of Singapore (AVA) and the National Environment Agency (NEA). Update on investigation into group B *Streptococcus* cases. July 24, 2015 [cited 2016 Aug 10]. http://www.nea.gov.sg/corporate-functions/newsroom/news-releases/year/2015/month/7/category/food-hygiene/ update-on-investigation-into-group-b-streptococcus-cases
- Tan S, Lin Y, Foo K, Koh HF, Tow C, Zhang Y, et al. Group B Streptococcus serotype III sequence type 283 bacteremia associated with consumption of raw fish, Singapore. Emerg Infect Dis. 2016;22:1970–3. http://dx.doi.org/10.3201/eid2211.160210
- Rajendram P, Mar Kyaw W, Leo YS, Ho H, Chen WK, Lin R, et al. Group B Streptococcus sequence type 283 disease linked to consumption of raw fish, Singapore. Emerg Infect Dis. 2016; 22:1974–7. http://dx.doi.org/10.3201/eid2211.160252
- Tan K, Wijaya L, Chiew HJ, Sitoh YY, Shafi H, Chen RC, et al. Diffusion-weighted MRI abnormalities in an outbreak of *Streptococcus agalactiae* serotype III, multilocus sequence type 283 meningitis. J Magn Reson Imaging. 2017;45:507–14. http://dx.doi.org/10.1002/jmri.25373
- Manning SD, Neighbors K, Tallman PA, Gillespie B, Marrs CF, Borchardt SM, et al. Prevalence of group B *Streptococcus* colonization and potential for transmission by casual contact in healthy young men and women. Clin Infect Dis. 2004;39:380–8. http://dx.doi.org/10.1086/422321
- Public Health Agency of Canada. Streptococcus agalactiae: pathogen safety data sheet—infectious substances. April 30, 2012 [cited 2016 Aug 10]. http://www.phac-aspc.gc.ca/lab-bio/res/ psds-ftss/streptococcus-agalactiae-eng.php
- Ballard MS, Schønheyder HC, Knudsen JD, Lyytikäinen O, Dryden M, Kennedy KJ, et al.; International Bacteremia Surveillance Collaborative. The changing epidemiology of group B Streptococcus bloodstream infection: a multi-national population-

based assessment. Infect Dis (Lond). 2016;48:386–91. http://dx.doi.org/10.3109/23744235.2015.1131330

- Phares CR, Lynfield R, Farley MM, Mohle-Boetani J, Harrison LH, Petit S, et al.; Active Bacterial Core surveillance/Emerging Infections Program Network. Epidemiology of invasive group B streptococcal disease in the United States, 1999–2005. JAMA. 2008;299:2056–65. http://dx.doi.org/10.1001/jama.299.17.2056
- Foxman B, Gillespie BW, Manning SD, Marrs CF. Risk factors for group B streptococcal colonization: potential for different transmission systems by capsular type. Ann Epidemiol. 2007; 17:854–62. http://dx.doi.org/10.1016/j.annepidem.2007.05.014
- Delannoy CM, Crumlish M, Fontaine MC, Pollock J, Foster G, Dagleish MP, et al. Human *Streptococcus agalactiae* strains in aquatic mammals and fish. BMC Microbiol. 2013;13:41. http://dx.doi.org/10.1186/1471-2180-13-41
- Ip M, Cheuk ES, Tsui MH, Kong F, Leung TN, Gilbert GL. Identification of a *Streptococcus agalactiae* serotype III subtype 4 clone in association with adult invasive disease in Hong Kong. J Clin Microbiol. 2006;44:4252–4. http://dx.doi.org/10.1128/ JCM.01533-06
- Kalimuddin S, Chen SL, Lim CTK, Koh TH, Tan TY, Kam M, et al.; Singapore Group B Streptococcus Consortium. 2015 epidemic of severe *Streptococcus agalactiae* ST283 infections in Singapore associated with the consumption of raw freshwater fish: a detailed analysis of clinical, epidemiological and bacterial sequencing data. Clin Infect Dis. 2017;64(Suppl_2):S145–52. http:// dx.doi.org/10.1093/cid/cix021
- 13. Joint news release between the Agri-Food and Veterinary Authority of Singapore (AVA) and the Ministry of Health Singapore (MOH) and the National Environment Agency (NEA). Freshwater fish banned from ready-to-eat raw fish dishes. December 5, 2015 [cited 2016 Aug 10]. http://www.nea.gov.sg/corporate-functions/ newsroom/news-releases/freshwater-fish-banned-from-ready-toeat-raw-fish-dishes
- Agri-Food and Veterinary Authority of Singapore. Sale of food act, chapter 283, section 56 (1), food regulations. December 20, 2016 [cited 2017 Jan 9]. http://www.ava.gov.sg/legislation
- British Columbia Centre for Disease Control. Food processing plants: guideline for live retail fish holding system. November 2013 [cited 2016 Aug 17]. http://www.bccdc.ca/resource-gallery/ Documents/Educational%20Materials/EH/FPS/Food/ RetailFishHoldingTankGuidelines_Nov2013trs.pdf
- Da Cunha V, Davies MR, Douarre PE, Rosinski-Chupin I, Margarit I, Spinali S, et al.; DEVANI Consortium. *Streptococcus agalactiae* clones infecting humans were selected and fixed through the extensive use of tetracycline. Nat Commun. 2014;5:4544. http://dx.doi.org/10.1038/ncomms5544

- Mehershahi KS, Hsu LY, Koh TH, Chen SL. Complete genome sequence of *Streptococcus agalactiae* serotype III, multilocus sequence type 283 strain SG-M1. Genome Announc. 2015; 3:e01188–15.
- Abuseliana AF, Daud HH, Aziz SA, Bejo SK, Alsaid M. Pathogenicity of *Streptococcus agalactiae* isolated from a fish farm in Selangor to juvenile red tilapia (*Oreochromis* sp.). Journal of Animal and Veterinary Advances. 2011;10:914–9. http://dx.doi.org/ 10.3923/javaa.2011.914.919
- Liu G, Zhang W, Lu C. Complete genome sequence of *Streptococcus agalactiae* GD201008-001, isolated in China from tilapia with meningoencephalitis. J Bacteriol. 2012;194:6653. http://dx.doi.org/10.1128/JB.01788-12
- Wang R, Li L, Huang Y, Luo F, Liang W, Gan X, et al. Comparative genome analysis identifies two large deletions in the genome of highly-passaged attenuated *Streptococcus agalactiae* strain YM001 compared to the parental pathogenic strain HN016. BMC Genomics. 2015;16:897. http://dx.doi.org/10.1186/ s12864-015-2026-y
- Martins C, Eding EH, Verdegem MC, Heinsbroek LT, Schneider O, Blancheton J-P, et al. New developments in recirculating aquaculture systems in Europe: a perspective on environmental sustainability. Aquacultural Engineering. 2010;43:83–93. http://dx.doi.org/10.1016/ j.aquaeng.2010.09.002
- Little D, Edwards P. Integrated livestock-fish farming systems. Rome: Food and Agriculture Organization of the United Nations; 2003 [cited 2017 Aug 15]. http://www.fao.org/docrep/006/y5098e/ y5098e00.htm
- 23. International Commission on Microbiological Specifications for Foods. Micro-organisms in foods 2: sampling for microbiological analysis; principles and specific applications. 2nd ed. 1986 [cited 2016 Aug 11]. http://www.icmsf.org/pdf/icmsf2.pdf
- Food and Drug Administration. Bad bug book: foodborne pathogenic microorganisms and natural toxins. 2nd ed. Silver Spring (MD): The Administration; 2012.
- 25. Chau ML, Aung KT, Hapuarachchi HC, Lee PSV, Lim PY, Kang JSL, et al. Microbial survey of ready-to-eat salad ingredients sold at retail reveals the occurrence and the persistence of *Listeria monocytogenes* sequence types 2 and 87 in pre-packed smoked salmon. BMC Microbiol. 2017;17:46. http://dx.doi.org/10.1186/ s12866-017-0956-z

Address for correspondence: Lee C. Ng, Environmental Health Institute, National Environment Agency, 11 Biopolis Way, #06-05/08, Helios Block, Singapore 138667, Singapore; email: Ng Lee Ching@nea.gov.sg Article DOI: https://doi.org/10.3201/eid2312.170596

Group B *Streptococcus* Infections Caused by Improper Sourcing and Handling of Fish for Raw Consumption, Singapore, 2015– 2016

Technical Appendix

Materials and Methods

Collection of Fish and Fish Tank Water Samples

Samples from Eating Establishments

Sliced fish samples (n = 328) from food stalls, restaurants, and snack bars were analyzed for microbiological safety and quality between freshwater and saltwater fish slices prepared by these retail eating establishments. Eating establishments linked to human cases (implicated establishments) and eating establishments that sold similar dishes but were not linked to human cases were included (control establishments). However, statistical analyses for implicated and control establishments were not performed because the number of samples from implicated establishments was small (26 food stall samples; 5 restaurant/snack bar samples). Limited samples were available because retailers were advised by health authorities to stop selling readyto-eat raw fish dishes containing Asian bighead carp and snakehead while the outbreak investigation was underway.

Samples from Fresh Produce Markets and Sashimi Suppliers

Although fresh produce markets refer to fish stalls at the ports and wet markets, as well as fresh produce sections of supermarkets that food stalls are known to procure fish from, sashimi suppliers refer to companies that local restaurants and snack bars are known to procure sashimi grade fish from. Fish parts or whole fish (n = 83) were analyzed for microbiological safety and quality for freshwater and saltwater fish sampled at these locations. Subsamples consisting of surface swab specimens, muscles, or organs, were obtained from each whole fish or fish part sample and subjected to microbial analyses (online Technical Appendix Table 2). During sampling of fish at fresh produce markets, samples of water in which live freshwater fish were kept were also collected. Microbial safety and quality of water were analyzed to assess their potential to cause cross-contamination among live fish in the market.

Samples from Ports

A total of 586 whole freshwater fish samples and 98 holding water samples were collected at ports in Singapore. These samples were tested for group B *Streptococcus* (GBS), GBS serotype III, and GBS sequence type (ST) 283.

Transport of Samples

Whole fish and fish part samples were purchased in their original packaging; sliced fish were collected in sterile bags, and fish tank water samples were collected in sterile bottles. All samples were transported on ice, kept refrigerated in laboratories, and analyzed on the day of collection.

Testing of Samples

Microbiological testing of fish and fish tank water samples was performed by 3 laboratories: the Environmental Health Institute (EHI) of the National Environment Agency, the Veterinary Public Health Laboratory of the Agri-Food and Veterinary Authority of Singapore (AVA), and an accredited commercial laboratory. The Veterinary Public Health Laboratory performed testing for GBS in whole fish and fish tank water samples collected from ports, and EHI performed testing for GBS on all other samples. Standard plates counts (SPCs) for *Escherichia coli, Staphylococcus aureus, Listeria monocytogenes, Salmonella* spp., *Vibrio cholerae*, and *V. parahaemolyticus* from fish slices collected from eating establishments was performed by an accredited commercial laboratory. SPCs for *E. coli, S. aureus, Aeromonas* spp., *L. monocytogenes, Salmonella* spp., *V. cholerae*, and *V. parahemolyticus* in whole fish and fish parts collected from ports, wet markets, supermarkets, and sashimi suppliers was performed by EHI. *S. aureus, Salmonella* spp., *V. cholerae*, and *V. parahaemolyticus* isolated by EHI and the accredited commercial laboratory were further characterized at EHI.

GBS Analysis

Fish and fish tank water samples collected from ports were analyzed by Agri-Food and Veterinary Authority of Singapore by using a proprietary method that involved screening for GBS and GBS serotype III with a multiplex PCR method and isolation of viable GBS ST283 by culture method. GBS analyses on all other samples were conducted at EHI by using a modified protocol (*1*). Each fish surface swab sample was prepared by applying a premoistened sponge swab (3M Company, Maplewood. MN, USA) onto the entire surface area of each whole fish or fish part sample. The sponge swab was then suspended in 9 mL of Todd Hewitt broth (Acumedia, Lansing, MI, USA) containing colistin sulfate (7.5 μ g/mL) and nalidixic acid (5 μ g/mL) and homogenized by using a Stomacher (model 400; Seward Medical, Worthing, UK) with a paddle speed of 230 rpm for 30 s. A total of 25 g each of fish muscle sample, fish organ composite sample, and sliced fish sample was homogenized with 225 mL of Todd Hewitt broth containing antimicrobial drugs by using a Stomacher.

Each 500-mL fish tank water sample was concentrated by using membrane filtration with sterile 0.45-µm pore size filters made of mixed cellulose esters (Merck Millipore, Darnstadt, Germany). Membranes were then suspended in 20 mL of Todd Hewitt broth containing antimicrobial drugs. Homogenized aliquots and membranes were incubated at 37°C for 24 h. Secondary enrichment was performed by transferring 1 mL of the enriched Todd Hewitt broth to 5 mL brain heart infusion broth (Acumedia) and incubating at 37°C for 24 h. Nucleic acids were extracted from enriched brain heart infusion broth and selected preenriched sample suspensions in Todd Hewitt broth containing antimicrobial drugs by using a method described (2).

GBS was detected by using a singleplex PCR specific for the *dltR* gene (3). All GBS PCR-positive samples were cultured to obtain viable isolates. Culture involved plating of 100 μ L of serially diluted (0–10⁵ dilutions) enriched brain heart infusion broth onto Brilliance GBS Agar (Thermo Fisher Scientific, Darmstadt, Germany) and incubating the agar at 37°C for 24 h. Presumptive mauve colonies were confirmed as GBS by serologic analysis using the Streptococcal Grouping Kit (Oxoid, Basingstoke, UK) according to the manufacturer's instructions.

Although 103 of 522 enriched samples tested by EHI were positive for GBS by PCR, only 35 of these samples yielded GBS isolates when plated on chromogenic agar; GBS was not

recovered from the other 68 PCR-positive samples. A high background flora could have masked GBS in these 68 enriched samples. This masking is supported by the observation that of the 68 isolate-negative samples, 57 initially showed PCR-negative results, but then showed PCR-positive results after a 2-day enrichment. This change in results indicated low amounts of GBS in the original samples. Presampling manipulation, such as freezing, could have also contributed to the low yield.

Because recovery of viable isolates was not always successful for GBS PCR-positive samples, the enriched broth of these samples was further screened for GBS serotype III by using published primers (4). This screening was performed to estimate the highest possible contamination rate of GBS serotype III ST283 in fish.

A total of 84 GBS isolates were characterized by using serotyping, multilocus sequence typing (MLST), and whole-genome sequencing methods. Serotyping of GBS isolates was performed by using the IMMULEX STREP-B Antisera Set (Staten Serum Institut Diagnostica, Hillerød, Denmark) according to the manufacturer's instructions. Results were confirmed by using a published multiplex PCR protocol (*5*). MLST was performed by using published primers and Sanger sequencing (*6*). Aligned nucleotide sequences of amplified products were compared with sequences in the PubMLST *Streptococcus agalactiae* Database (http://pubmlst.org/sagalactiae/) for identification of sequence types.

Whole-genome sequencing was performed by the Genome Institute of Singapore and the Agency for Science, Technology and Research by using a NextSeq 500 sequencer (Illumina, San Diego, CA, USA) according to protocols described (7). A total of 1 mL of each GBS overnight culture in brain heart infusion broth was centrifuged, and the pellet of bacterial cells was lysed by using enzymatic lysis buffer at 37°C for 45 min, followed by extraction using the DNeasy Blood and Tissue Kit (QIAGEN, Valencia, CA, USA) according to the manufacturer's instructions. Genomic DNA shearing was performed by using an M220 Focused Ultrasonicator (Covaris, Woburn, MA, USA), and library preparation was performed by using the TruSeq Nano DNA LT Library Prep Kit Pooled libraries (Illumina). Samples were then sequenced by using a NextSeq 500 sequencer with 2×151 -bp reads. All sequencing data are available in GenBank under BioProject PRJNA293392.

Sequence Analysis

All primary sequence analysis was performed by the Genome Institute of Singapore with the Efficient Rapid Microbial Sequencing Platform. Reference-based analyses were performed by using the SG-M1 genome (8) as a reference. FASTQ files were mapped by using Burrows-Wheeler Aligner version 0.7.10 software (9). Indel realignment and single-nucleotide polymorphism (SNP) calling was performed by using Lofreq* version 2.1.2 with default parameters (10). Maximum-likelihood SNP trees (ignoring indels) of GBS ST283 strains were created by using SNPhylo (11). Neighbor-joining trees of all GBS strains were created by using APE version 3.5 (12). All phylogenetic trees were visualized with GGTREE version 3.2 (13) in R version 3.2.2 (https://www.R-project.org). Publicly available sequences of GBS were downloaded from the GenBank Short Read Archive (for Illumina datasets; http://www.ncbi.nlm.nih.gov/sra) and the Genbank FTP site (for assembled genomes; https://www.ncbi.nlm.nih.gov/genome/genomes/186?). MLST and resistance gene predictions were made by using SRST2 version 0.1.8(14) for Illumina sequenced strains or manually by using BLASTN (15) for fully assembled reference sequences, the recommended MLST database (http://pubmlst.org/sagalactiae/) (16), and the ARGannot resistance gene database (17) included with SRST2.

SPCs and E. coli and S. aureus Counts

The protocols used by EHI are described below. Each fish surface swab sample was prepared by applying a premoistened sponge swab onto a 10 cm \times 10 cm surface area of each fish part or whole fish sample. The sponge swab was then suspended in 9 mL Butterfield buffer and homogenized by using a Stomacher. A total of 25 g of each fish muscle and sliced fish sample was homogenized with 225 mL of universal preenrichment broth (Acumedia) by using a Stomacher. Each 500-mL fish tank water sample was concentrated by membrane filtration using 0.45-µm pore size filters and suspended in 20 mL of universal preenrichment broth. SPCs and *E. coli* and *S. aureus* counts were determined performed according to described methods (*18*). The presence of the methicillin resistance gene and enterotoxin-producing genes (*sea, seb, sec, sed, see, seg, seh, sei, sej,* and *sel*) of 18 *S. aureus* isolates from 17 fish was determined by using published primers (*19–21*).

Detection of Aeromonas spp., Listeria monocytogenes, Salmonella spp., Vibrio cholerae, and V. parahaemolyticus

Each fish surface swab sample was prepared by applying a premoistened sponge swab onto the entire surface area of each fish part or whole fish sample. The sponge swab was then suspended in 9 mL of universal preenrichment broth and homogenized by using a Stomacher. All stomached aliquots (surface swab specimens, fish muscle, fish organs, sliced fish) resuspended in universal preenrichment broth were incubated at 37°C for 24 h for detection of *Aeromonas* spp. (*A. caviae, A. hydrophila*, and *A. sobria*), *L. monocytogenes, Salmonella* spp., *V. cholerae*, and *V. parahaemolyticus*. A 10-µL loopful of an enriched culture in universal preenrichment broth was subcultured onto starch-ampicillin agar (Himedia, Mumbai, India), PALCAM agar (Acumedia), and xylose-lysine-desoxycholate (XLD) agar (Oxoid) for isolation of *Aeromonas* spp., *L. monocytogenes*, and *Salmonella* spp.; and onto thiosulfate-citrate-bile salt-sucrose agar (Acumedia) for isolation of and *V. cholerae* and *V. parahaemolyticus*. All selective agars were incubated at 37°C for 24 h, except for PALCAM agar, which was incubated at 37°C for 48 h.

After incubation, colonies on starch-ampicillin agar were flooded with diluted (1:20) iodine solution (Oxoid) to detect colonies with zones of clearance. Presumptive *Aeromonas* spp. colonies were streaked onto tryptic soy agar (Acumedia) for purification and biochemical confirmation by using API 20 E (bioMérieux, Marcy l'Étoile, France). Presumptive gray–green colonies surrounded by dark halos on PALCAM agar were picked for identification of *L. monocytogenes* by using a *Listeria* genus–specific PCR and a *L. monocytogenes* species–specific PCR that targeted the *prs* gene (22) and the *inlA* gene (23).

All other bacteria isolates, *Salmonella* spp. (n = 2), *V. cholerae* (n = 16), and *V. parahaemolyticus* (n = 6), were verified biochemically by using described methods (*18*). Sequence types of *Salmonella* spp. isolates were determined by using primers and Sanger sequencing, as well as comparison of sequences in the MLST database (http://mlst.warwick.ac.uk/mlst/dbs/Senterica/documents/primersEnterica_html). Sixteen *V. cholerae* isolates from 9 fish and 1 fish tank water samples were screened for virulence genes (*ctxA*, *ctxB*, and *tcpA*) by using published primers (*24*–26). Six *V. parahaemolyticus* isolates from 5 fish samples were screened for virulence genes (*tdh*, *trh1*, and *trh2*) by using published primers (*27*,*28*). Nucleic acids of all bacterial isolates, including GBS, were extracted by using the DNeasy Blood and Tissue Kit (QIAGEN). Amplified product sizes were analyzed by using

the QIAxcel DNA Screening Kit (QIAGEN) or the QIAxcel DNA High Resolution Kit (QIAGEN) as appropriate.

SPCs in Fish Pieces before and after Rinsing

We analyzed 21 sets of saltwater fish muscle samples to determine if SPC could be reduced by rinsing with water. Each set of samples consisted of 2 muscle pieces (each 5 cm \times 5 cm \times 1 cm). One piece served as a control (before rinsing), and the other piece was rinsed with 500 mL of sterile water. We analyzed fish muscle samples for SPCs according a described method (*18*).

References

- van der Mee-Marquet N, Domelier A-S, Salloum M, Violette J, Arnault L, Gaillard N, et al.; Bloodstream Infection Study Group of the Reseau des Hygienistes de la Region Centre. Molecular characterization of temporally and geographically matched *Streptococcus agalactiae* strains isolated from food products and bloodstream infections. Foodborne Pathog Dis. 2009;6:1177–83. PubMed http://dx.doi.org/10.1089/fpd.2009.0287
- 2. Chau ML, Hartantyo SH, Yap M, Kang JSL, Aung KT, Gutiérrez RA, et al. Diarrheagenic pathogens in adults attending a hospital in Singapore. BMC Infect Dis. 2016;16:32. <u>PubMed</u> <u>http://dx.doi.org/10.1186/s12879-016-1354-0</u>
- 3. Lamy M-C, Dramsi S, Billoët A, Réglier-Poupet H, Tazi A, Raymond J, et al. Rapid detection of the "highly virulent" group B *Streptococcus* ST-17 clone. Microbes Infect. 2006;8:1714–22. <u>PubMed</u> <u>http://dx.doi.org/10.1016/j.micinf.2006.02.008</u>
- 4. Poyart C, Tazi A, Réglier-Poupet H, Billoët A, Tavares N, Raymond J, et al. Multiplex PCR assay for rapid and accurate capsular typing of group B streptococci. J Clin Microbiol. 2007;45:1985–8.
 <u>PubMed http://dx.doi.org/10.1128/JCM.00159-07</u>
- Imperi M, Pataracchia M, Alfarone G, Baldassarri L, Orefici G, Creti R. A multiplex PCR assay for the direct identification of the capsular type (Ia to IX) of *Streptococcus agalactiae*. J Microbiol Methods. 2010;80:212–4. <u>PubMed http://dx.doi.org/10.1016/j.mimet.2009.11.010</u>
- 6. Jones N, Bohnsack JF, Takahashi S, Oliver KA, Chan M-S, Kunst F, et al. Multilocus sequence typing system for group B *Streptococcus*. J Clin Microbiol. 2003;41:2530–6. <u>PubMed</u> <u>http://dx.doi.org/10.1128/JCM.41.6.2530-2536.2003</u>

- Kalimuddin S, Chen SL, Lim CTK, Koh TH, Tan TY, Kam M, et al.; Singapore Group B Streptococcus Consortium. 2015 epidemic of severe *Streptococcus agalactiae* ST283 infections in Singapore associated with the consumption of raw freshwater fish: a detailed analysis of clinical, epidemiological and bacterial sequencing data. Clin Infect Dis. 2017;64(Suppl_2):S145– 52. <u>PubMed http://dx.doi.org/10.1093/cid/cix021</u>
- Mehershahi KS, Hsu LY, Koh TH, Chen SL. Complete genome sequence of *Streptococcus agalactiae* serotype III, multilocus sequence type 283 strain SG-M1. Genome Announc. 2015;3:e01188–15.
 <u>PubMed</u>
- Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics. 2009;25:1754–60. <u>PubMed http://dx.doi.org/10.1093/bioinformatics/btp324</u>
- 10. Wilm A, Aw PPK, Bertrand D, Yeo GHT, Ong SH, Wong CH, et al. LoFreq: a sequence-quality aware, ultra-sensitive variant caller for uncovering cell-population heterogeneity from highthroughput sequencing datasets. Nucleic Acids Res. 2012;40:11189–201. <u>PubMed</u> <u>http://dx.doi.org/10.1093/nar/gks918</u>
- 11. Lee T-H, Guo H, Wang X, Kim C, Paterson AH. SNPhylo: a pipeline to construct a phylogenetic tree from huge SNP data. BMC Genomics. 2014;15:162. <u>PubMed http://dx.doi.org/10.1186/1471-2164-15-162</u>
- Paradis E, Claude J, Strimmer K. APE: analyses of phylogenetics and evolution in R language. Bioinformatics. 2004;20:289–90. <u>PubMed http://dx.doi.org/10.1093/bioinformatics/btg412</u>
- 13. Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y. GGTREE: an R package for visualization and annotation of phylogenetic trees with their covariates and other associated data. Methods in Ecology and Evolution. 2016 [cited 2017 Aug 16]. https://www.researchgate.net/publication/306241714_ggtree_an_R_package_for_visualization_a nd_annotation_of_phylogenetic_trees_with_their_covariates_and_other_associated_data
- 14. Inouye M, Dashnow H, Raven L-A, Schultz MB, Pope BJ, Tomita T, et al. SRST2: Rapid genomic surveillance for public health and hospital microbiology labs. Genome Med. 2014;6:90. <u>PubMed</u> <u>http://dx.doi.org/10.1186/s13073-014-0090-6</u>
- 15. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+: architecture and applications. BMC Bioinformatics. 2009;10:421. <u>PubMed http://dx.doi.org/10.1186/1471-2105-10-421</u>

- Jolley KA, Maiden MC. BIGSdb: scalable analysis of bacterial genome variation at the population level. BMC Bioinformatics. 2010;11:595. <u>PubMed http://dx.doi.org/10.1186/1471-2105-11-595</u>
- 17. Gupta SK, Padmanabhan BR, Diene SM, Lopez-Rojas R, Kempf M, Landraud L, et al. ARG-ANNOT, a new bioinformatic tool to discover antibiotic resistance genes in bacterial genomes. Antimicrob Agents Chemother. 2014;58:212–20. <u>PubMed http://dx.doi.org/10.1128/AAC.01310-13</u>
- 18. Aung KT, Lo JA, Chau ML, Kang JS, Yap HM, Gutiérrez RA, et al. Microbiological safety assessment and risk mitigation of Indian rojak (deep fried ready-to-eat food) in Singapore. Southeast Asian J Trop Med Public Health. 2016;47:1231–45.
- Rosec JP, Gigaud O. Staphylococcal enterotoxin genes of classical and new types detected by PCR in France. Int J Food Microbiol. 2002;77:61–70. <u>PubMed http://dx.doi.org/10.1016/S0168-</u> <u>1605(02)00044-2</u>
- 20. Cremonesi P, Luzzana M, Brasca M, Morandi S, Lodi R, Vimercati C, et al. Development of a multiplex PCR assay for the identification of *Staphylococcus aureus* enterotoxigenic strains isolated from milk and dairy products. Mol Cell Probes. 2005;19:299–305. <u>PubMed</u> <u>http://dx.doi.org/10.1016/j.mcp.2005.03.002</u>
- Strommenger B, Kettlitz C, Werner G, Witte W. Multiplex PCR assay for simultaneous detection of nine clinically relevant antibiotic resistance genes in *Staphylococcus aureus*. J Clin Microbiol. 2003;41:4089–94. <u>PubMed http://dx.doi.org/10.1128/JCM.41.9.4089-4094.2003</u>
- 22. Doumith M, Buchrieser C, Glaser P, Jacquet C, Martin P. Differentiation of the major *Listeria monocytogenes* serovars by multiplex PCR. J Clin Microbiol. 2004;42:3819–22. <u>PubMed</u> <u>http://dx.doi.org/10.1128/JCM.42.8.3819-3822.2004</u>
- 23. Liu D, Lawrence ML, Wiedmann M, Gorski L, Mandrell RE, Ainsworth AJ, et al. *Listeria monocytogenes* subgroups IIIA, IIIB, and IIIC delineate genetically distinct populations with varied pathogenic potential. J Clin Microbiol. 2006;44:4229–33. <u>PubMed http://dx.doi.org/10.1128/JCM.01032-06</u>
- 24. Hoshino K, Yamasaki S, Mukhopadhyay AK, Chakraborty S, Basu A, Bhattacharya SK, et al. Development and evaluation of a multiplex PCR assay for rapid detection of toxigenic *Vibrio cholerae* O1 and O139. FEMS Immunol Med Microbiol. 1998;20:201–7. <u>PubMed</u> <u>http://dx.doi.org/10.1111/j.1574-695X.1998.tb01128.x</u>

- 25. Imani FA, Iman ID, Hosseini DR, Karami A, Marashi SM. Design of a multiplex PCR method for detection of toxigenic-pathogenic in *Vibrio cholerae*. Asian Pac J Trop Med. 2013;6:115–8. <u>PubMed http://dx.doi.org/10.1016/S1995-7645(13)60005-X</u>
- 26. Rivera IN, Chun J, Huq A, Sack RB, Colwell RR. Genotypes associated with virulence in environmental isolates of *Vibrio cholerae*. Appl Environ Microbiol. 2001;67:2421–9. <u>PubMed</u> <u>http://dx.doi.org/10.1128/AEM.67.6.2421-2429.2001</u>
- 27. Bej AK, Patterson DP, Brasher CW, Vickery MC, Jones DD, Kaysner CA. Detection of total and hemolysin-producing *Vibrio parahaemolyticus* in shellfish using multiplex PCR amplification of *tl, tdh* and *trh.* J Microbiol Methods. 1999;36:215–25. <u>PubMed http://dx.doi.org/10.1016/S0167-7012(99)00037-8</u>
- 28. Tada J, Ohashi T, Nishimura N, Shirasaki Y, Ozaki H, Fukushima S, et al. Detection of the thermostable direct hemolysin gene (*tdh*) and the thermostable direct hemolysin-related hemolysin gene (*trh*) of *Vibrio parahaemolyticus* by polymerase chain reaction. Mol Cell Probes. 1992;6:477–87. PubMed http://dx.doi.org/10.1016/0890-8508(92)90044-X
- 29. Ward RD. FISH-BOL, a case study for DNA barcodes. Methods Mol Biol. 2012;858:423–39. <u>PubMed http://dx.doi.org/10.1007/978-1-61779-591-6_21</u>
- 30. Agri-Food and Veterinary Authority of Singapore. Sale of food act, chapter 283, section 56 (1), food regulations. December 20, 2016 [cited 2017 Jan 9]. http://www.ava.gov.sg/legislation
- 31. Da Cunha V, Davies MR, Douarre PE, Rosinski-Chupin I, Margarit I, Spinali S, et al.; DEVANI Consortium. *Streptococcus agalactiae* clones infecting humans were selected and fixed through the extensive use of tetracycline. Nat Commun. 2014;5:4544. <u>PubMed</u> <u>http://dx.doi.org/10.1038/ncomms5544</u>

Technical Appendix Table 1. Characteristics of freshwater and saltwater fish tested during investigation of group B Streptococcus infections, Singapore, 2015-2016

		No.	
Sampling location	Type of sample	samples	Type of fish collected
Noneating establishment			
Port	Whole fish	586	Freshwater fish sold as Asian bighead carp (<i>Hypophthalmichthys nobilis</i>), grass carp (<i>Ctenopharyngodon idella</i>), Pangasius dory
			(<i>Pangasius</i> spp.), snakehead (<i>Channa</i> spp.), and tilapia (<i>Oreochromis</i> spp.)
Fresh produce markets (fish stalls at	Whole fish, fish	39	Freshwater fish sold as Asian bighead carp,
ports and wet markets and fresh produce	parts		snakehead-toman (C. micropeltes), snakehead-
sections of supermarkets)			haruan (<i>C. striata</i>), silver carp (<i>Hypophthalmichthys molitrix</i>), and tilapia
Fresh produce markets (fish stalls at	Whole fish, fish	23	Saltwater fish sold as mackerel tuna (Euthynnus
ports and wet markets and fresh produce sections of supermarkets)	parts		affinis), ribbonfish (<i>Trichiurus</i> spp.), salmon (<i>Salmo</i> salar), and wolf herring (<i>Chirocentrus</i> spp.)
Sashimi suppliers*	Whole fish, fish	21	Saltwater fish sold as salmon and tuna (Thunnus
	parts		spp.)
Eating establishment			
Food stalls in (in hawker centers, coffee	Sliced fish	25	Freshwater fish sold as Asian bighead carp, grass
shops,and eating houses)			carp, Pangasius dory, and snakehead
Food stalls in (in hawker centers, coffee	Sliced fish	21	Saltwater fish sold as salmon and Spanish mackerel
shops,and eating houses)			(Scomberomorus spp.), and wolf herring.
Restaurants and snack bars*	Sliced fish	282	Saltwater fish sold as salmon, tuna, and <i>taï</i> †

*These locations did not sell freshwater fish meant for raw consumption.

†Generally refers to red sea bream (Pagrus major) in authentic Japanese cuisine, but it can be substituted with other fish by retailers in Singapore.

Technical Appendix Table 2. Testing of subsamples during investigation of group B Streptococcus infections, Singapore, 2015-2016*

	Purpose of testing								
				Measurement of					
	Detection of organism	Detection of food safety	Detection of food	total microbial load					
Characteristic	implicated in outbreak	indicators	hygiene indicators	in food					
Testing parameter	GBS, GBS, serotype III, and GBS ST283	Aeromonas spp. (A. caviae, A. hydrophila, and A. sobria); Listeria monocytogenes; Salmonella spp.; Vibrio cholerae and V. parahaemolyticus	Escherichia coli and Staphylococcus aureus counts	Standard plate count					
Type of sample Whole fish, fish parts									
Organs (brain, eyes, spleen,	Yes	No	No	No					
or kidneys)									
Surface swab specimen	Yes	Yes†	Yes†	No					
Muscle	Yes	Yes†	Yes†	Yes†					
Sliced fish	Yes	Yes‡	Yes	Yes					
Water	Yes	Yes‡	Yes	Yes					

*GBS, group B Streptococcus; ST, sequence type. †These parameters were tested on whole fish/fish part samples collected at only fresh produce markets and sashimi suppliers; whole fish samples collected from ports were excluded.

‡All food safety indicators were tested except for Aeromonas spp.

									Antimicrobial
	No. isolates			Sold as RTE		Sold as			drug resistance
Sample ID	analyzed	Sample description	Type of sample	dish	Sampling site†	freshwater fish	Serotype	ST	gene‡
SGEHI2015-63§	2	Fish parts (slab) sold as snakehead-toman	Muscle	No	Wet market stall	Yes	la	7	_
SGEHI2015-IV87	2	Sliced fish sold as salmon	Muscle	Yes (sashimi)	Restaurant/snack bar	No	la	7	_
SGEHI2015-IV170	2	Sliced fish sold as taï (tilapia)¶	Muscle	Yes (sashimi)	Restaurant/snack bar	No	la	7	_
SGEHI2015-49§	1	Fish parts (slab) sold as snakehead-toman	Surface swab	No	Wet market stall	Yes	la	23	tetM
SGEHI2015-50§	1	Fish parts (slab) sold as snakehead-toman	Muscle	No	Wet market stall	Yes	la	23	tetM
SGEHI2015-243	10	Whole fish sold asmackerel	Surface swab	No	Fish stall (port)	No	la	23	lsaC, tetM
SGEHI2015-IV89	5	Sliced fish sold as tuna	Muscle	Yes (sashimi)	Restaurant/snack	No	la	103	tetO#
SGEHI2015-IV211	2	Sliced fish sold as tuna	Muscle	Yes (sashimi)	Restaurant/snack	No	la	103	-
SGEHI2015-IV232	2	Sliced fish sold as salmon	Muscle	Yes (<i>lo-hei</i>)	Restaurant/snack	No	la	103	-
SGEHI2015-77§	1	Fish part (slab) sold as	Muscle	No	Wet market stall	Yes	II	28	tetM
SGEHI2015-II47	1	Sliced fish sold as grass	Muscle	Yes (yusheng)	Food stall	Yes	II	652	tetL, tetM
LG01	1	Whole fish sold as Asian	Organs	No	Port	Yes	Ш	283	_
LG02	1	Whole fish sold as Asian	Muscle	No	Port	Yes	Ш	283	_
LG03	1	Whole fish sold as Asian	Surface swab	No	Port	Yes	Ш	283	-
LG04	1	Whole fish sold as Asian	Surface swab	No	Port	Yes	Ш	283	-
LG05	1	Whole fish sold as Asian bighead carp	Surface swab	No	Port	Yes	Ш	283	-
LG06	1	Whole fish sold as Asian	Organs	No	Port	Yes	Ш	283	-
LG07	1	Water for holding live	NA	NA	Port	NA	Ш	283	-
LG08	1	Water for holding live	NA	NA	Port	NA	Ш	283	-
LG09	1	Water for holding live freshwater fish	NA	NA	Port	NA	Ш	283	_
LG10	1	Water for holding live freshwater fish	NA	NA	Port	NA	Ш	283	-
LG11	1	Water for holding live freshwater fish	NA	NA	Port	NA	Ш	283	-

Technical Appendix Table 3. Characteristics of group B Streptococcus isolates from fish and water samples, Singapore, 2015–2016*

	No. isolates			Sold as RTE		Sold as			Antimicrobial drug resistance
Sample ID	analyzed	Sample description	Type of sample	dish	Sampling site†	freshwater fish	Serotype	ST	gene‡
LG12	1	Water for holding live freshwater fish	NA	NA	Port	NA			-
SGEHI2015- NWC941§	3	Sliced fish sold as grass carp	Muscle	Yes (yusheng)	Food stall	Yes	III	283	-
SGEHI2015-107§	1	Whole live fish sold as snakehead-haruan	Muscle	No	Wet market stall	Yes	Ш	283	-
SGEHI2015-95§	1	Fish part (slab) sold as silver carp	Muscle	No	Supermarket	Yes	Ш	283	-
SGEHI2015-22§	2	Fish part (tail) sold as	Muscle	No	Fish stall (port)	Yes	Ш	283	-
SGEHI2015-25§	1	Fish part (head) sold as Asian bighead carp	Muscle	No	Fish stall (port)	Yes	Ш	283	-
SGEHI2015-51§	1	Fish part (head) sold as Asian bighead carp	Surface swab	No	Supermarket	Yes	Ш	283	-
SGEHI2015-31§	4	Fish part (slab) sold as	Muscle	No	Fish stall (port)	Yes	Ш	283	-
SGEHI2015-57§	1	Whole fish sold as black	Muscle	No	Wet market stall	Yes	Ш	283	-
SGEHI2015-101§	1	Whole fish sold as black	Muscle	No	Wet market stall	Yes	Ш	283	-
SGEHI2015-113§	1	Whole fish sold as black	Muscle	No	Supermarket	Yes	Ш	283	tetM
SGEHI2015-114§	1	Whole fish sold as black	Organs	No	Supermarket	Yes	Ш	283	tetM
SGEHI2015-29§	2	Whole fish sold as red	Organs	No	Fish stall (port)	Yes	Ш	283	-
SGEHI2015–59§	1	Whole fish sold as black	Surface swab	No	Wet market stall	Yes	Ш	283	-
SGEHI2015-60§	1	Whole fish sold as black	Muscle	No	Wet market stall	Yes	la	24	tetM
SGEHI2015-II33	1	Sliced fish sold as	Muscle	Yes (yusheng)	Food stall	Yes	Ш	335	tetM
SGEHI2015-IV45	5	Sliced fish sold as taï (tilania)¶	Muscle	Yes (sashimi)	Restaurant/snack	No	Ш	651	ermB, tetL
SGEHI2015-IV118	2	Sliced fish sold as salmon	Muscle	Yes (sashimi)	Restaurant/snack	No	Ш	861	lsaC, msrD, tetO
SGEHI2015-IV100	3	Sliced fish sold as salmon	Muscle	Yes (sashimi)	Restaurant/snack	No	V, la	1, 485	tetM
SGEHI2015-IV104	2	Sliced fish sold as salmon	Muscle	Yes (sashimi)	Restaurant/snack	No	V	1	tetM
SGEHI2015-II55	2	Sliced fish sold as wolf herring	Muscle	Yes (yusheng)	Food stall	No	V	1	tetM
SGEHI2015-II56	1	Sliced fish sold as Asian	Muscle	No	Food stall	Yes	V	1	tetM
SGEHI2015-IV74	1	Sliced fish sold as salmon	Muscle	Yes (sashimi)	Restaurant/snack bar	No	VII	1	tetM

									Antimicrobial
	No. isolates			Sold as RTE		Sold as			drug resistance
Sample ID	analyzed	Sample description	Type of sample	dish	Sampling site†	freshwater fish	Serotype	ST	gene‡
SGEHI2015-IV72	2	Sliced fish sold as salmon	Muscle	Yes (sashimi)	Restaurant/snack bar	No	VII	1	tetM, tetO
SGEHI2015-IV132	2	Sliced fish sold as salmon	Muscle	Yes (sashimi)	Restaurant/snack bar	No	V	24	tetM
SGEHI2015-IV227	1	Sliced fish sold as wolf herring	Muscle	Yes (<i>lo-hei</i>)	Restaurant/snack bar	No	VI	167	tetM

*NA, not applicable; RTE, ready-to-eat; ST, sequence type; *lo-hei*, Lunar New Year festive dish consisting of raw fish slices served with raw vegetables and condiments; *yusheng*, Chinese style sliced raw fish dish served separately with porridge.

†Food stalls refer to stalls housed within larger eating establishments that include hawker centers, coffee shops, and eating houses. Supermarkets refer to fresh produce sections of supermarkets and exclude sashimi and sushi counters of supermarkets.

[‡]-, no genes encoding resistance from the ARGannot database (including those acting on aminoglycosides, β-lactams, colistin, fosfomycin, fluoroquinolones, glycopeptides, macrolides, lincosamide, streptogramines, phenicols, rifampin, sulfonamides, tetracyclines, and trimethoprim) were detected by SRST2 from the genome sequencing data.

Sisolates also described in a study (7) on the analysis of clinical, epidemiologic, and bacterial sequencing data obtained during investigation of group B Streptococcus infections.

Fish identified as tilapia (Oreochromis spp.) by using the DNA barcoding method (29).

#Gene encoding resistance to tetracycline (tetO) was detected in 3 of the 5 isolates.



Technical Appendix Figure 1. Group B *Streptococcus* infections caused by handling and consumption of raw fish, Singapore, 2015–2016. SPC of fish muscle before and after rinsing with water. Dashed horizontal line indicates regulatory limit of Singapore for SPCs in ready-to-eat foods (<5 log₁₀ CFU/g) (*30*). Top and bottom of boxes in plots indicate 25th and 75th percentiles, horizontal lines indicate medians, and error bars indicate minimum and maximum values. Open circles indicate outliers. NS, not significant (p>0.05); SPC, standard plate count; SG, Singapore government.



Technical Appendix Figure 2. Group B *Streptococcus* infections caused by handling and consumption of raw fish, Singapore, 2015–2016. Global phylogenetic tree (neighbor-joining tree) with SNPs relative to the SG-M1 reference genome for 1,369 GBS strains. Red circles indicate strains isolated from fish and fish tank water samples sequenced during this study. Blue circles indicate complete publicly available genome sequences. All other strains are sequences from the GenBank Short Read Archive. Vertical black bars indicate CCs (*31*) and STs. Black arrows indicate several relatively rare fish strains from this study that fall outside the major CCs and have relatively few neighboring publicly available sequences. Seventeen of these strains (just below CC17) are placed among a set of strains that are more likely not to have come from humans (fish and camel strains). CC, clonal complex; GBS, group B *Streptococcus*; SNPs, single-nucleotide polymorphisms; ST, sequence type.