

Pneumonia-Specific *Escherichia coli* with Distinct Phylogenetic and Virulence Profiles, France, 2012–2014

Béatrice La Combe, Olivier Clermont, Jonathan Messika, Matthieu Eveillard, Achille Kouatchet, Sigismond Lasocki, Stéphane Corvec, Karim Lakhal, Typhaine Billard-Pomares, Romain Fernandes, Laurence Armand-Lefevre, Sandra Bourdon, Jean Reignier, Vincent Fihman, Nicolas de Prost, Julien Bador, Julien Goret, Frederic Wallet, Erick Denamur, Jean-Damien Ricard, on behalf of the COLOCOLI group¹

In a prospective, nationwide study in France of *Escherichia coli* responsible for pneumonia in patients receiving mechanical ventilation, we determined *E. coli* antimicrobial susceptibility, phylotype, O-type, and virulence factor gene content. We compared 260 isolates with those of 2 published collections containing commensal and bacteremia isolates. The preponderant phylogenetic group was B2 (59.6%), and the predominant sequence type complex (STc) was STc73. STc127 and STc141 were overrepresented and STc95 underrepresented in pneumonia isolates compared with bacteremia isolates. Pneumonia isolates carried higher proportions of virulence genes *sfa/foc*, *papGIII*, *hlyC*, *cnf1*, and *iroN* compared with bacteremia isolates. Virulence factor gene content and antimicrobial drug resistance were higher in pneumonia than in commensal isolates. Genomic

and phylogenetic characteristics of *E. coli* pneumonia isolates from critically ill patients indicate that they belong to the extraintestinal pathogenic *E. coli* pathovar but have distinguishable lung-specific traits.

Nosocomial infections remain a major threat for patients and a burden on healthcare institutions, hampering the public health economy. In intensive care units (ICUs), the most common life-threatening nosocomial infection is ventilator-associated pneumonia; the attributable mortality rate is ≈13%, partly because of increased durations of mechanical ventilation and ICU stays (1,2), all of which generate considerable additional costs (3).

Until the early 2000s, *Enterobacteriaceae* were not considered as major pathogens responsible for ventilator-assisted pneumonia (2); as such, pathophysiological studies focused mainly on *Pseudomonas aeruginosa*, *Staphylococcus aureus*, and *Acinetobacter baumannii*. However, recent data have consistently shown that *Enterobacteriaceae* are now frequent etiologic agents of ventilator-assisted pneumonia, more frequent than *P. aeruginosa* and *S. aureus* (4–6). According to the World Health Organization, *Enterobacteriaceae*, including *Escherichia coli*, are among the critical priority antibiotic-resistant bacteria (7). Therefore, to optimize patient management, in-depth epidemiologic knowledge of the phenotypic and genotypic characteristics of these bacteria is warranted. Although most *E. coli* responsible for symptomatic extraintestinal infections (8), including the urinary tract (9), bloodstream (10,11), cerebral spinal fluid (12,13), and peritoneum (14), have been extensively studied, less is known about *E. coli* strains responsible for pneumonia (15), especially in the context of highly virulent and resistant clones such as sequence type (ST) 69 and ST131 (16).

Author affiliations: Infection, Antimicrobiens, Modélisation, Évolution, Paris, France (B. La Combe, O. Clermont, J. Messika, T. Billard-Pomares, R. Fernandes, L. Armand-Lefevre, E. Denamur, J.-D. Ricard); Université Paris Diderot, Paris (B. La Combe, O. Clermont, J. Messika, T. Billard-Pomares, R. Fernandes, L. Armand-Lefevre, E. Denamur, J.-D. Ricard); Hôpital Louis Mourier, Colombes, France (B. La Combe, J. Messika, T. Billard-Pomares, R. Fernandes, J.-D. Ricard); Centre Hospitalier Universitaire, Angers, France (M. Eveillard, A. Kouatchet, S. Lasocki); Centre Hospitalier Universitaire, Nantes, France (S. Corvec, J. Reignier); Hôpital Laënnec, Nantes (K. Lakhal); Hôpital Bichat, AP-HP, Paris (L. Armand-Lefevre, E. Denamur); Centre Hospitalier Départemental Vendée, La Roche-sur-Yon, France (S. Bourdon); Hôpital Henri Mondor, AP-HP, Créteil, France (V. Fihman, N. de Prost); Centre Hospitalier Universitaire Bocage Central, Dijon, France (J. Bador); Centre Hospitalier Universitaire Pellegrin, Bordeaux, France (J. Goret); Centre Hospitalier Régional Universitaire, Lille, France (F. Wallet)

DOI: <https://doi.org/10.3201/eid2504.180944>

¹Group members are listed at the end of this article.

Our previous monocentric prospective study in France found a predominance of B2 phylogenetic group (66%) among *E. coli* pneumonia isolates, one third of which belonging to ST complex (STc) 127 (15). To obtain further insights in the physiopathology of ventilator-assisted pneumonia, we conducted a multicentric prospective epidemiologic study of genomic and phylogenetic characteristics of *E. coli* strains responsible for pneumonia across France.

Materials and Methods

Patients and *E. coli* Isolates

We conducted this prospective study in 14 ICUs throughout France, in collaboration with their hospital laboratories. We selected these ICUs to guarantee appropriate geographic coverage of metropolitan France. The same geographic selection was also applied to the ICUs in the Paris area. The ethics committee of the French Intensive Care Society approved the study (SRLF-CE 12-388), which was registered at ClinicalTrials.gov (NCT03303937). Formal consent was not required because of the observational, noninterventional study design (no change in practices, and all procedures already routinely performed). Patients or family/relatives were informed of the nature of the study and its purpose and objectives and had the option of declining participation.

During a 38-month period (2012–2014), any *E. coli* isolate responsible for pneumonia in a mechanically ventilated patient was collected, regardless of the method of sampling (quantitative cultures of tracheal suctioning, bronchoalveolar lavage, or protected telescoping catheter). Demographic and clinical data for the patient were recorded. We defined ventilator-assisted pneumonia as pneumonia occurring ≥ 48 hours after initiation of invasive ventilation, hospital-acquired pneumonia as pneumonia occurring ≥ 48 hours after hospital admission but within the first 48 hours of invasive ventilation, and community-acquired pneumonia as pneumonia that occurred either before or within the first 48 hours of hospitalization (17). Each *E. coli* isolate was stored at -80°C in brain-heart infusion broth containing glycerol 20% (the COLOCOLI collection).

E. coli Phylotyping, O-Typing, and Virulence Factor Gene Content Determination

We used quadruplex PCR to determine the *E. coli* phylogenetic group (A, B1, B2, C, D, E, F) or belonging to clade I (18). Among the strains, we determined the B1 clonal complex 87 (CC87) (Institut Pasteur MLST schema nomenclature corresponding to the ST58 and 155 in the Achtman schema [19]), the 10 main B2 subgroups (20), and the clonal group A (clonal group A) from the D phylogroup (21). The exhaustive correspondence between this typing

approach and STc membership according to the currently used Achtman MLST schema is available in (22).

We used PCR to determine the most anticipated serotypes in isolates from patients with extraintestinal infections (Table 1) (23). Multiplex PCR was used to detect genes encoding for 11 frequently encountered extraintestinal virulence factors (*sfa/foc*, *papC*, *papGII*, *papGIII*, *fyuA*, *iroN*, *aer*, *traT*, *neuC*, *hlyC*, and *cnfI*), which belong to the main classes of virulence factors (adhesins, toxins, iron acquisition systems, and protectins) (11). For each isolate, we calculated the virulence score, which was defined by the number of virulence factors present among the 11 tested.

Antimicrobial Resistance Phenotypes

We determined the antimicrobial susceptibility of each isolate by using the disk-diffusion method according to the French Society of Microbiology (<https://www.sfm-microbiologie.org>) (Table 2). Resistance score was defined as the sum of inactive in vitro antimicrobial agents for each isolate. A score of 1 indicates resistant; 0.5, intermediary; and 0, sensitive. A higher score indicates a more resistant isolate. Detection of gene sequences coding for the TEM, SHV, and CTX-M enzymes was performed by PCR with genomic DNA. The oligonucleotide primer sets specific for the β -lactamase genes used in the PCR assays have been published (24,25).

Characteristics of Other *E. coli* Strain Collections

To learn more about *E. coli* pneumonia strains, we compared the isolates from the COLOCOLI collection with

Table 1. Characterization of the main *Escherichia coli* B2 phylogroup clones, by combination of subgroup and O-type, among patients with extraintestinal infections, France, 2012–2014*

Subgroup and O-type	B2 clones, no. (%), n = 155
I-O25b	15 (62.5)
I-O6a	2 (8.3)
I-O16	3 (12.5)
II-O22	2 (5)
II-O2b	4 (10)
II-O6a	26 (65)
III-O6a	16 (100)
IV-O2b	19 (95)
VI-O4	13 (92.9)
VII-O18	2 (66.6)
VII-O75	1 (33.3)
IX-O1	7 (50)
IX-O18	3 (21.4)
IX-O2a	3 (21.4)

*Data are presented as no. (%) of each clone within the corresponding subgroup. Roman numerals correspond to the B2 subgroup. The correspondence with the Achtman multilocus sequence type schema is as follows: subgroup I, sequence type complex (STc) 131; subgroup II, STc73; subgroup III, STc127; subgroup IV, STc141; subgroup VI, STc12; subgroup VII, STc14; subgroup IX, STc95 (22). The most anticipated serotypes in extraintestinal infections were searched by PCR: O1, O2a, O2b, O4, O6, O7, O12, O15, O16, O17, O18, O22, O25a, O25b, O45a, O75, O78.

Table 2. Resistance and virulence traits of the 260 *Escherichia coli* isolates responsible for pneumonia in patients receiving mechanical ventilation, according to phylogenetic group, France, 2012–2014*

Trait	Phylogenetic group			Phylogenetic group				
	B2, n = 155	Non-B2, n = 105	p value	A, n = 22	B1, n = 26	C, n = 20	D, n = 25	F, n = 11
<i>iroN</i>	130 (83.9)	52 (49.5)	<0.0001	10 (45.4)	13 (50)	16 (80)	6 (24)	6 (54.5)
<i>sfa/foc</i>	109 (70.3)	0	<0.0001	0	0	0	0	0
<i>neuC</i>	41 (26.4)	0	<0.0001	0	0	0	0	0
<i>fyuA</i>	152 (98.1)	51 (48.6)	<0.0001	7 (31.8)	9 (34.6)	16 (80)	10 (40)	8 (72.7)
<i>hlyC</i>	98 (63.2)	2 (1.9)	<0.0001	0	2 (7.7)	0	0	0
<i>cnf1</i>	91 (58.7)	1 (1)	<0.0001	0	1 (3.8)	0	0	0
<i>aer</i>	65 (41.9)	67 (63.8)	0.0006	14 (63.6)	15 (57.7)	14 (70)	15 (60)	8 (72.7)
<i>papC</i>	100 (64.5)	27 (25.7)	<0.0001	7 (31.8)	3 (11.5)	8 (40)	8 (32)	1 (9.1)
<i>papGI</i>	27 (17.4)	4 (3.8)	0.0007	0	0	0	4 (16)	0
<i>papGIII</i>	64 (41.3)	0	<0.0001	0	0	0	0	0
<i>traT</i>	70 (45.2)	82 (78.1)	<0.0001	15 (68.2)	20 (76.9)	17 (85)	20 (80)	9 (81.8)
Virulence score, median (IQR)†	7 (5–7)	3 (2–4)	<0.0001	2.5 (2–4)	3 (1–4)	4 (3–5)	3 (2–3)	3 (2.5–3)
Antimicrobial resistance								
Amoxicillin	75 (48.4)	83 (79)	<0.0001	19 (86.4)	19 (73)	16 (80)	19 (76)	10 (90.9)
Amoxicillin/clavulanic acid	66 (42.6)	68 (64.8)	0.0006	18 (81.8)	13 (50)	15 (75)	14 (56)	8 (72.7)
Piperacillin/tazobactam	21 (13.5)	27 (25.7)	0.02	6 (27.3)	6 (23.1)	7 (35)	3 (12)	5 (45.4)
Cefotaxime	11 (7.1)	17 (16.2)	0.02	4 (18.2)	4 (15.4)	2 (10)	3 (12)	4 (36.4)
Ceftazidime	12 (7.7)	17 (16.2)	0.04	4 (18.2)	4 (15.4)	2 (10)	3 (12)	4 (36.4)
Imipenem	0	1 (1)	0.4	0	1 (3.8)	0	0	0
Gentamicin	4 (2.6)	10 (9.5)	0.02	3 (13.6)	1 (3.8)	2 (10)	1 (4)	3 (27.3)
Amikacin	3 (1.9)	1 (1)	0.6	0	0	0	0	1 (9.1)
Ofloxacin	15 (9.7)	28 (26.7)	0.0005	7 (31.8)	6 (23.1)	6 (30)	2 (8)	6 (54.5)
Ciprofloxacin	13 (8.4)	24 (22.9)	0.002	7 (31.8)	5 (19.2)	6 (30)	2 (8)	4 (36.4)
Resistance score, median (IQR)‡	1.5 (0–4)	4.5 (2.5–7)	<0.0001	5 (3.5–8)	4 (1–6)	4.5 (3.5–7.5)	4 (1.5–5.5)	7.5 (5–9)
ESBL phenotype	10 (6.4)	12 (11.4)	0.2	4 (18.2)	1 (3.8)	1 (4)	2 (8)	3 (27.3)
WT phenotype	84 (54.2)	24 (22.9)	<0.0001	4 (18.2)	7 (26.9)	6 (24)	7 (28)	1 (9.1)

*Values are no. (%) except as indicated. ESBL, extended-spectrum beta-lactamase; IQR, interquartile range; WT, wild type (susceptible to all tested antimicrobials).

†Virulence score was calculated, defined by the number of present virulence factors among the 11 tested.

‡Resistance score was defined as the sum of inactive in vitro antimicrobial agents for each isolate. Tested antimicrobials were gentamicin, amikacin, minocycline, nalidixic acid, ofloxacin, ciprofloxacin, fosfomycin, furans, trimethoprim, amoxicillin, amoxicillin-clavulanic acid, ticarcillin, piperacillin, piperacillin-tazobactam, imipenem, cefotaxime, and ceftazidime. A score of 1 was attributed for a resistant, 0.5 for an intermediary, and 0 for a sensitive isolate; a higher score thus indicated a more resistant isolate. For each antimicrobial, resistance is defined by the sum of resistant or intermediary isolates.

those of 2 published collections, originating from the Paris area in France. In 2010, a total of 280 *E. coli* strains were isolated from fecal samples from community adults and can be considered as commensal strains (COLIVILLE collection) (26). In 2005, a total of 373 *E. coli* strains were isolated from the blood of 373 in-patients in 14 hospitals during the course of bacteremia (COLIBAFI collection) (27). Of note, 20.6% of isolates from patients with bacteremia were nosocomial and 57% were of urinary origin. Among patients with bacteremia, the portal of entry was pulmonary for <2% (most patients were not in an ICU).

For these strains, we determined the phylogroup/subgroup membership, the presence of the 11 virulence factors cited above, and the susceptibility to 6 antimicrobial drugs (amoxicillin, amoxicillin/clavulanic acid, cefotaxime, amikacin, ofloxacin, and ceftazidime). For all strains, we calculated a virulence score.

Statistical Analyses

For our analyses we used GraphPad Prism7 software (<https://www.graphpad.com>). For quantitative variables, results are presented as the median and range, and for

categorical variables, as frequency and proportion. Variables were compared according to whether they were nosocomial or community isolates and whether they were of phylogenetic group B2 or not B2. As we compared the virulence factor gene content of the 3 collections, we also compared the proportion of phylogenetic groups and subgroups and of resistant strains. We used the Student *t* test to compare continuous variables and the Fisher exact test to compare categorical variables. Because of the observational design of the study and its exploratory aim, we did not adjust for multiple testing (28). We considered $p < 0.05$ to be significant.

Results

Host Characteristics

During the study period, we collected 260 *E. coli* isolates from 243 patients with a median age of 64 years (interquartile range 52–73 years) (Table 3). Of these isolates, 117 were responsible for ventilator-assisted pneumonia, 61 for hospital-acquired pneumonia, and 82 for community-acquired pneumonia. The main reasons for ICU admission

Table 3. Demographics and clinical characteristics of 243 pneumonia patients requiring mechanical ventilation, from whom *Escherichia coli* was isolated, France, 2012–2014*

Characteristic	Value
Age, y, median (IQR)	64 (52–73)
Sex	
M	183 (75.3)
F	60 (24.7)
SAPS II at admission, median (IQR)	57 (42–69)
Comorbid conditions	
Chronic alcohol consumption	56 (23)
Diabetes mellitus	45 (18.5)
Neoplastic disease	43 (17.7)
Immunosuppression†	77 (31.7)
Cirrhosis	12 (4.9)
Chronic kidney disease	18 (7.4)
Dialysis	5 (2.1)
Chronic respiratory disease	33 (13.6)
Chronic heart failure	43 (17.7)
Reason for ICU admission	
Acute respiratory failure	61 (25.1)
Coma	48 (19.8)
Septic shock	44 (18.1)
Cardiac arrest	28 (11.5)
Cardiogenic shock	14 (5.8)
Polytrauma	22 (9.1)
Postoperative care	8 (3.3)
Hemorrhagic shock	5 (2.1)
Exposure to antimicrobial drug therapy in previous 3 mo	98 (40.3)
Amoxicillin	6 (2.5)
Amoxicillin/clavulanic acid	38 (15.6)
Third-generation cephalosporin	19 (7.8)
Aminoglycosides	29 (11.9)
Piperacillin/tazobactam	24 (9.9)
Quinolone	10 (4.1)
Carbapenem	11 (4.5)
Polymicrobial sampling	57 (23.5)
ICU length of stay, d (IQR)	17 (7–33)
Hospital length of stay, d (IQR)	24 (10–45)
Death	
While in ICU	90 (37)
While in hospital	99 (40.7)
Associated with <i>E. coli</i>	27 (11.1)

*Values are no. (%) except as indicated. ICU, intensive care unit; IQR, interquartile range; SAPS II, Simplified Acute Physiology Score II.

†Defined by ≥ 1 immunosuppression factor among neoplastic disease, hematologic malignancy, HIV infection, immunosuppressive therapy, corticosteroids therapy.

were acute respiratory failure ($n = 61$, 25.1%), coma ($n = 48$, 19.8%), and septic shock ($n = 44$, 18.1%). A total of 98 (40.3%) patients had received antimicrobial drugs in the previous 3 months.

E. coli Characteristics

E. coli alone was isolated from 76.5% of the respiratory samples, whereas 23.5% of the samples were polymicrobial. The monomicrobial and polymicrobial samples did not differ in terms of phylogroup, virulence factor content, or antimicrobial resistance. We compiled classifications of the different phylogenetic groups/subgroups (Table 4) and details about their community or nosocomial status (Appendix Table 1, <https://wwwnc.cdc.gov/EID/>

article/25/4/18-0944-App1.pdf). The main phylogenetic groups were B2 ($n = 155$, 59.6%), B1 ($n = 26$, 10%), and D ($n = 25$, 9.6%). The most commonly identified lineages were STc73 (subgroup II, $n = 40$, 25.8% of B2 isolates), STc131 (subgroup I, $n = 24$, including 18 ST131), STc69 (clonal group A [29], $n = 20$), STc141 (subgroup IV, $n = 20$), and STc127 (subgroup III, $n = 16$). STc95 (subgroup IX) encompassed 14 strains (9% of B2 isolates). Community and nosocomial isolates did not differ in terms of phylogenetic group, except for C phylogroup isolates, which had a community predisposition (14.6% community vs. 4.5% nosocomial; $p = 0.01$).

We identified the O-type of 163 strains. We identified B2 phylogroup strains at the clonal level as having a combination of subgroup and O-type, as previously described (27) (Table 1). Among B2 strains, clones II-O6a were predominant, followed by IV-O2b, I-O25b (which belongs to ST131), III-O6a (which belongs to the highly virulent archetypal strain 536 [30]), and VI-O4.

Resistance and virulence traits of the 260 strains are detailed in Table 2 and Appendix Table 2. B2 phylogroup strains carried more virulence factor genes (virulence score 7 [5–7]) than non-B2 phylogroup strains (3 [2–4]; $p < 0.0001$). However, *traT* genes were significantly more present in non-B2 (78.1%) than B2 phylogroup (45.2%) isolates ($p < 0.0001$), as were *aer* genes (non-B2 63.8% and B2 41.9%; $p = 0.0006$). In nearly three quarters of strains, mainly B2 strains, we found *iroN* (70%) and *fyuA* (78.1%). Community and nosocomial isolates did not differ in terms of virulence and resistance scores. A total of 22 (8.5%) isolates were producers of extended-spectrum β -lactamases, including 13 CTX-M-1 group, 6 CTX-M-9 group, and 3 TEM. One isolate produced an OXA-48 carbapenemase, 28 (10.8%) isolates were resistant to cefotaxime, and 48 (18.5%) strains were resistant to piperacillin-tazobactam. B2 phylogroup strains were more sensitive to antimicrobials (resistance score 1.5 [0–4]) than were non-B2 phylogroup strains (resistance score 4.5 [2.5–7]; $p < 0.0001$).

General Comparisons of the COLOCOLI, COLIBAFI, and COLIVILLE Collections

When comparing the pneumonia *E. coli* isolates with those from the 2 other collections, we found strong differences (Table 5; Figure). The B2 phylogroup was overrepresented in pneumonia strains (59.6%) compared with commensal strains (32.1%; $p < 0.0001$) but not with bacteremia strains (52%; $p = 0.06$). Among B2 phylogroup pneumonia strains, subgroup III (STc127) was significantly overrepresented (10.3%) compared with the 2 other collections (bacteremia 4.1%, $p = 0.03$; commensal 2.2%, $p = 0.02$). The proportion of subgroup IV (STc141) isolates was significantly higher among pneumonia strains (12.9%) than bacteremia strains (2.6%; $p = 0.0002$), whereas subgroup IX (STc95) isolates

(9%) were underrepresented compared with the bacteremia B2 phylogroup strains (29.4%; $p < 0.0001$). Within the D phylogenetic group, the proportions of clonal group A pneumonia isolates (80%) were greater than those of commensal isolates (40%; $p = 0.001$).

Virulence scores of pneumonia isolates (5 [3–7]) were significantly higher than those of commensal isolates (3 [1–5]; $p < 0.0001$) but not different from those of the bacteremia isolates (4 [2–7]; $p = 0.3$). However, some adhesins (*sfa/foc*, *papGIII*), some toxins (*hlyC*, *cnfI*), and *iroN* were significantly overrepresented in pneumonia *E. coli* strains compared with strains in the 2 other collections.

Pneumonia *E. coli* isolates were more resistant than commensal isolates to all tested antimicrobial drugs except amikacin. Pneumonia isolates and bacteremia isolates did not differ in terms of antimicrobial drug susceptibility, except for resistance to cefotaxime (pneumonia isolate resistance 10.8%) and ceftaxime (pneumonia isolate resistance 11.5%) compared with bacteremia isolates (5.1%; $p = 0.009$ for cefotaxime, $p = 0.004$ for ceftaxime). Of note, when the 260 pneumonia isolates were compared with the 220 bacteremia strains of urinary tract origin, they were still distinguishable in terms of phylogroups/subgroups and virulence factor content (Appendix).

Discussion

This prospective nationwide study provides data on *E. coli* pneumonia isolates in critically ill patients. With regard to the characteristics of *E. coli* pneumonia isolates, we found the following: 1) a preponderance of phylogenetic group B2 (59.6%); 2) a predominant STc73 (subgroup II) lineage and threatening emergence of ST131 (within subgroup I), STc69 (clonal group A), and STc127 (subgroup III), along with STc141 (subgroup IV); 3) a much lower proportion of STc95 (subgroup IX) in B2 pneumonia than in bacteremia isolates; 4) a specific virulence factor gene content in pneumonia versus bacteremia strains. Taken together, these epidemiologic, phylogenetic, genotypic, and experimental data argue for inclusion of *E. coli* pneumonia isolates in the extraintestinal pathogenic *E. coli* (ExPEC) pathovar but with distinguishable lung-specific traits.

In 2010, Croxen and Finlay reviewed the molecular mechanisms of *E. coli* pathogenicity (31). Among ExPEC, numerous pathovars were listed, including uropathogenic and neonatal meningitis pathogenic *E. coli*, but none concerned the lungs. Pneumonia was not even cited as a possible disease caused by *E. coli*.

The situation is now clearly different. In 2012, Hamet et al. reported that *Enterobacteriaceae* accounted for a quarter of the 323 episodes of ventilator-assisted pneumonia occurring in their ICU (6). Our group showed that over a 5-year analysis of ventilator-assisted pneumonia episodes, finding *Enterobacteriaceae* as the responsible

Table 4. Phylogenetic groups/subgroups of *Escherichia coli* isolated from 260 pneumonia patients requiring mechanical ventilation, France, 2012–2014*

Phylogroup	No. (%)
A	22 (8.5)
B1	26 (10)
CC87†	11 (42.3)
Non-CC87	15 (57.7)
B2	155 (59.6)
I ST131†	18 (11.6)
I non-ST131	6 (3.9)
II	40 (25.8)
III	16 (10.3)
IV	20 (12.9)
V	1 (0.6)
VI	14 (9)
VII	3 (1.9)
IX	14 (9)
Unassigned	23 (14.8)
C	20 (7.7)
D	25 (9.6)
CGA†	20 (80)
Non-CGA	5 (20)
F	11 (4.2)
Clade I	1 (0.4)

*Nosocomial isolates encompass ventilator-associated pneumonia and hospital-acquired pneumonia isolates. CC87, clonal complex 87 (sequence type complex [STc] 155) (19); CGA, clonal group A (STc69) (21); ST131, sequence type 131.
The correspondence with the Achtman MLST schema is as follows: subgroup I, STc131; subgroup II, STc73; subgroup III, STc127; subgroup IV, STc141; subgroup V, STc144; subgroup VI, STc12; subgroup VII, STc14; subgroup IX, STc95. No isolate belonged to subgroup VIII (STc452), X (STc372), or E phylogroup (22).
†Proportions of subgroups are reported as fractions of the respective phylogroups.

pathogen increased significantly (5). In an international multicenter study, Kollef et al. also confirmed that *Enterobacteriaceae* were the leading pathogens of ventilator-assisted pneumonia in the ICU (4). Among them, *E. coli* is a major threat, recently highlighted by the World Health Organization, because of its ever-increasing resistance to antimicrobial drugs (7).

ExPEC are characterized by pathogenic virulence factor genes coding for various combinations of adhesins, toxins, iron-acquisition systems, capsule production, and toxins that enable them to cause disease once outside the host gut reservoir (32). ExPEC virulence factors are encoded on the bacterial chromosome, where they are usually located within pathogenicity-associated islands (PAIs) or plasmids. Most ExPEC isolates belong to the B2 phylogroup and, to a lesser extent, the D phylogroup. More in-depth analysis of ExPEC strains has enabled characterization of particular STs of ExPEC isolates including ST131, ST73, and ST127 (33). Our most striking finding was the specificity of pneumonia *E. coli* strains, compared with bacteremia ones, within the ExPEC family (Table 5). First, although B2 phylogroup strains are preponderant in ExPEC (15,34), we found a trend toward an even greater proportion of B2 isolates among pneumonia isolates (59.6%) than among bacteremia isolates, whatever the origin (52%; $p = 0.06$)

Table 5. Proportion of phylogenetic groups, B2 subgroups, D CGA, and B1 CC87 among pneumonia, bacteremia, or commensal isolates*

Phylogenetic characteristics	Pneumonia isolates	Bacteremia isolates	Commensal isolates
A	–	+	++
B1	+	–	+
CC87†	+++	+++	+
B2	++++	++++	+++
I	+	+	+
II	++	++	++
III	+	–	–
IV	+	–	+
IX	–	++	+
C	–	–	–
D	–	+	+
CGA*	++++	++++	+++
E	–	–	–
F	–	–	–

*CGA, clonal group A (sequence type complex [STc] 69) (21); CC87, clonal complex 87 (STc155) (19); –, <10%; +, 10%–20%; ++, 20%–30%; +++ 30–50%; +++++, >50% of isolates. Correspondence with the Achtman multilocus sequence typing schema is subgroup I = STc131; subgroup II = STc73; subgroup III = STc127; subgroup IV = STc141; and subgroup IX = STc95 (22).

†Proportions of subgroups are calculated within each respective phylogroup.

(Table 3; Figure). This finding is in the range of what is observed in urosepsis isolates (62%) (11) but a little less than in neonatal meningitis strains (68%) (13). Then, among B2 phylogroup isolates (Table 4), if the predominance of subgroup II (STc73) was expected, we found high proportions of specific phylogenetic group B2 clones among other lineages, which could be worrisome (Table 1). The ST131 O25b:H4 clone represented 9.7% of the B2 phylogroup isolates, 53.3% of them producing an extended-spectrum β -lactamase. Subgroup IV (STc141) was significantly more present among pneumonia strains than among bacteremia strains. Challenging the hypothesis of a commensal character, with a low level of human invasiveness (27), this finding indicates that subgroup IV isolates may have a high affinity for the respiratory tract. Last, we must highlight the greater proportion of subgroup III among pneumonia isolates (namely STc127) compared with bacteremia and commensal isolates, in agreement with previous findings from our group (15). Contrary to isolates from the other 2 collections, pneumonia isolates were composed of fewer IX subgroup strains (STc95). Whereas subgroup IX is usually well represented among commensal and other pathogenic *E. coli* strains (26,27,35), these data suggest that the respiratory tract is less suitable than other tissues for subgroup IX implantation, a finding in agreement with our previous report in which subgroup IX was not represented (15).

Among D phylogroup strains, the higher prevalence of clonal group A (STc69) in pneumonia isolates than in bacteremia and commensal isolates was unexpected. Among 571 D phylogroup *E. coli* responsible for extraintestinal infection, Johnson et al. reported only 144 clonal group A *E. coli* (25.2%) (29). The multidrug-resistant nature of these pathogens is of increasing concern (29,36).

Regarding virulence factor gene content, the literature suggests that PAIs involved in ExPEC causing pneumonia differ from those involved in urinary tract and blood-

stream infections (37,38). Using PAI deletion mutants in a rat model of pneumonia, Phillips-Houlbracq et al. related pneumonia pathogenicity to the presence of PAIs I and III (37). In our study (Figure 1), pneumonia isolates differed from bacteremia isolates because they significantly more often carried *sfa/foc*, *iroN* (both belonging to PAI III), *papGIII* and *cnfI* (belonging to PAI II), and *hlyC* (belonging to PAI I and II). The role of α -hemolysin in experimentally induced pneumonia in rats has been reported (15,39). Of note, the virulence factor content of the pneumonia strains still differs when we consider only the urosepsis isolates (Appendix). Although additional studies are required to confirm, these findings do suggest a coherent molecular trait for the isolates' lung specificity.

Consistent with data in the literature (15), we found that pneumonia B2 isolates were less resistant to antimicrobial drugs (B2 resistance score 1.5 [0–4] vs. non-B2 resistance score 4.5 [2.5–7]; $p < 0.0001$) but carried more virulence factor genes (B2 virulence score 7 [5–7] vs. non-B2 virulence score 3 [2–4]; $p < 0.0001$) (Table 2). However, contrary to an old belief, this trade-off does not mean that antimicrobial drug resistance decreases with increasing virulence (40). Indeed, virulence and antimicrobial-drug resistance were both higher in pneumonia isolates than in commensal isolates and the following were more highly represented: *sfa/foc*, *papC*, *papGIII*, *fyuA*, *iroN*, *hlyC*, and *cnfI*.

Our study should be interpreted within the context of its limitations. First, although our collection of *E. coli* pneumonia isolates is large, results regarding some subgroup analyses will require confirmation because of their sample size. Our choice of PCR rather than whole-genome characterization was governed by our wish to compare the 2 other collections with the same type of data. Despite these limitations, our analysis based on 14 centers representing France on a population and geographic level, its prospective design, and the comparison of our large number of isolates




Figure. Comparison of *Escherichia coli* isolates among pneumonia patients with extraintestinal infections, France, 2012–2014, with commensal *E. coli* (COLIVILLE collection) and *E. coli* bacteremia isolates (COLIBAFI collection). A) Proportions of *E. coli* phylogenetic groups and subgroups; B) B2 subgroups, D CGA, and B1 CC87; C) virulence factors; and D) antimicrobial drug resistance. Roman numerals correspond to the Achtman multilocus sequence typing schema is as follows: subgroup I, STc131; subgroup II, STc73; subgroup III, STc127; subgroup IV, STc141; subgroup VI, STc12; subgroup VII, STc14; subgroup IX, STc95 (22). Proportions of subgroups are reported as fractions of the respective phylogroups. Asterisks indicate a significant difference between respiratory isolates and strains responsible for bacteremia (COLIBAFI collection) or commensal strains (COLIVILLE collection): * $p < 0.05$; ** $p < 0.005$; *** $p < 0.0005$. AMC, amoxicillin/clavulanic acid; AMK, amikacin; AMX, amoxicillin; CTX, cefotaxime; FOX, cefoxitin; OFX, ofloxacin; STc, sequence type complex.

with those from recently published collections enable us to draw valid conclusions. We did not assess the functionality and expression of the encoded virulence factors in these isolates. Our team has consistently demonstrated these features in several murine models of infection (including pneumonia) and observed a strong correlation between the presence of these genes and death (15,34,37).

Our data raise the question of why certain clonal lineages were overrepresented in patients with respiratory tract infection. Patients acquire *E. coli* infection from their own digestive tract (41). This event implies an upward retrograde motion of the bacterial cells to reach the oropharynx and the lung parenchyma and suggests particular metabolic-adaptation and response-to-stress characteristics (e.g., to overcome the acidity of the stomach). We have previously shown that some *E. coli* strains are capable of high growth capacities in relation to metabolic pathways (42) while others are highly resistant to stress (43). For both studies, however, no link to specific clones could be

established. The specific organ tropism therefore more probably results from a combination of genetic background and virulence factors.

In summary, we identified emerging pneumonia-causing pathogenic *E. coli* whose main characteristics define them as ExPEC. Their specificities include a very strong proportion of B2 phylogroup isolates; a high proportion of subgroups II (STc73), I (STc131), IV (STc141), and III (STc127); and consequent proportions of clonal group A (STc69) isolates within the D phylogroup. Virulence factor gene content of pneumonia isolates also appeared to be singular compared with that of bacteremia isolates, among them urosepsis isolates. These epidemiologic data underline the specificity of pneumonia *E. coli* populations and may help with the design of more targeted therapies.

Acknowledgments

We are indebted to the physicians (microbiologists and intensivists) who took part in the COLOCOLI Study. We are

also grateful to the patients and their families, and we gratefully acknowledge Dominique Poignard and Marie-Claire Hipeaux for technical assistance and all the nurses and assistant nurses of the ICU for their help in obtaining the microbiological samples.

Collaborators for the COLOCOLI group: Catherine Branger, Luce Landraud, Alexandre Bleibtreu, Françoise Jauréguay (Infection, Antimicrobiens, Modélisation, Évolution, Paris, France); Pierre Asfar (Centre Hospitalier Universitaire, Angers, France); Didier Dreyfuss, Guilène Barnaud, Fatma Magdoud (Hôpital Louis Mourier, AP-HP, Colombes, France); Michel Wolff, Jean-François Timsit (Hôpital Bichat, AP-HP, Paris, France); Stéphanie Martin (Centre Hospitalier Départemental Vendée, La Roche-sur-Yon, France); Pierre-Emmanuel Charles (Centre Hospitalier Universitaire Bocage Central, Dijon, France); Alexandre Boyer (Centre Hospitalier Universitaire Pellegrin, Bordeaux, France); Emmanuelle Jaillette, Saad Nseir (Centre Hospitalier Régional Universitaire, Lille, France); Raymond Ruimy, Pierre-Eric Danin, Jean Dellamonica (Centre Hospitalier Universitaire, Nice, France); Julie Cremitter, Jean-Pierre Frat (Centre Hospitalier Universitaire, Poitiers, France); Christophe Clec'h (Hôpital Avicenne, AP-HP, Bobigny, France), Dominique Decré, Eric Maury (Hôpital Saint-Antoine, AP-HP, Paris, France).

This work was supported by a grant from the Fonds de Dotation Recherche en Santé Respiratoire/Fondation du Souffle to (B.L.C.). This work was partly supported by a grant from the Fondation pour la Recherche Médicale (to E.D., Équipe FRM 2016, grant no. DEQ20161136698).

About the Author

Dr. La Combe is an intensive care physician who works at the Infection, Antimicrobials, Modelling, Evolution laboratory of the French Institute for Medical Research in Paris. Her work focuses on the virulence and antimicrobial resistance of respiratory *E. coli* isolates and the pharmacodynamics and genetics of chlorhexidine resistance of oropharyngeal isolates in patients receiving mechanical ventilation.

References

- Melsen WG, Rovers MM, Groenwold RHH, Bergmans DCJJ, Camus C, Bauer TT, et al. Attributable mortality of ventilator-associated pneumonia: a meta-analysis of individual patient data from randomised prevention studies. *Lancet Infect Dis*. 2013;13:665–71. [http://dx.doi.org/10.1016/S1473-3099\(13\)70081-1](http://dx.doi.org/10.1016/S1473-3099(13)70081-1)
- Chastre J, Fagon J-Y. Ventilator-associated pneumonia. *Am J Respir Crit Care Med*. 2002;165:867–903. <http://dx.doi.org/10.1164/ajrcm.165.7.2105078>
- Russo TA, Johnson JR. Medical and economic impact of extraintestinal infections due to *Escherichia coli*: focus on an increasingly important endemic problem. *Microbes Infect*. 2003;5:449–56. [http://dx.doi.org/10.1016/S1286-4579\(03\)00049-2](http://dx.doi.org/10.1016/S1286-4579(03)00049-2)
- Kollef MH, Ricard J-D, Roux D, Francois B, Ischaki E, Rozgonyi Z, et al. A randomized trial of the amikacin fosfomicin inhalation system for the adjunctive therapy of gram-negative ventilator-associated pneumonia: IASIS Trial. *Chest*. 2017;151:1239–46. <http://dx.doi.org/10.1016/j.chest.2016.11.026>
- Fihman V, Messika J, Hajage D, Tournier V, Gaudry S, Magdoud F, et al. Five-year trends for ventilator-associated pneumonia: correlation between microbiological findings and antimicrobial drug consumption. *Int J Antimicrob Agents*. 2015;46:518–25. <http://dx.doi.org/10.1016/j.ijantimicag.2015.07.010>
- Hamet M, Pavon A, Dalle F, Pechinot A, Prin S, Quenot J-P, et al. *Candida* spp. airway colonization could promote antibiotic-resistant bacteria selection in patients with suspected ventilator-associated pneumonia. *Intensive Care Med*. 2012;38:1272–9. <http://dx.doi.org/10.1007/s00134-012-2584-2>
- World Health Organization. Global priority list of antibiotic-resistant bacteria to guide research, discovery, and development of new antibiotics [cited 2017 Sep 29]. <http://www.who.int/medicines/publications/global-priority-list-antibiotic-resistant-bacteria/en/>
- Johnson JR, Russo TA. Extraintestinal pathogenic *Escherichia coli*: “the other bad *E. coli*.” *J Lab Clin Med*. 2002;139:155–62. <http://dx.doi.org/10.1067/mlc.2002.121550>
- Johnson JR, Kuskowski MA, Gajewski A, Soto S, Horcajada JP, Jimenez de Anta MT, et al. Extended virulence genotypes and phylogenetic background of *Escherichia coli* isolates from patients with cystitis, pyelonephritis, or prostatitis. *J Infect Dis*. 2005;191:46–50. <http://dx.doi.org/10.1086/426450>
- Johnson JR, Kuskowski MA, O'Bryan TT, Maslow JN. Epidemiological correlates of virulence genotype and phylogenetic background among *Escherichia coli* blood isolates from adults with diverse-source bacteremia. *J Infect Dis*. 2002;185:1439–47. <http://dx.doi.org/10.1086/340506>
- Lefort A, Panhard X, Clermont O, Woerther P-L, Branger C, Mentré F, et al.; COLIBAFI Group. Host factors and portal of entry outweigh bacterial determinants to predict the severity of *Escherichia coli* bacteremia. *J Clin Microbiol*. 2011;49:777–83. <http://dx.doi.org/10.1128/JCM.01902-10>
- Bidet P, Mahjoub-Messai F, Blanco J, Dehem M, Aujard Y, et al. Combined multilocus sequence typing and O serogrouping distinguishes *Escherichia coli* subtypes associated with infant urosepsis and/or meningitis. *J Infect Dis*. 2007;196:297–303. <http://dx.doi.org/10.1086/518897>
- Bingen E, Picard B, Brahim N, Mathy S, Desjardins P, Elion J, et al. Phylogenetic analysis of *Escherichia coli* strains causing neonatal meningitis suggests horizontal gene transfer from a predominant pool of highly virulent B2 group strains. *J Infect Dis*. 1998;177:642–50. <http://dx.doi.org/10.1086/514217>
- Bert F, Johnson JR, Ouattara B, Leflon-Guibout V, Johnston B, Marcon E, et al. Genetic diversity and virulence profiles of *Escherichia coli* isolates causing spontaneous bacterial peritonitis and bacteremia in patients with cirrhosis. *J Clin Microbiol*. 2010;48:2709–14. <http://dx.doi.org/10.1128/JCM.00516-10>
- Messika J, Magdoud F, Clermont O, Margetis D, Gaudry S, Roux D, et al. Pathophysiology of *Escherichia coli* ventilator-associated pneumonia: implication of highly virulent extraintestinal pathogenic strains. *Intensive Care Med*. 2012;38:2007–16. <http://dx.doi.org/10.1007/s00134-012-2699-5>
- Blanco J, Mora A, Mamani R, López C, Blanco M, Dahbi G, et al. National survey of *Escherichia coli* causing extraintestinal infections reveals the spread of drug-resistant clonal groups O25b:H4-B2-ST131, O15:H1-D-ST393 and CGA-D-ST69 with high virulence gene content in Spain. *J Antimicrob Chemother*. 2011;66:2011–21. <http://dx.doi.org/10.1093/jac/dkr235>
- American Thoracic Society, Infectious Diseases Society of America. Guidelines for the management of adults with

- hospital-acquired, ventilator-associated, and healthcare-associated pneumonia. *Am J Respir Crit Care Med.* 2005;171:388–416. <http://dx.doi.org/10.1164/rccm.200405-644ST>
18. Clermont O, Christenson JK, Denamur E, Gordon DM. The Clermont *Escherichia coli* phylo-typing method revisited: improvement of specificity and detection of new phylo-groups. *Environ Microbiol Rep.* 2013;5:58–65. <http://dx.doi.org/10.1111/1758-2229.12019>
 19. Skurnik D, Clermont O, Guillard T, Launay A, Danilchanka O, Pons S, et al. Emergence of antimicrobial-resistant *Escherichia coli* of animal origin spreading in humans. *Mol Biol Evol.* 2016;33:898–914. <http://dx.doi.org/10.1093/molbev/msv280>
 20. Clermont O, Christenson JK, Daubié A-S, Gordon DM, Denamur E. Development of an allele-specific PCR for *Escherichia coli* B2 sub-typing, a rapid and easy to perform substitute of multilocus sequence typing. *J Microbiol Methods.* 2014;101:24–7. <http://dx.doi.org/10.1016/j.mimet.2014.03.008>
 21. Johnson JR, Owens K, Manges AR, Riley LW. Rapid and specific detection of *Escherichia coli* clonal group A by gene-specific PCR. *J Clin Microbiol.* 2004;42:2618–22. <http://dx.doi.org/10.1128/JCM.42.6.2618-2622.2004>
 22. Clermont O, Gordon D, Denamur E. Guide to the various phylogenetic classification schemes for *Escherichia coli* and the correspondence among schemes. *Microbiology.* 2015;161:980–8. <http://dx.doi.org/10.1099/mic.0.000063>
 23. Clermont O, Johnson JR, Menard M, Denamur E. Determination of *Escherichia coli* O types by allele-specific polymerase chain reaction: application to the O types involved in human septicemia. *Diagn Microbiol Infect Dis.* 2007;57:129–36. <http://dx.doi.org/10.1016/j.diagmicrobio.2006.08.007>
 24. Eckert C, Gautier V, Saladin-Allard M, Hidri N, Verdet C, Ould-Hocine Z, et al. Dissemination of CTX-M-type beta-lactamases among clinical isolates of *Enterobacteriaceae* in Paris, France. *Antimicrob Agents Chemother.* 2004;48:1249–55. <http://dx.doi.org/10.1128/AAC.48.4.1249-1255.2004>
 25. Rasheed JK, Jay C, Metchock B, Berkowitz F, Weigel L, Crellin J, et al. Evolution of extended-spectrum beta-lactam resistance (SHV-8) in a strain of *Escherichia coli* during multiple episodes of bacteremia. *Antimicrob Agents Chemother.* 1997;41:647–53. <http://dx.doi.org/10.1128/AAC.41.3.647>
 26. Massot M, Daubié A-S, Clermont O, Jauréguy F, Couffignal C, Dahbi G, et al.; The COLIVILLE Group. Phylogenetic, virulence and antibiotic resistance characteristics of commensal strain populations of *Escherichia coli* from community subjects in the Paris area in 2010 and evolution over 30 years. *Microbiology.* 2016;162:642–50. <http://dx.doi.org/10.1099/mic.0.000242>
 27. Clermont O, Couffignal C, Blanco J, Mentré F, Picard B, Denamur E; COLIVILLE and COLIBAFI groups. Two levels of specialization in bacteraemic *Escherichia coli* strains revealed by their comparison with commensal strains. *Epidemiol Infect.* 2017;145:872–82. <http://dx.doi.org/10.1017/S0950268816003010>
 28. Bender R, Lange S. Adjusting for multiple testing—when and how? *J Clin Epidemiol.* 2001;54:343–9. [http://dx.doi.org/10.1016/S0895-4356\(00\)00314-0](http://dx.doi.org/10.1016/S0895-4356(00)00314-0)
 29. Johnson JR, Menard ME, Lauderdale T-L, Kosmidis C, Gordon D, Collignon P, et al.; Trans-Global Initiative for Antimicrobial Resistance Analysis Investigators. Global distribution and epidemiologic associations of *Escherichia coli* clonal group A, 1998–2007. *Emerg Infect Dis.* 2011;17:2001–9. <http://dx.doi.org/10.3201/eid1711.110488>
 30. Brzuszkiewicz E, Brüggemann H, Liesegang H, Emmerth M, Olschläger T, Nagy G, et al. How to become a uropathogen: comparative genomic analysis of extraintestinal pathogenic *Escherichia coli* strains. *Proc Natl Acad Sci U S A.* 2006;103:12879–84. <http://dx.doi.org/10.1073/pnas.0603038103>
 31. Croxen MA, Finlay BB. Molecular mechanisms of *Escherichia coli* pathogenicity. *Nat Rev Microbiol.* 2010;8:26–38. <http://dx.doi.org/10.1038/nrmicro2265>
 32. Russo TA, Johnson JR. Proposal for a new inclusive designation for extraintestinal pathogenic isolates of *Escherichia coli*: ExPEC. *J Infect Dis.* 2000;181:1753–4. <http://dx.doi.org/10.1086/315418>
 33. Dale AP, Woodford N. Extra-intestinal pathogenic *Escherichia coli* (ExPEC): disease, carriage and clones. *J Infect.* 2015;71:615–26. <http://dx.doi.org/10.1016/j.jinf.2015.09.009>
 34. Picard B, Garcia JS, Gouriou S, Duriez P, Brahimi N, Bingen E, et al. The link between phylogeny and virulence in *Escherichia coli* extraintestinal infection. *Infect Immun.* 1999;67:546–53.
 35. Ciesielczuk H, Jenkins C, Chattaway M, Doumith M, Hope R, Woodford N, et al. Trends in ExPEC serogroups in the UK and their significance. *Eur J Clin Microbiol Infect Dis.* 2016;35:1661–6. <http://dx.doi.org/10.1007/s10096-016-2707-8>
 36. Kallonen T, Brodrick HJ, Harris SR, Corander J, Brown NM, Martin V, et al. Systematic longitudinal survey of invasive *Escherichia coli* in England demonstrates a stable population structure only transiently disturbed by the emergence of ST131. *Genome Res.* 2017;27:1437–49. <http://dx.doi.org/10.1101/gr.216606.116>
 37. Phillips-Houlbracq M, Ricard J-D, Fourcier A, Yoder-Himes D, Gaudry S, Bex J, et al. Pathophysiology of *Escherichia coli* pneumonia: respective contribution of pathogenicity islands to virulence. *Int J Med Microbiol.* 2018;308:290–6. <http://dx.doi.org/10.1016/j.ijmm.2018.01.003>
 38. Turret J, Diard M, Garry L, Matic I, Denamur E. Effects of single and multiple pathogenicity island deletions on uropathogenic *Escherichia coli* strain 536 intrinsic extra-intestinal virulence. *Int J Med Microbiol.* 2010;300:435–9. <http://dx.doi.org/10.1016/j.ijmm.2010.04.013>
 39. Russo TA, Davidson BA, Genagon SA, Warholc NM, Macdonald U, Pawlicki PD, et al. *E. coli* virulence factor hemolysin induces neutrophil apoptosis and necrosis/lysis in vitro and necrosis/lysis and lung injury in a rat pneumonia model. *Am J Physiol Lung Cell Mol Physiol.* 2005;289:L207–16. <http://dx.doi.org/10.1152/ajplung.00482.2004>
 40. Roux D, Danilchanka O, Guillard T, Cattoir V, Aschard H, Fu Y, et al. Fitness cost of antibiotic susceptibility during bacterial infection. *Sci Transl Med.* 2015;7:297ra114. <http://dx.doi.org/10.1126/scitranslmed.aab1621>
 41. Garrouste-Orgeas M, Chevret S, Arlet G, Marie O, Rouveau M, Popoff N, et al. Oropharyngeal or gastric colonization and nosocomial pneumonia in adult intensive care unit patients. A prospective study based on genomic DNA analysis. *Am J Respir Crit Care Med.* 1997;156:1647–55. <http://dx.doi.org/10.1164/ajrccm.156.5.96-04076>
 42. Sabarly V, Bouvet O, Glodt J, Clermont O, Skurnik D, Diancourt L, et al. The decoupling between genetic structure and metabolic phenotypes in *Escherichia coli* leads to continuous phenotypic diversity. *J Evol Biol.* 2011;24:1559–71. <http://dx.doi.org/10.1111/j.1420-9101.2011.02287.x>
 43. Bleibtreu A, Gros P-A, Laouénan C, Clermont O, Le Nagard H, Picard B, et al. Fitness, stress resistance, and extraintestinal virulence in *Escherichia coli*. *Infect Immun.* 2013;81:2733–42. <http://dx.doi.org/10.1128/IAI.01329-12>

Address for correspondence: Jean-Damien Ricard, Service de Réanimation Médicale, Hôpital Louis Mourier, 92700 Colombes, France; email: jean-damien.ricard@aphp.fr

Pneumonia-Specific *Escherichia coli* Isolates with Distinct Phylogenetic and Virulence Profiles, France, 2012–2014

Appendix

Results

Appendix Table 1. *E. coli* phylogenetic groups belonging according to the community or nosocomial status*

Phylogroups	All (n = 260)	Community isolates (n = 82)	Nosocomial isolates (n = 178)	p value
A	22 (8.5)	5 (6.1)	17 (9.6)	0.5
B1	26 (10)	4 (4.8)	22 (12.4)	0.08
CC87†	11 (42.3)	2 (50)	9 (40.9)	1
Non-CC87	15 (57.7)	2 (50)	13 (59.1)	
B2	155 (59.6)	54 (65.9)	101 (56.7)	0.2
I ST131†	18 (11.6)	7 (13)	11 (10.9)	0.8
I non-ST131	6 (3.9)	3 (5.6)	3 (3)	0.4
II	40 (25.8)	14 (26)	26 (25.7)	1
III	16 (10.3)	5 (9.3)	11 (10.9)	1
IV	20 (12.9)	4 (7.4)	16 (15.8)	0.2
V	1 (0.6)	1 (1.9)	0	0.3
VI	14 (9)	6 (11.1)	8 (7.9)	0.6
VII	3 (1.9)	1 (1.9)	2 (2)	1
IX	14 (9)	4 (7.4)	10 (9.9)	0.8
Unassigned	23 (14.8)	9 (16.7)	14 (13.9)	0.6
C	20 (7.7)	12 (14.6)	8 (4.5)	0.01
D	25 (9.6)	4 (4.9)	21 (11.8)	0.1
CGA†	20 (80)	3 (75)	17 (81)	1
Non-CGA	5 (20)	1 (25)	4 (19)	
F	11 (4.2)	3 (3.7)	8 (4.5)	1
Clade I	1 (0.4)	0	1 (0.6)	1

*Abbreviations: ST131, sequence type 131; CGA, clonal group A (STc69) [1] CC87, Clonal complex 87 (STc155) [18]. Nosocomial isolates encompass VAP and HAP isolates. Data are presented as n (%). The correspondence with the Achtman MLST schema is as follow: subgroup I = STc131; subgroup II = STc73; subgroup III = STc127; subgroup IV = STc141; subgroup V = STc144; subgroup VI = STc12; subgroup VII = STc14; subgroup IX = STc95. No isolate belonged to subgroup VIII (STc452), X (STc372) or to E phylogroup [2].

†Proportions of subgroups are reported as fractions of the respective phylogroups.

Appendix Table 2. Resistance and virulence traits of the 260 *E. coli* isolates, according to the community or nosocomial status*

Trait	All (n = 260)	Community isolates (n = 82)	Nosocomial isolates (n = 178)	p value
<i>iroN</i>	182 (70)	60 (73.2)	122 (68.5)	0.5
<i>sfa/foc</i>	109 (41.9)	38 (46.3)	71 (39.9)	0.3
<i>neuC</i>	41 (15.2)	12 (14.6)	29 (16.3)	0.9
<i>fyuA</i>	203 (78.1)	70 (85.4)	133 (74.7)	0.06
<i>hlyC</i>	100 (38.5)	35 (42.7)	65 (36.5)	0.4
<i>cnf1</i>	92 (35.4)	32 (39)	60 (33.7)	0.4
<i>aer</i>	132 (50.8)	38 (46.3)	94 (52.8)	0.3
<i>papC</i>	127 (48.8)	35 (42.7)	92 (51.7)	0.2
<i>papGII</i>	31 (11.9)	8 (9.8)	23 (12.9)	0.5
<i>papGIII</i>	64 (24.6)	19 (23.2)	45 (25.3)	0.8
<i>traT</i>	152 (58.5)	47 (57.3)	105 (59)	0.9
Virulence score	5 [3-7]	5 [3-7]	5 [3-7]	0.8
Resistance score	3.5 [0-5.5]	3 [0-5]	3.5 [0-5.5]	0.3
ESBL phenotype	22 (8.5)	4 (4.9)	18 (10.1)	0.2
WT phenotype	108 (41.5)	38 (46.3)	70 (39.3)	0.3

*Abbreviations: ESBL, extended-spectrum beta-lactamase; WT phenotype, Wild type phenotype (susceptibility to all tested antibiotics). Nosocomial isolates encompass VAP and HAP isolates. Data are presented as n (%) or median [interquartile range]. For each antibiotic, antimicrobial resistance is defined by the sum of resistant or intermediary isolates.

Comparison of phylogroups, subgroups, and virulence factor gene content of pneumonia isolates vs bacteraemia isolates of urinary origin

A former study from our group showed, among strains responsible for bacteremia, differences in the bacteremia strains of urinary tract origin with the presence of specific clones, whereas the bacteraemia strains of digestive origin resembled more the commensal strains (3).

Hence, we sought to compare *E. coli* pneumonia isolates and bacteraemia isolates of urinary origin (Appendix Table 3). Similarities lay in the phylogroup as well as in the subgroup distribution with two noticeable exceptions: subgroup IV was clearly over-represented in pneumonia isolates whereas this was the other way round for subgroup IX.

Differences were found in the distribution of virulence factor genes more than in the overall virulence score (5 [3-7] for the pneumonia isolates vs 4 [2-4] for the bacteremia isolates of urinary origin). Noticeable differences were related to a greater number of pneumonia isolates harbouring *sfa/foc*, *papGIII* and *cnf1*, and a greater number of bacteremia isolates from urinary origin harbouring *papC*, *papGII*.

Appendix Table 3. Comparison of phylogroups, subgroups, and virulence factor gene content of pneumonia isolates vs bacteraemia isolates of urinary origin*

Variable	Pneumonia isolates (n = 260)	Bacteremia isolates of urinary origin (n = 220)	p value
Phylogroups			
A	8.5	5.5	0.2
B1	10	5	0.06
B2	59.6	63.2	0.5
C	7.7	7.3	1
D	9.6	16.4	0.03
E	0	0	1
F	4.2	2.7	0.5
Subgroupst			
I	15.5	7.9	0.3
II	25.8	27.3	0.8
III	10.3	4.5	0.07
IV	12.9	1.4	0.0001
IX	9	34.5	<0.0001
CGA	80	69.4	0.4
CC87	42.3	27.3	0.5
Virulence factor genes			
<i>sfa/foc</i>	41.9	28.6	0.003
<i>papC</i>	48.8	67.7	<0.0001
<i>papGII</i>	11.9	59.1	<0.0001
<i>papGIII</i>	24.6	8.6	<0.0001
<i>fyuA</i>	78.1	88.2	0.004
<i>iroN</i>	70	62.3	0.08
<i>aer</i>	50.8	70	<0.0001
<i>traT</i>	58.5	70	0.01
<i>neuC</i>	15.2	25.5	0.009
<i>hlyC</i>	38.5	31.8	0.2
<i>cnf1</i>	35.4	20	0.0002

*The correspondence with the Achtman MLST schema is as follow: subgroup I = STc131; subgroup II = STc73; subgroup III = STc127; subgroup IV = STc141; subgroup V = STc144; subgroup VI = STc12; subgroup VII = STc14; subgroup IX = STc95. No isolate belonged to subgroup VIII (STc452), X (STc372) or to E phylogroup [2].

†Proportions of subgroups are reported as fractions of the respective phylogroups. Data are presented as n (%). Electronic supplemental material to "Emerging pneumonia-specific *Escherichia coli* isolates with distinct phylogenetic and virulence profiles: a nationwide study"

Results

Appendix Table 4. Phylogenetic groups, subgroups, serotypes, ESBL phenotype and virulence factor gene content of each isolate of the COLOCOLI collection*

Isolate ID	ESBL phenotype	Phylogenetic group	B2 subgroup	D subgroup	B1												
					subgroup	Serotype	<i>neuC</i>	<i>iroN</i>	<i>sfa</i>	<i>fyuA</i>	<i>hly</i>	<i>cnf1</i>	<i>aer</i>	<i>papC</i>	<i>papGII</i>	<i>papGIII</i>	<i>traT</i>
AN01	0	B2	III	non-D	non-B1	O6	0	1	1	1	1	1	0	1	0	1	1
AN02	0	B1	non-B2	non-D	CC87	ND	0	1	0	1	0	0	1	1	0	0	1
AN03	1	F	non-B2	non-D	non-B1	ND	0	1	0	1	0	0	1	0	0	0	1
AN04	0	B2	IV	non-D	non-B1	O2b	1	1	1	1	1	1	1	1	0	1	1
AN05	0	B2	IX	non-D	non-B1	O18	1	1	1	1	0	0	1	0	0	0	1
AN06	0	B2	UA	non-D	non-B1	O75	0	1	1	1	1	1	0	1	0	1	0
AN10	0	B2	III	non-D	non-B1	O6	0	0	0	1	0	0	0	1	0	1	0
AN12	0	B2	II	non-D	non-B1	O2b	0	0	0	1	1	1	1	1	1	0	1
AN13	0	D	non-B2	CGA	non-B1	O17	0	1	0	0	0	0	1	1	1	0	1
AN15	0	B1	non-B2	non-D	CC87	ND	0	1	0	0	0	0	1	0	0	0	1
AN16	0	B2	IX	non-D	non-B1	O2a	1	0	0	1	0	0	0	1	1	0	1
AN17	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	1	1	0	0	0	0
AN18	0	C	non-B2	non-D	non-B1	O78	0	1	0	1	0	0	1	0	0	0	1
AN19	0	C	non-B2	non-D	non-B1	ND	0	1	0	1	0	0	1	1	0	0	1
AN20	0	A	non-B2	non-D	non-B1	ND	0	0	0	1	0	0	0	0	0	0	1
AN21	0	B2	UA	non-D	non-B1	ND	0	0	0	0	0	0	0	1	0	0	0
AN22	0	B1	non-B2	non-D	no CC87	ND	0	0	0	0	0	0	0	0	0	0	1
AN24	0	A1	non-B2	non-D	non-B1	ND	0	1	0	1	0	0	1	0	0	0	1
AN25	0	C	non-B2	non-D	non-B1	ND	0	1	0	1	0	0	1	0	0	0	1
AN28	0	B2	IX	non-D	non-B1	O1	1	1	0	1	0	0	1	1	1	0	1
AN30	0	D	non-B2	CGA	non-B1	O25b	0	0	0	1	0	0	0	0	0	0	1
AN33	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	0	0	0	0	0	0
AN34	0	B2	UA	non-D	non-B1	O18	0	1	0	1	1	0	1	1	1	0	1
AN36	1	B2	I	non-D	non-B1	O25b	0	0	0	1	0	0	1	1	0	0	1
AN37	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	1	0	1	0	1	0
AN39	0	B2	IX	non-D	non-B1	O2a	1	1	0	1	0	0	1	1	1	0	1
AN41	0	B2	VI	non-D	non-B1	O4	0	1	1	1	1	1	1	1	1	0	0
AN43	0	D	non-B2	no CGA	non-B1	ND	0	1	0	0	0	0	0	0	0	0	1
AN44	0	C	non-B2	non-D	non-B1	ND	0	1	0	1	0	0	0	0	0	0	1
AN45	0	B2	UA	non-D	non-B1	O75	0	1	1	1	1	1	0	1	0	1	0
AN46	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	1	0	0	0	0	0
AN47	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	1	0	1	0	1	0
AN48	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	1	0	1	0	1	0
AN49	0	C	non-B2	non-D	non-B1	ND	0	1	0	0	0	0	0	0	0	0	1
AVC02	1	B2	I	non-D	non-B1	O25b	0	0	0	1	0	0	1	0	0	0	0
AVC03	1	F	non-B2	non-D	non-B1	ND	0	1	0	1	0	0	1	0	0	0	1
BCH811	0	B2	IV	non-D	non-B1	O2b	0	1	1	1	1	1	0	1	0	1	0
BCH818	0	B2	VII	non-D	non-B1	O18	0	1	1	1	0	0	0	1	1	0	0
BCH826	0	F	non-B2	non-D	non-B1	ND	0	0	0	1	0	0	0	0	0	0	1
BCH830	0	B2	VII	non-D	non-B1	O18	0	1	1	1	1	0	0	1	1	0	0
BCH838	0	F	non-B2	non-D	non-B1	ND	0	1	0	1	0	0	0	0	0	0	1
BCH841	0	F	non-B2	non-D	non-B1	ND	0	0	0	1	0	0	0	0	0	0	1

Isolate ID	ESBL phenotype	Phylogenetic group	B2 subgroup	D subgroup	B1		Serotype	<i>neuC</i>	<i>iroN</i>	<i>sfa</i>	<i>fyuA</i>	<i>hly</i>	<i>cnf1</i>	<i>aer</i>	<i>papC</i>	<i>papGII</i>	<i>papGIII</i>	<i>traT</i>
					subgroup													
BCH852	0	B2	IX	non-D	non-B1	O1	1	1	0	1	0	0	1	1	1	0	1	
BCH853	0	C	non-B2	non-D	non-B1	ND	0	1	0	1	0	0	0	0	0	0	1	
BCH855	0	D	non-B2	no CGA	non-B1	ND	0	0	0	0	0	0	0	0	0	0	0	
BCH857-15	0	B2	IX	non-D	non-B1	O1	1	1	0	1	0	0	1	1	1	0	1	
BCH857-72	0	B2	UA	non-D	non-B1	O75	0	1	1	1	1	1	0	1	0	1	0	
BCH858	0	F	non-B2	non-D	non-B1	ND	0	0	0	1	0	0	1	0	0	0	1	
BCH859	0	C	non-B2	non-D	non-B1	ND	0	0	0	1	0	0	1	1	0	0	1	
BCH860	0	B2	IX	non-D	non-B1	O1	1	1	0	1	0	0	1	1	1	0	1	
BCH861	0	B1	non-B2	non-D	CC87	ND	0	0	0	0	0	0	0	0	0	0	1	
BCH862	0	A1	non-B2	non-D	non-B1	ND	0	1	0	0	0	0	0	0	0	0	1	
BCH865	0	B2	IX	non-D	non-B1	O1	1	1	0	1	0	0	1	1	1	0	1	
BCH869	0	B2	I	non-D	non-B1	O6	0	1	0	1	0	0	0	0	0	0	1	
BCH876	0	B1	non-B2	non-D	no CC87	ND	0	0	0	0	0	0	0	0	0	0	0	
BCH877	1	B2	I	non-D	non-B1	O16	0	0	0	1	0	0	1	0	0	0	1	
BCH885	0	C	non-B2	non-D	non-B1	ND	0	1	0	1	0	0	1	1	0	0	1	
BCH887	0	D	non-B2	CGA	non-B1	ND	0	0	0	1	0	0	1	0	0	0	1	
BCH892	0	D	non-B2	CGA	non-B1	ND	0	0	0	1	0	0	1	0	0	0	1	
BCH898	0	C	non-B2	non-D	non-B1	ND	0	1	0	1	0	0	1	1	0	0	0	
BCH902	1	F	non-B2	non-D	non-B1	O2b	0	0	0	0	0	0	1	1	0	0	0	
BCH906	0	B2	VII	non-D	non-B1	O75	0	0	0	1	0	0	1	0	0	0	1	
BCH907	0	B2	II	non-D	non-B1	O2b	0	1	1	1	1	1	1	1	0	1	0	
BCH914	0	B2	I	non-D	non-B1	O16	0	0	0	1	0	0	1	0	0	0	1	
BCH916	0	B2	I	non-D	non-B1	O25b	0	0	0	1	0	0	1	0	0	0	0	
BCH921	0	B2	III	non-D	non-B1	O6	0	1	1	1	1	0	0	1	0	0	0	
BCH924	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	1	0	1	1	0	1	
BCH925	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	1	1	1	1	0	1	
BCH926-49	0	C	non-B2	non-D	non-B1	ND	0	1	0	1	0	0	1	0	0	0	1	
BCH926-55	0	B2	III	non-D	non-B1	O6	0	1	1	1	1	1	0	1	0	1	1	
BCH927	0	B2	III	non-D	non-B1	O6	0	1	1	1	1	1	0	1	0	0	0	
BCH933	0	B2	III	non-D	non-B1	O6	0	1	1	1	1	1	0	1	0	0	0	
BCH935	0	B2	VI	non-D	non-B1	O4	0	1	0	1	1	1	0	1	0	1	1	
BCH943	0	D	non-B2	CGA	non-B1	O17	0	1	0	0	0	0	1	1	1	0	1	
BCH957	0	B2	UA	non-D	non-B1	O75	0	1	1	1	1	1	0	1	0	1	0	
BCH964	0	B1	non-B2	non-D	no CC87	ND	0	0	0	1	0	0	0	0	0	0	1	
BDX01	1	A1	non-B2	non-D	non-B1	ND	0	0	0	0	0	0	1	1	0	0	1	
BDX02	0	B2	I	non-D	non-B1	O25b	0	1	0	1	0	0	1	1	0	1	1	
BDX03	0	D	non-B2	CGA	non-B1	O17	0	1	0	0	0	0	1	0	0	0	1	
BDX04	0	A0	non-B2	non-D	non-B1	ND	0	0	0	0	0	0	0	0	0	0	0	
BDX07	1	D	non-B2	CGA	non-B1	O25b	0	0	0	0	0	0	1	0	0	0	1	
BDX09	0	F	non-B2	non-D	non-B1	ND	0	0	0	1	0	0	1	0	0	0	1	
BDX10	0	A	non-B2	non-D	non-B1	ND	0	0	0	1	0	0	1	0	0	0	0	
BDX11	0	B2	VI	non-D	non-B1	O4	0	1	1	1	1	1	0	1	0	1	0	
BDX13	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	1	0	1	0	1	0	
BDX14	0	B2	II	non-D	non-B1	O4	0	1	1	1	0	0	0	0	0	0	1	
BDX15	0	C	non-B2	non-D	non-B1	ND	0	0	0	0	0	0	0	0	0	0	0	
BDX18	1	D	non-B2	CGA	non-B1	ND	0	0	0	1	0	0	0	0	0	0	1	
BDX19	1	B2	I	non-D	non-B1	O25b	0	0	0	1	1	1	1	1	1	0	1	
BDX20	0	C	non-B2	non-D	non-B1	ND	0	1	0	1	0	0	1	1	0	0	1	

Isolate ID	ESBL phenotype	Phylogenetic group	B2 subgroup	D subgroup	B1		Serotype	<i>neuC</i>	<i>iroN</i>	<i>sfa</i>	<i>fyuA</i>	<i>hly</i>	<i>cnf1</i>	<i>aer</i>	<i>papC</i>	<i>papGII</i>	<i>papGIII</i>	<i>traT</i>
					subgroup	no CC87												
BDX21	0	B1	non-B2	non-D	no CC87	ND	0	0	0	0	0	0	1	1	0	0	0	1
BDX22	1	B2	I	non-D	non-B1	O25b	0	0	0	1	0	0	1	0	0	0	0	0
BDX23	0	B2	II	non-D	non-B1	ND	0	1	1	1	0	0	1	0	0	0	0	1
BDX25	0	B2	UA	non-D	non-B1	O18	0	0	0	1	0	0	0	0	0	0	0	0
BDX26	0	D	non-B2	CGA	non-B1	O2b	0	0	0	1	0	0	1	0	0	0	0	1
BDX27	0	A1	non-B2	non-D	non-B1	ND	0	1	0	1	0	0	1	1	0	0	0	1
DIJ03	0	B2	II	non-D	non-B1	O2a	0	1	1	1	1	1	1	1	1	1	1	0
DIJ04	0	C	non-B2	non-D	non-B1	ND	0	1	0	1	0	0	1	0	0	0	0	1
DIJ06	0	B2	II	non-D	non-B1	O25a	0	1	1	1	1	1	0	0	0	0	0	0
DIJ07	0	D	non-B2	CGA	non-B1	O17	0	0	0	0	0	0	1	0	0	0	0	1
DIJ09	0	B2	II	non-D	non-B1	ND	0	1	1	1	0	0	0	0	0	0	0	1
DIJ10	0	B2	VI	non-D	non-B1	O4	0	1	1	1	1	1	0	1	0	1	1	0
DIJ11	0	B2	UA	non-D	non-B1	O6	0	1	1	1	1	1	0	1	0	1	1	1
DIJ12	0	B2	IV	non-D	non-B1	O2b	0	0	0	1	1	1	0	1	0	1	1	0
DIJ13	0	B2	I	non-D	non-B1	ND	1	1	0	1	0	0	0	0	0	0	0	0
DIJ14	0	B2	IV	non-D	non-B1	O2b	1	1	1	1	1	1	1	1	0	1	1	1
DIJ15	0	B2	IV	non-D	non-B1	O2b	1	1	1	1	0	0	0	0	0	0	0	0
DIJ16	0	B2	I	non-D	non-B1	O6	1	1	1	1	1	1	0	1	1	1	0	1
DIJ17	0	B2	VI	non-D	non-B1	O4	0	1	1	1	1	1	1	1	1	1	0	0
DIJ18	0	B1	non-B2	non-D	CC87	ND	0	1	0	1	0	0	1	0	0	0	0	1
H06	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	1	0	1	1	0	1	1
H08	0	B2	I	non-D	non-B1	O25b	0	0	0	1	0	0	1	0	0	0	0	1
H09	0	C	non-B2	non-D	non-B1	ND	0	0	0	1	0	0	0	0	0	0	0	0
H11	0	F	non-B2	non-D	non-B1	ND	0	1	0	0	0	0	1	0	0	0	0	1
H12	0	C	non-B2	non-D	non-B1	ND	0	1	0	1	0	0	1	0	0	0	0	1
H14	0	B1	non-B2	non-D	CC87	ND	0	0	0	1	1	0	1	0	0	0	0	1
H15	0	B2	II	non-D	non-B1	O6	0	1	1	1	0	0	1	0	0	0	0	1
H16	0	B2	UA	non-D	non-B1	O75	0	1	1	1	1	1	0	1	0	1	0	0
H17	0	B2	UA	non-D	non-B1	O16	0	0	0	0	0	0	0	0	0	0	0	1
H18	0	B2	I	non-D	non-B1	ND	1	1	0	1	0	0	0	0	0	0	0	1
H19	0	B2	III	non-D	non-B1	O6	0	1	1	1	1	1	0	1	0	1	1	0
H20	0	B2	VI	non-D	non-B1	O4	0	1	1	1	1	1	0	1	0	1	1	0
H21	0	B2	I	non-D	non-B1	ND	1	1	0	1	0	0	1	0	0	0	0	1
H22	0	A	non-B2	non-D	non-B1	ND	0	1	0	0	0	0	1	0	0	0	0	0
H23	0	B2	UA	non-D	non-B1	O75	0	1	1	0	1	1	0	1	0	1	1	0
H24	0	D	non-B2	CGA	non-B1	O25b	0	0	0	0	0	0	1	1	1	1	0	1
H25	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	0	1	1	1	1	0	1
H27	0	A	non-B2	non-D	non-B1	ND	0	0	0	1	0	0	1	1	0	0	0	1
H28	0	C	non-B2	non-D	non-B1	ND	0	1	0	1	0	0	1	1	0	0	0	1
H29	0	C	non-B2	non-D	non-B1	ND	0	0	0	0	0	0	0	0	0	0	0	1
H30	0	B2	III	non-D	non-B1	O6	0	1	1	1	1	1	0	1	0	1	1	1
H31	0	B2	I	non-D	non-B1	O25b	0	0	0	1	0	0	1	1	0	0	0	1
H32	1	C	non-B2	non-D	non-B1	O78	0	1	0	1	0	0	1	1	0	0	0	1
H33	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	1	1	1	0	0	0	0
H34	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	1	1	0	0	0	0	0
H35	0	B2	IV	non-D	non-B1	O2b	1	1	1	1	1	1	0	1	0	1	1	0
H37	0	A1	non-B2	non-D	non-B1	ND	0	0	0	1	0	0	1	0	0	0	0	1
LI01	0	B2	UA	non-D	non-B1	O18	0	0	0	1	0	0	0	0	0	0	0	0

Isolate ID	ESBL phenotype	Phylogenetic group	B2 subgroup	D subgroup	B1		Serotype	<i>neuC</i>	<i>iroN</i>	<i>sfa</i>	<i>fyuA</i>	<i>hly</i>	<i>cnf1</i>	<i>aer</i>	<i>papC</i>	<i>papGII</i>	<i>papGIII</i>	<i>traT</i>
					subgroup	no CC87												
LI02	0	F	non-B2	non-D	non-B1	ND	0	1	0	1	0	0	1	0	0	0	0	0
LI03	1	B2	I	non-D	non-B1	O25b	0	0	0	1	1	0	1	1	0	0	0	0
LI04	0	B2	II	non-D	non-B1	O7	1	1	1	1	0	0	0	0	0	0	0	0
LI05	0	B2	I	non-D	non-B1	O25b	0	1	0	1	0	0	1	1	0	0	0	1
LI06	0	B2	VI	non-D	non-B1	O4	0	1	1	1	1	1	0	1	0	1	0	0
LI07	0	B2	UA	non-D	non-B1	ND	0	1	1	1	1	1	1	0	0	0	0	1
LI08	0	B1	non-B2	non-D	no CC87	ND	0	1	0	0	0	0	1	0	0	0	0	1
LI09	0	B1	non-B2	non-D	no CC87	ND	0	1	0	0	0	0	1	0	0	0	0	1
LI10	1	B2	I	non-D	non-B1	O25b	0	0	0	1	0	0	1	1	1	0	0	1
LI11	0	B2	II	non-D	non-B1	O6	0	1	1	1	0	0	1	0	0	0	0	1
LI12	0	B2	III	non-D	non-B1	O6	0	1	1	1	1	1	0	1	0	1	0	0
LM01	0	B2	I	non-D	non-B1	ND	1	1	0	1	1	0	1	0	0	0	0	1
LM02	0	D	non-B2	CGA	non-B1	O17	0	0	0	1	0	0	0	0	0	0	0	1
LM03	1	A0	non-B2	non-D	non-B1	ND	0	0	0	0	0	0	1	0	0	0	0	1
LM08	0	D	non-B2	CGA	non-B1	O25b	0	0	0	1	0	0	0	1	0	0	0	1
LM09	0	B2	IX	non-D	non-B1	O2a	1	1	0	1	0	0	1	1	1	0	0	1
LM10	0	D	non-B2	CGA	non-B1	O15	0	0	0	0	0	0	1	1	0	0	0	1
LM12	0	B2	III	non-D	non-B1	O6	0	1	1	1	1	1	0	1	0	1	0	0
LM14	0	B2	IV	non-D	non-B1	O2b	1	1	1	1	1	1	0	1	0	1	0	0
LM17	0	B2	III	non-D	non-B1	O6	0	1	1	1	1	1	0	1	0	1	0	0
LM19	0	B1	non-B2	non-D	CC87	ND	0	1	0	1	0	0	1	0	0	0	0	1
LM31	0	B1	non-B2	non-D	CC87	ND	0	1	0	1	0	0	1	0	0	0	0	1
LM33	1	B2	I	non-D	non-B1	O25b	0	1	0	1	0	0	1	1	0	1	1	1
LM34	0	B2	I	non-D	non-B1	O25b	0	0	0	1	0	0	1	0	0	0	0	1
LM36	1	B2	I	non-D	non-B1	O25b	0	1	0	1	0	0	1	1	0	1	1	1
LM38	0	B2	VI	non-D	non-B1	O4	0	1	0	1	1	1	0	1	0	1	1	1
LM39	0	B1	non-B2	non-D	CC87	ND	0	1	0	1	0	0	1	0	0	0	0	1
LM43	0	C	non-B2	non-D	non-B1	ND	0	1	0	1	0	0	1	1	0	0	0	1
LM45	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	1	1	1	0	0	0	0
LM49	0	B2	UA	non-D	non-B1	O2b	0	1	1	1	1	1	0	1	0	1	0	0
LM51	0	B1	non-B2	non-D	CC87	ND	0	0	0	0	0	0	0	0	0	0	0	1
LM54	0	B2	III	non-D	non-B1	O6	0	1	1	1	1	1	0	1	0	1	0	0
LM57	0	A	non-B2	non-D	non-B1	ND	0	0	0	0	0	0	0	0	0	0	0	0
LM58	0	D	non-B2	CGA	non-B1	O7	0	0	0	0	0	0	0	0	0	0	0	0
LM61	0	B2	I	non-D	non-B1	O16	0	0	0	1	0	0	1	0	0	0	0	1
LM64	0	B2	UA	non-D	non-B1	ND	0	1	1	1	1	1	0	1	0	1	0	0
LM67	0	B2	III	non-D	non-B1	O6	0	1	1	1	1	1	0	1	0	1	0	0
LM68	0	B2	II	non-D	non-B1	O22	0	1	1	1	1	1	0	0	0	0	0	0
LM69	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	1	1	1	1	0	0	1
NAN02	1	A1	non-B2	non-D	non-B1	ND	0	1	0	1	0	0	1	1	0	0	0	1
NAN03	0	B2	IX	non-D	non-B1	O18	1	1	1	1	1	1	1	0	0	0	0	1
NAN04	0	B2	IV	non-D	non-B1	O2b	1	1	1	1	1	1	0	1	0	1	0	0
NAN05	1	B1	non-B2	non-D	no CC87	ND	0	1	0	0	0	0	1	0	0	0	0	1
NAN06	0	A1	non-B2	non-D	non-B1	ND	0	1	0	0	0	0	1	0	0	0	0	1
NAN08	0	D	non-B2	CGA	non-B1	ND	0	0	0	1	0	0	1	0	0	0	0	1
NAN09	0	B2	II	non-D	non-B1	O6	0	1	1	1	0	0	0	0	0	0	0	0
NAN11	0	D	non-B2	no CGA	non-B1	ND	0	0	0	1	0	0	0	1	0	0	0	0
NAN12	0	B2	IV	non-D	non-B1	ND	1	1	1	1	1	1	0	1	0	1	0	0

Isolate ID	ESBL phenotype	Phylogenetic group	B2 subgroup	D subgroup	B1		Serotype	<i>neuC</i>	<i>iroN</i>	<i>sfa</i>	<i>fyuA</i>	<i>hly</i>	<i>cnf1</i>	<i>aer</i>	<i>papC</i>	<i>papGII</i>	<i>papGIII</i>	<i>traT</i>
					subgroup	subgroup												
NAN13	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	1	1	1	0	0	0	0
NAN14	0	B2	VI	non-D	non-B1	ND	0	1	1	1	1	1	1	0	1	0	1	1
NAN15	0	B2	IV	non-D	non-B1	O2b	0	1	1	1	1	1	1	0	1	0	1	0
NAN16	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	1	1	1	0	0	0	0
NAN20	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	1	1	1	0	0	0	0
NAN21	0	B2	IX	non-D	non-B1	O1	1	1	0	1	0	0	1	1	1	1	0	1
NAN22	0	B2	IV	non-D	non-B1	O2b	1	1	1	1	0	0	1	0	0	0	0	0
NAN25	0	B2	IV	non-D	non-B1	O2b	1	1	1	1	1	1	1	0	1	0	1	0
NAN26	0	B2	IV	non-D	non-B1	O2b	1	1	1	1	1	1	0	1	0	1	1	0
NAN28	0	B1	non-B2	non-D	no CC87	ND	0	0	0	0	1	1	1	0	0	0	0	1
NAN29	0	A0	non-B2	non-D	non-B1	ND	0	1	0	0	0	0	0	0	0	0	0	1
NAN30	0	B2	I	non-D	non-B1	O25b	0	1	0	1	0	0	1	1	0	1	1	1
NAN31	0	D	non-B2	CGA	non-B1	O17	0	0	0	0	0	0	1	1	0	0	0	1
NAN32	0	B2	VI	non-D	non-B1	O4	0	1	1	1	1	1	1	0	1	0	1	0
NAN33	0	A1	non-B2	non-D	non-B1	ND	0	1	0	0	0	0	1	0	0	0	0	1
NAN34	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	1	1	0	0	0	0	0
NAN35	0	B2	VI	non-D	non-B1	O4	0	1	1	1	1	1	1	1	0	1	1	1
NAN36	0	B2	II	non-D	non-B1	O2b	0	1	1	1	0	0	0	0	0	0	0	0
NAN37	0	D	non-B2	CGA	non-B1	O17	0	0	0	1	0	0	1	1	1	1	0	1
NAN39	0	B2	UA	non-D	non-B1	O75	0	1	1	1	1	1	0	1	0	1	0	0
NAN40	0	A1	non-B2	non-D	non-B1	ND	0	0	0	0	0	0	0	0	0	0	0	0
NAN41	0	B2	VI	non-D	non-B1	O4	0	1	1	1	1	1	0	1	1	1	1	1
NAN42	0	B1	non-B2	non-D	no CC87	ND	0	0	0	0	0	0	0	0	0	0	0	1
NAN43	0	D	non-B2	CGA	non-B1	O17	0	1	0	0	0	0	1	0	0	0	0	1
NAN44	0	B2	UA	non-D	non-B1	O75	0	1	1	1	1	1	0	1	0	1	0	0
NAN45	0	B2	II	non-D	non-B1	ND	0	1	1	1	1	1	1	0	0	0	0	0
NAN46	0	B1	non-B2	non-D	CC87	ND	0	1	0	1	0	0	1	0	0	0	0	1
NAN47	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	1	1	0	0	0	0	1
NAN50	0	B1	non-B2	non-D	CC87	ND	0	1	0	1	0	0	1	1	0	0	0	1
NAN51	0	B2	II	non-D	non-B1	O2b	0	1	1	1	1	1	1	1	0	1	0	0
NIC01	1	B2	IV	non-D	non-B1	O2b	1	1	1	1	1	1	1	0	0	0	0	1
NIC03	0	B2	IX	non-D	non-B1	O1	1	0	0	1	0	0	0	1	1	1	0	1
NIC04	0	B2	IX	non-D	non-B1	O18	1	1	1	1	0	0	1	0	0	0	0	1
NIC05	0	B2	III	non-D	non-B1	O6	0	1	1	1	1	1	0	1	0	0	0	1
NIC06	0	A1	non-B2	non-D	non-B1	ND	0	1	0	0	0	0	1	1	0	0	0	1
NIC07	0	A1	non-B2	non-D	non-B1	ND	0	1	0	0	0	0	1	1	0	0	0	1
NIC08	1	B1	non-B2	non-D	no CC87	ND	0	1	0	0	0	0	1	0	0	0	0	1
NIC09	0	A1	non-B2	non-D	non-B1	ND	0	0	0	0	0	0	1	1	0	0	0	1
NIC11	0	B1	non-B2	non-D	no CC87	ND	0	0	0	0	0	0	0	0	0	0	0	0
NIC12	0	B2	IV	non-D	non-B1	O2b	1	1	1	1	1	1	0	1	0	1	1	1
PO01	0	B2	UA	non-D	non-B1	O7	1	1	1	1	0	0	0	0	0	0	0	1
PO02	0	F	non-B2	non-D	non-B1	ND	0	1	0	0	0	0	1	0	0	0	0	1
PO03	0	B1	non-B2	non-D	no CC87	ND	0	1	0	0	0	0	0	0	0	0	0	0
PO04	0	B2	II	non-D	non-B1	O25b	0	1	1	1	1	1	0	1	0	1	0	0
RY02	1	A1	non-B2	non-D	non-B1	O16	0	0	0	0	0	0	0	0	0	0	0	0
RY03	0	B1	non-B2	non-D	no CC87	ND	0	0	0	0	0	0	0	0	0	0	0	0
RY05	0	Clade	non-B2	non-D	non-B1	ND	0	0	0	0	0	0	0	0	0	0	0	0
RY06	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	1	1	0	0	0	0	0

Isolate ID	ESBL phenotype	Phylogenetic group	B2 subgroup	D subgroup	B1		Serotype	neuC	iroN	sfa	fyuA	hly	cnf1	aer	papC	papGII	papGIII	traT
					subgroup	subgroup												
RY07	0	C	non-B2	non-D	non-B1	ND	0	1	0	0	0	0	1	0	0	0	0	1
RY08	0	B2	II	non-D	non-B1	O22	0	1	1	1	0	0	0	0	0	0	0	0
RY09	0	B2	VI	non-D	non-B1	O4	0	1	1	1	1	1	0	1	0	1	0	0
RY10	0	B2	UA	non-D	non-B1	O2b	1	1	1	1	1	1	0	0	0	0	0	1
RY11	0	B2	UA	non-D	non-B1	O2b	0	0	0	1	0	0	0	0	0	0	0	0
RY12	0	D	non-B2	CGA	non-B1	O17	0	1	0	0	0	0	1	0	0	0	0	1
RY13	0	D	non-B2	no CGA	non-B1	ND	0	0	0	0	0	0	0	0	0	0	0	0
RY14	0	B2	III	non-D	non-B1	O6	0	1	1	1	1	1	0	1	0	1	0	0
RY15	0	B1	non-B2	non-D	no CC87	ND	0	1	0	1	0	0	1	0	0	0	0	1
RY19	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	1	1	0	0	0	0	0
RY20	0	B2	IX	non-D	non-B1	O45a	1	1	0	1	0	0	1	1	1	0	1	0
RY21	0	D	non-B2	no CGA	non-B1	ND	0	0	0	0	0	0	0	0	0	0	0	0
RY22	0	B1	non-B2	non-D	no CC87	ND	0	0	0	0	0	0	0	0	0	0	0	0
RY24	0	B23	UA	non-D	non-B1	O4	0	1	1	1	1	1	0	1	0	1	0	1
RY25	0	B2	UA	non-D	non-B1	O2a	0	1	1	1	0	0	0	0	0	0	0	1
RY26	0	B2	UA	non-D	non-B1	O75	1	0	1	1	1	1	0	0	0	0	0	1
RY27	0	B2	IV	non-D	non-B1	O2b	1	1	1	1	0	0	1	0	0	0	0	0
RY28	0	B2	V	non-D	non-B1	O83	1	1	1	1	0	0	1	0	0	0	0	1
RY29	0	B2	III	non-D	non-B1	O6	0	1	1	1	0	0	0	1	0	1	1	1
RY33	0	B2	II	non-D	non-B1	O6	0	1	1	1	1	1	1	0	0	0	0	0
RY34	0	A1	non-B2	non-D	non-B1	ND	0	0	0	0	0	0	0	0	0	0	0	0
RY3501	0	B2	IV	non-D	non-B1	O2b	1	1	1	1	1	1	0	1	0	1	0	0
RY3502	0	B2	IV	non-D	non-B1	O2b	1	1	1	1	1	1	0	1	0	1	0	0
RY36	0	B2	VI	non-D	non-B1	O4	0	1	1	1	1	1	0	1	0	1	0	0
RY37	0	B2	IV	non-D	non-B1	O2b	1	1	1	1	1	1	0	1	0	1	0	0
STA1	0	B2	IV	non-D	non-B1	O2b	1	1	1	1	1	1	0	1	0	1	0	0

*Abbreviations: UA, Unassigned; ND, Not determined; The roman number corresponds to the B2 subgroup. The correspondence with the Achtman MLST schema is as follow: subgroup I = STc131; subgroup II = STc73; subgroup III = STc127; subgroup IV = STc141; subgroup VI = STc12; subgroup VII = STc14; subgroup IX = STc95 [2]. The most anticipated serotypes in extra-intestinal infections were searched by PCR: O1, O2a, O2b, O4, O6, O7, O12, O15, O16, O17, O18, O22, O25a, O25b, O45a, O75, O78.

References

1. Blanco J, Mora A, Mamani R, López C, Blanco M, Dahbi G, et al. National survey of *Escherichia coli* causing extraintestinal infections reveals the spread of drug-resistant clonal groups O25b:H4-B2-ST131, O15:H1-D-ST393 and CGA-D-ST69 with high virulence gene content in Spain. *J Antimicrob Chemother.* 2011;66:2011–21. [PubMed http://dx.doi.org/10.1093/jac/dkr235](http://dx.doi.org/10.1093/jac/dkr235)
2. Johnson JR, Owens K, Manges AR, Riley LW. Rapid and specific detection of *Escherichia coli* clonal group A by gene-specific PCR. *J Clin Microbiol.* 2004;42:2618–22. [PubMed http://dx.doi.org/10.1128/JCM.42.6.2618-2622.2004](http://dx.doi.org/10.1128/JCM.42.6.2618-2622.2004)

3. Clermont O, Couffignal C, Blanco J, Mentré F, Picard B, Denamur E; COLIVILLE and COLIBAFI groups. Two levels of specialization in bacteraemic *Escherichia coli* strains revealed by their comparison with commensal strains. *Epidemiol Infect.* 2017;145:872–82. [PubMed](#)
<http://dx.doi.org/10.1017/S0950268816003010>