
Community Responses during Early Phase of COVID-19 Epidemic, Hong Kong

Kin On Kwok, Kin Kit Li, Henry Ho Hin Chan, Yuan Yuan Yi,
Arthur Tang, Wan In Wei, Samuel Yeung Shan Wong

During the early phase of the coronavirus disease epidemic in Hong Kong, 1,715 survey respondents reported high levels of perceived risk, mild anxiety, and adoption of personal-hygiene, travel-avoidance, and social-distancing measures. Widely adopted individual precautionary measures, coupled with early government actions, might slow transmission early in the outbreak.

Hong Kong was relatively successful in mitigating transmission early in the outbreak of coronavirus disease (COVID-19). Confirmed cases were first reported in the city of Wuhan, China, in December 2019 (1). Situated at the southern tip of China, Hong Kong was at risk for importing COVID-19, given its shared border and high infrastructural and social connectivity with China. In 2019, >236 million passengers crossed the border between China and Hong Kong by land (2). Hong Kong is also vulnerable to virus transmission owing to its high population density and heavy reliance on public transportation. Despite these risks, as of March 20, 2020, transmission control efforts in Hong Kong, as reflected in the numbers of confirmed cases and deaths (256 cases, 4 deaths) (3), had been relatively successful compared with nearby countries and regions, including mainland China (80,967 cases, 3,248 deaths), South Korea (8,652 cases, 94 deaths), and Japan (950 cases, 33 deaths, in addition to the 712 cases from a cruise ship) (4).

JC School of Public Health and Primary Care, The Chinese University of Hong Kong, Hong Kong, China (K.O. Kwok, H.H.H. Chan, Y.Y. Yi, W.I. Wei, S.Y.S. Wong); Stanley Ho Centre for Emerging Infectious Diseases, The Chinese University of Hong Kong, Hong Kong (K.O. Kwok); Shenzhen Research Institute of The Chinese University of Hong Kong, Shenzhen, China (K.O. Kwok); City University of Hong Kong College of Liberal Arts and Social Sciences, Hong Kong (K.K. Li); Sungkyunkwan University College of Software, Seoul, South Korea (A. Tang)

DOI: <https://doi.org/10.3201/eid2607.200500>

Health officials in Hong Kong have enacted multipronged interventions to slow disease spread (5). Adopted strategies include border screening (measuring body temperature, imposing a health declaration form system, imposing a 14-day mandatory quarantine period on persons entering Hong Kong from mainland China; parts of Korea, Japan, France, Germany, and Spain; and all of Italy and Iran), social distancing (shutting down the border, reducing cross-border commuting services, delaying the resumption of classes in schools, arranging telework for civil servants, and suspending of public services), and extending the Enhanced Laboratory Surveillance Program to adult patients with fever and mild respiratory symptoms at emergency departments or general outpatient clinics in the public sector.

The behaviors of the public are important for outbreak management, particularly during the early phase when no treatment or vaccination is available and nonpharmaceutical interventions are the only options. The efficacy of nonpharmaceutical interventions depends on persons' degree of engagement and compliance in precautionary behaviors, such as face-mask wearing, hand hygiene, and self-isolation. Willingness to engage in precautionary behaviors voluntarily depends on risk perception toward the current health threat. In fact, risk perception is a main theme in common health behavior theories (6,7). In addition, with advanced information technology in recent years comes the uncertainty of how risk perception is shaped by various information sources. Hong Kong's experience with outbreaks of novel pathogens (e.g., 2003 severe acute respiratory syndrome [SARS] and 2009 pandemic influenza) also provides a reference point to evaluate the risk perceptions of COVID-19. In comparison, Hong Kong was more affected by SARS than COVID-19 thus far. In 2003, a total of 1,755 persons in Hong Kong contracted SARS, resulting in 299 deaths (8).

In light of the importance of persons' behavior in mitigating transmission and the goal of informing policy formation in a timely manner, we examined risk perceptions and behavioral responses of the general community during the early phase of the COVID-19 epidemic in Hong Kong. Considering the rapid development of the epidemic during the survey period and the potential variability in the adoption of preventive measures among persons, we also examined the temporal changes in anxiety levels, the factors associated with adoption of preventive measures, and sources of information about the epidemic.

The Study

District councilors distributed an online survey including measures of preventive behaviors, general anxiety, risk perceptions, and information exposure to the residents of Hong Kong within 36 hours after detection of the first confirmed case of COVID-19 in Hong Kong (Appendix, <https://wwwnc.cdc.gov/>

EID/article/26/7/20-0500-App1.pdf). The survey was conducted for 3 weeks. We compiled a chronology of major events related to COVID-19 both inside and outside Hong Kong and the number of confirmed cases in Hong Kong before and during the period covered by the survey (Figure 1).

Analysis of 1,715 respondents' data indicated high levels of perceived susceptibility to (89%) and severity of (97%) COVID-19 (Table 1). However, the general anxiety level, measured by the Hospital Anxiety and Depression Scale (9), was mild (9.01 out of 21). Most respondents ($\geq 98\%$) had their daily routines disrupted and were alert to COVID-19. The most trusted information sources were doctors (84%) and radio broadcasts (57%), but they were not the sources by which respondents typically received their information (doctors 5%, broadcast 34%).

Among preventive measures and their perceived efficacy, enhanced personal hygiene (from 78% of respondents disinfecting their homes to 99% wearing




Figure 1. Chronology of major events during the early phase of the coronavirus disease epidemic and laboratory-confirmed cases in Hong Kong, December 30, 2019–February 14, 2020. A, unexplained pneumonia reported in Wuhan, China; B, HK begins temperature screenings at border checkpoints for travelers from Wuhan; C, HK launches preparedness and response plan for novel infectious disease of public health significance, serious response level; D, first death reported in Wuhan; E, World Health Organization (WHO) names disease 2019-nCoV acute respiratory disease and the virus 2019-nCoV (refer to Y for subsequent renaming); F, China confirms human-to-human transmission; G, HK introduces health declaration form system on inbound travelers by air from Wuhan; H, WHO declines to declare COVID-19 a public health emergency of international concern; I, first first confirmed COVID-19 case in HK, halt of sale of high-speed rail tickets to and from Wuhan; J, HK activates emergency response level; K, HK closes public leisure and cultural facilities until further notice; L, WHO declares COVID-19 a public health emergency of international concern; M, United States declares COVID-19 a public health emergency, imposes entry restriction; N, HK imposes 4-week school suspension, 1-week extension for home-office arrangement for civil servants; O, first COVID-19 death outside China in the Philippines; P, HK medical workers strike to call for border shutdown; Q, first COVID-19 death in HK, closure of 4 more border control points; R, 46 foreign airlines cancelled flights to mainland China; S, HK implements further port hygiene measures; T, HK offers home-office arrangement for civil servants until February 16; U, first death of a doctor in China (Wuhan); V, HK begins mandatory 14-day quarantine on persons entering from China; W, HK reports COVID-19 cluster involving 9 people in a gathering on January 26; X, HK reports COVID-19 cluster involving 5 residents (2 families) in the same building; Y, WHO and ICTV rename disease COVID-19 and virus SARS-CoV-2; Z, HK extends home-office arrangement for civil servants until February 23, school suspension until March 16. HK, Hong Kong.

Table 1. Risk perception of the community toward COVID-19 during the early phase of the COVID-19 epidemic in Hong Kong*

Characteristic	No. (%) respondents				
	Level 1	Level 2	Level 3	Level 4	Level 5
Perceived susceptibility (assuming no preventive measure)					
How likely you will be infected†	776 (45)	751 (44)	160 (9)	23 (1)	5 (0)
How likely your families will be infected‡	924 (54)	660 (38)	113 (7)	14 (1)	4 (0)
Perceived severity					
Seriousness of symptoms caused by SARS-CoV-2‡	1102 (64)	569 (33)	33 (2)	7 (0)	4 (0)
Chance of having COVID-19 cured§	190 (11)	552 (32)	708 (41)	239 (14)	26 (2)
Chance of survival if infected with COVID-19§	136 (8)	476 (28)	788 (46)	290 (17)	25 (1)

*COVID-19, coronavirus disease; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2.

†Level 1, very likely; level 2, likely; level 3, neutral; level 4, unlikely; level 5, very unlikely.

‡Level 1, very serious; level 2, serious; level 3, neutral; level 4, not serious; level 5, not serious at all.

§Level 1, very low; level 2, low; level 3, neutral; level 4, high; level 5, very high.

facemasks) and travel avoidance (from 90% avoiding Hubei Province, China, to 92% avoiding mainland China altogether) were frequently adopted and were considered effective (>90%) (Figure 2). The adoption of social-distancing measures was moderate to high (from 39% respondents avoiding public transportation to 93% avoiding contact with persons with respiratory disease symptoms). Higher levels of adoption of social-distancing measures were associated with being female, living in the New Territories (1 of the 3 geographic regions in Hong Kong that shares the border with mainland China), perceiving oneself as having a good understanding of COVID-19, and being more anxious (Table 2).

Conclusions

The relative success in transmission control in Hong Kong could be attributed to the widely adopted precautionary behaviors of the public, together with early government interventions (e.g., border control

and compulsory quarantine for those from affected regions). Unlike in many other countries, visitors from mainland China have never been fully banned from entering Hong Kong. The citizens of Hong Kong assumed responsibility for infection control on their own and became very attentive to personal preventive measures. Our findings showed that nearly all respondents adopted enhanced personal hygiene (e.g., wearing facemasks) and travel avoidance. The experience in outbreak management during the 2003 SARS epidemic might also have contributed to these swift and strong psychological and behavioral responses. Metaphorically, these responses resembled a secondary immune response, which is fast and strong during re-exposure to the same pathogen.

The case of Hong Kong demonstrates the extent to which voluntary preventive measures by persons might be required for slowing transmission (e.g., ≥78% adoption of enhanced personal-hygiene measures, ≥90% adoption of travel-avoidance, and




Figure 2. Perceived efficacy and actual adoption of precautionary measures to prevent transmission of severe acute respiratory syndrome coronavirus 2 and avoid contracting coronavirus disease, Hong Kong.

Table 2. Factors associated with greater adoption of social-distancing interventions during the early phase of the COVID-19 epidemic in Hong Kong*

Characteristic	aOR (95% CI)	p value†
Sex		
M	Referent	
F	1.31 (1.06–1.63)	0.01
Age group, y		
18–24	Referent	
25–34	1.26 (0.97–1.63)	0.08
35–44	1.17 (0.88–1.56)	0.28
45–54	1.34 (0.94–1.92)	0.11
>55	0.93 (0.61–1.41)	0.74
District of residence		
Hong Kong Island	Referent	
Kowloon East	0.96 (0.68–1.36)	0.83
Kowloon West	0.95 (0.62–1.46)	0.82
New Territories East	1.57 (1.18–2.11)	0.00
New Territories West	1.37 (1.02–1.85)	0.04
Left Hong Kong in the previous month		
No	Referent	
Yes	0.72 (0.57–0.91)	0.01
Made regular visits to mainland China		
No	Referent	
Yes	0.48 (0.24–0.91)	0.03
Perceived understanding about COVID-19		
Not well or not well at all	Referent	
Neutral	1.07 (0.76–1.51)	0.70
Well or very well	1.80 (1.27–2.56)	0.00
Presence of chronic diseases		
No	Referent	
Yes	0.77 (0.55–1.06)	0.11
Anxiety level		
Normal	Referent	
Mild	1.38 (1.08–1.76)	0.01
Moderate or severe	1.71 (1.34–2.17)	0.00

*aOR, adjusted odds ratio; COVID-19, coronavirus disease.

†By 2-tailed *t*-test.

39%–93% adoption of social-distancing). Being in agreement with the findings of Anderson et al. (10), we hope that these behavioral standards are useful in promoting person-level preventive measures for countries in the early phase of the COVID-19 outbreak, especially when border-control measures are not viable. This high level of civil engagement toward disease control also enables most businesses to continue as usual, which reduces the economic toll from strict quarantine measures.

In addition, we consider the increased anxiety levels reported as a double-edged sword. On one hand, anxiety can motivate precautionary measures. On the other hand, it might adversely affect school, work, or family life. Besides providing accurate information about the epidemic, public health institutions (e.g., Hong Kong Department of Health) also should promote a healthy lifestyle and psychological well-being. Further discussion of the interpretation of some specific findings, including assessing the sustainability of the preventive measures, the general anxiety level of the public in different outbreaks, the effective

communication channels for COVID-19 information, and the drivers of social-distancing behaviors are provided (Appendix).

In conclusion, we identified high levels of risk perception regarding COVID-19 in the community in Hong Kong. Most respondents were alert to the disease progression of COVID-19 and adopted self-protective measures. Our findings contribute to the body of research examining the psychobehavioral responses of the public, in addition to the already widely studied biological and mechanistic aspects of COVID-19, during the early phase of the current COVID-19 epidemic. The timely psychological and behavioral assessment of the community can inform subsequent intervention and risk-communication strategies as the epidemic progresses.

Acknowledgments

We thank Edelweiss Chui, Ivy Huang, and Margaret Tsoi for technical support.

K.O.K. acknowledges support from the Research Fund for the Control of Infectious Diseases, Hong Kong (grant no. INF-CUHK-1), the General Research Fund (reference no. 14112818), the Wellcome Trust Fund (reference no. 200861/Z/16/Z), and internal funding from the Chinese University of Hong Kong.

K.O.K., K.K.L., A.T., W.I.W., and S.Y.S.W. conceptualized the study; K.O.K. and W.I.W. performed data curation; K.O.K., H.H.H.C., Y.Y.Y., and W.I.W. analyzed the data; K.O.K., W.I.W., and S.Y.S.W. wrote the first draft of the manuscript; K.K.L., H.H.H.C., Y.Y.Y., and A.T. edited the manuscript.

About the Author

Dr. Kwok is an assistant professor affiliated with the JC School of Public Health and Primary Care at The Chinese University of Hong Kong. His primary research interests include infectious disease epidemiology and infection control, with a current focus on mitigating emerging infectious disease outbreaks from the mathematic, epidemiologic, and behavioral perspectives.

References

1. Wuhan Municipal Health Commission. Briefing on the current pneumonia epidemic situation. 2019 [cited 21 Mar 2020]. <http://wjw.wuhan.gov.cn/front/web/showDetail/2019123108989>
2. Immigration Department (Hong Kong). Immigration clearance. 2020 [cited 21 Mar 2020]. <https://www.immd.gov.hk/eng/facts/control.html>
3. Hong Kong Centre for Health Protection. Latest situation of cases of COVID-19. 2020 [cited 21 Mar 2020]. https://www.chp.gov.hk/files/pdf/local_situation_covid19_en.pdf

4. Hong Kong Centre for Health Protection. Countries/ areas with reported cases of coronavirus disease-2019 (COVID-19). 2020 [cited 21 Mar 2020]. https://www.chp.gov.hk/files/pdf/statistics_of_the_cases_novel_coronavirus_infection_en.pdf
5. Government of the Hong Kong Special Administrative Region. Hong Kong's multi-pronged response to COVID-19. 2020 [cited 21 Mar 2020]. https://www.hketojakarta.gov.hk/doc/pdf/Factsheet_coronavirus_Mar_17_E.pdf
6. Skinner CS, Tiro J, Champion VL. The health belief model. In: Glanz K, Rimer BK, Viswanath K, editors. Health behavior: theory, research, and practice. 5th edition. San Francisco: Jossey-Bass; 2015. p. 75-94.
7. Rogers RW, Prentice-Dunn S. Protection motivation theory. In: Gochman DS, editor. Handbook of health behavior research I: personal and social determinants. New York: Springer; 1997. p. 113-32.
8. Leung GM, Ho LM, Lam TH, Hedley AJ. Epidemiology of SARS in the 2003 Hong Kong epidemic. *Hong Kong Med J*. 2009;15(Suppl 9):12-6.
9. Leung CM, Wing YK, Kwong PK, Lo A, Shum K. Validation of the Chinese-Cantonese version of the hospital anxiety and depression scale and comparison with the Hamilton Rating Scale of Depression. *Acta Psychiatr Scand*. 1999;100:456-61. <https://doi.org/10.1111/j.1600-0447.1999.tb10897.x>
10. Anderson RM, Heesterbeek H, Klinkenberg D, Hollingsworth TD. How will country-based mitigation measures influence the course of the COVID-19 epidemic? *Lancet*. 2020;395:931-4. [https://doi.org/10.1016/S0140-6736\(20\)30567-5](https://doi.org/10.1016/S0140-6736(20)30567-5)

Address for correspondence: Kin On Kwok, Room 419, 4/F, JC School of Public Health and Primary Care Building, Prince of Wales Hospital, Shatin, N.T., Hong Kong; email: kkokwok@cuhk.edu.hk; or Wan In Wei, Room 427, 4/F, JC School of Public Health and Primary Care Building, Prince of Wales Hospital, Shatin, N.T., Hong Kong; email: vivian1628@cuhk.edu.hk

etymologia

prions, *Plasmodium knowlesi*, cholera, tularemia, *Eptesicus fuscus*, *synecytium*, *Klebsiella*, Kaposi, *Leptospira*, sapovirus, yaws, *Rickettsia*, *Vibrio vulnificus*, Quinine, variola, *Campylobacter*, *Acinetobacter*, Chagas disease, rotavirus, Lyssavirus, *Aspergillus*, botulism, *Escherichia coli*, sypphilis, knemidocoptic mange, *Babesia*, hemozoin, *Naegleria fowleri*, *Ehrlichia*, Leishmaniasis, *Anopheles*, *Bordetella*, rabies, Verona integrin, vaccination, Artemisinin, Dengue, Zika virus, Herpesvirus, Borna disease virus, Ebola, *Franciscella tularensis*, typhus, *Shigella*, orf, *Coxiella burnetii*, kobuvirus, *Candida*, Rickettsia, *Orientia tsutsugamushi*, Bocavirus, chimera, Q fever, Norovirus, tuberculosis, quarantine, Mange, *Brucella*, Malaria, measles, tetanus, Chikungunya, pertactin, *Borrelia*, Leprosy, influenza, Calcivirus, quarantine, Peste des petits ruminants, meliodosis, Diphtheria, *O'nyong-nyong virus*, *Pseudoterranova azarasi*, pertussis, Merkel cells, *Ignatzschineria*, Glanders, *Yersinia*

featured monthly in **EMERGING INFECTIOUS DISEASES** <http://wwwnc.cdc.gov/eid/articles/etymologia>

Community Responses during Early Phase of COVID-19 Epidemic, Hong Kong

Appendix

Methods

Subject Recruitment

A cross-sectional online survey was conducted within 36 hours after the first confirmed Coronavirus Disease 2019 (COVID-19) case was reported in Hong Kong. To ensure good coverage of the general community in Hong Kong, chairpersons and vice-chairpersons of all eighteen district councils and all individual councilors of the 452 District Council Constituency Areas (DCCAs) were approached by electronic mails and their contact numbers listed in the District Council Web sites (<https://www.districtcouncils.gov.hk/index.html>) for survey dissemination. District councilors were invited to share our survey link and promotion messages on their webpages, social media platforms or any channels which they usually use to convey information to their targeted residents, but in general there was no restriction on their dissemination. Individuals who were aged 18 or above, understood Chinese and lived (on average) over 5 days per week in Hong Kong in the last month were eligible to participate. Respondents were compensated with a HKD10 cash coupon if they indicated willingness for receipt. To avoid duplicated responses from the same respondent, the survey could only be taken once from the same electronic device. To start this survey, respondents were asked to indicate their willingness for participation by answering this first question “Are you willing to participate?.” Only those who answered “Yes” could go on with the survey.

Respondent Characteristics

Respondents were asked about their demographics (including sex, age, living district, education attainment, household income), self-perceived health status, travel history in the past month, occurrence of respiratory symptoms in the past fourteen days. The scales and response sets of these measures are self-explanatory in the respective result tables. Respondents were also

asked about anxiety level using the Chinese-Cantonese version of the Hospital, Anxiety and Depression scale - Anxiety (HADS-A) (0–7 = Normal or no anxiety; 8–10 = mild anxiety; 11–14 = moderate anxiety; 15–21 = severe anxiety) (1). Although HADS-A is intended for screening clinically significant anxiety symptoms in clinical populations, many studies have showed that it is valid for community populations (2,3), including employees (4), general population aged 65–80 years in Sweden (5), an Italian community sample aged 18–85 years (6). As a complementary measure, the state anxiety level of a subset of respondents was assessed on a four-point scales (1 = almost never; 4 = almost always) with the validated Chinese version of State-Trait Anxiety Inventory (STAI) (7) adapted from the original STAI (8). State anxiety from a 10-item STAI was used when comparing to past studies wherever applicable.

Risk Perception

Risk perception toward COVID-19 was measured by two very relevant psychological dimensions (9): (i) perceived susceptibility, and (ii) perceived severity. The first dimension was proxied by how likely one considered oneself (his/her families) would be infected with COVID-19 if no preventive measure was taken. The second dimension was proxied by how one rated the seriousness of symptoms caused by COVID-19, their perceived chance of having COVID-19 cured and that of survival if infected with COVID-19. The items are shown in Table 1 (<https://wwwnc.cdc.gov/EID/article/26/7/20-0500-T1.htm>). Subjects were also asked to rate the relative severity of COVID-19 compared with common non-communicable diseases (NCDs) and previous outbreaks by novel pathogens in Hong Kong. Responses were captured with a five-point Likert scale.

Information Exposure

Respondents were asked about the sources from which they obtained information about COVID-19, and how much they trust those sources. They were also asked about the types of information that they wanted to receive.

Preventive Measures

Respondents were asked whether they performed precautionary measures and what their perceived efficacy of those measures are. Three types of precautionary measures were considered: hygienic practices, social distancing and travel avoidance.

Ethics Consideration

This study has been approved by the Survey and Behavioral Research Ethics Committee of The Chinese University of Hong Kong.

Statistical Analysis

Frequency and proportions of responses were tabulated. Demographics of respondents were compared to the 2016 population by-census in Hong Kong with Cohen's w effect size (small: 0.1; medium: 0.3; large: 0.5) (10). Regression models were used to test for temporal change in anxiety level and to identify factors associated with greater adoption of social-distancing preventive measures (proxied by adopting five or more social-distancing precautionary measures). For temporal change in anxiety level, the HADS-A score was considered the outcome with the survey date being the exposure. Adjusted odds ratio (aOR) and 95% confidence interval (CI) were estimated. Candidate variables included: demographics of respondents, self-perceived health status, travel history and anxiety level. A statistical significance of 0.05 was specified. Analysis was performed in R.

Results

The survey was conducted from 24 January 2020 to 13 February 2020 (Figure 1, <https://wwwnc.cdc.gov/EID/article/26/7/20-0500-F1.htm>). Our survey period covers important clinical incidences, including first local death case and first overseas death case (Philippines), and social incidences, including healthcare workers on strike to call for entire border shutdown. It was also amid of the start-up of large-scale social-distancing interventions, including halt of sales of high-speed rail tickets to and from Wuhan, closure of public cultural and leisure facilities and deferral of school resumption. Meanwhile, alongside the launch of this survey was the escalating official threat tone on COVID-19: The World Health Organization (WHO) declared the COVID-19 epidemic as a public health emergency of international concern, with Hong Kong activated the emergency response level.

Respondent Characteristics

There were initially 2478 clicks of the survey link which fulfill the inclusion criteria, with 6%–31% missingness on demographics variables. Therefore, for a complete-case analysis, 1715 responses were analyzed. Appendix Table 1 shows the respondent characteristics. Many of the

respondents are female (69%; 1176/1715), of young age (18–44 years) (80%; 1380/1715), working population (68%; 1168/1715). The study sample is moderately comparable to the population in terms of living district and sex (effect size = 0.27). Appendix Table 2 shows the background health conditions and travel history of respondents. The majority perceived their health status as good or very good (78%; 1331/1715), a quarter of them experienced respiratory symptoms in the past 14 days (25%; 423/1715) and traveled outside Hong Kong in the previous month (24%; 408/1715). Among the 408 respondents who were abroad, at least 24% of them went to the Mainland China excluding Macau.

Risk Perception

Table 1 (<https://wwwnc.cdc.gov/EID/article/26/7/20-0500-T1.htm>) shows the perceived susceptibility and perceived severity toward COVID-19 among respondents. Most respondents regarded themselves as likely to be infected with COVID-19 (very likely/likely: 89%), and most considered the symptoms of COVID-19 (if infected) as serious (very serious/serious: 97%). Less than a quarter of the respondents thought that it was likely to have COVID-19 cured (if infected) (15%), and only 18% thought that it was likely to survive through COVID-19. When referencing to existing diseases (Appendix Table 3), almost all respondents (>98%) consider equivalent disease severity between COVID-19 and SARS. This magnitude was similar to other deadly NCDs (85%–96%), but much higher than the annual seasonal influenza (66%).

Most respondents were worried about COVID-19 (97%; 1667/1715), and they claimed that their daily routines were slightly (42%; 727/1715) or greatly (56%; 955/1715) disrupted. The average HADS-A score is 9.01 out of 21 (standard deviation [SD]:4.23); while the average score of state anxiety by the full-version STAI, from 804 complete responses, is 2.66 (SD: 0.58). A significantly increasing time trend in HADS-A score is identified ($p < 0.05$) (Appendix Figure 1).

Information Exposure

Nearly all respondents were continuously alert to the disease progression of COVID-19 (99.5%; 1707/1715) and actively searched for related information (83%; 1431/1715). Appendix Table 4 lists the types of COVID-19 information wanted by the 1639 (96%) respondents who indicated such need. Information which respondents were most interested were: distribution of

cases (92%), number of infected individuals (91%), infection control interventions undertaken by local officials (88%), and preventive measures (87%).

Appendix Figure 2 shows the sources from which respondents obtained information about COVID-19, and how well the information sources were trusted. The most trusted sources were doctors (84%; very reliable/reliable: 1449/1715), but only 5% (87/1715) respondents could obtain information from them. The next two most trusted sources were broadcast (57%) and newspaper (54%), but they were used by less than 40% of the respondents. On the other hand, the two most common information sources were social platforms (94%; 1608/1715) and Web sites (regardless of official or unofficial) (90%; 1539/1715), but they were rated as reliable or very reliable by only 26% and 16%–23% of the respondents respectively. Only 16% (269/1715) of respondents found information from official Web sites reliable or very reliable.

Preventive Measures

Figure 2 (<https://wwwnc.cdc.gov/EID/article/26/7/20-0500-F2.htm>) shows the adoption of precautionary measures by respondents and their perceived efficacy. Enhanced personal hygiene practices (including wearing masks, cleaning hands and better coughing and sneezing etiquette) and avoid traveling to Mainland China were adopted by most respondents (>89%), and these practices were considered very effective or effective (>90%). For social-distancing measures, although they were considered useful in preventing COVID-19 (very effective/effective: $\geq 70\%$), their actual adoption was moderate-to-high (range 39%–93%).

Table 2 (<https://wwwnc.cdc.gov/EID/article/26/7/20-0500-T2.htm>) shows the regression analysis results for greater adoption (five or more) of social-distancing interventions during the early phase of this COVID-19 epidemic. Being female (aOR 1.31; 95% CI 1.06–1.63), living in the NT (aOR 1.37–1.57), perceived as having good understanding of COVID-19 (aOR 1.80; 95% CI 1.27–2.56), being more anxious (aOR 1.38–1.71) were positively associated with greater adoption.

Discussion

This study provides timely assessment of the risk perception, information exposure and adoption of precautionary measures during the initial phase of the COVID-19 epidemic in Hong Kong. Despite disease uncertainty (including transmissibility, route of transmission and

pathogenicity) at the early stage, individuals in the community had high perceived risk toward COVID-19 at large, viz: high perceived susceptibility and high perceived severity. A slightly increasing general anxiety level was observed over the 3-week study period. Enhanced personal hygiene and travel avoidance were adopted by nearly all respondents, higher propensity of adopting greater degree of social-distancing measures were associated with being female, living in the New Territories, perceived as having good understanding of COVID-19 well, work status except students and being more anxious. Our results have several immediate and significant public health implications.

First, our results provide the baseline psychological and behavioral responses of the community against which current infection control strategies fit in. With the high perceived risk and large proportion of individuals adopting preventive measures in the community at the beginning, during which the accumulated number of local cases is 130 (almost 7 weeks since the first case) with a significant initial portion of them being imported cases (11), we have an edge to block local transmission. This suggests that efforts to curb imported cases were efficient at the early phase of this outbreak. Following the enactment of a 14-day quarantine period for individuals entering Hong Kong from the Mainland China, Italy, Iran, and other regions with outbreaks, and the emergence of clustered local cases, the next important strategy on the agenda is to stabilize the supply of preventive materials, such as masks, so that the blockage of local transmission chain can be sustained. Besides sustainable supply, how much longer the public can maintain this high adoption of preventive measures without seeing the light at the end of tunnel has become a concern. Follow-up studies on the sustainability of such behaviors are needed.

Second, our results reveal the risk perception in the community, which is an important piece of information to enhance epidemic control (12). Although the epicenter of the COVID-19 epidemic is Wuhan, the perceived risk of the community in Hong Kong was high. For emotional status, the HADS-A score in our survey (9.01 out of 21) suggests that the community had mild anxiety. The community was more anxious about the current COVID-19 epidemic (mean of the 10-item state anxiety from STAI = 2.57; SD = 0.62) than the 2003 SARS outbreak (mean of the 10-item state anxiety from STAI = 2.24; SD = 0.58) (p-value from two-sample t-test <0.05) (13). The significant time trend associated with HADS-A (Appendix Figure 1) suggests that the community became more and more anxious as new cases and new incidences came up (Figure 1).

Third, our results suggest an alternative strategy for better risk communication. The large proportion of respondents were alert to COVID-19 (99.5%) or actively searching for related information (83%) highlighted the role of social media in shaping risk perception and epidemic-related emotion. It is particularly important amid of much disease uncertainty as mass scares can be triggered easily. Considering the high level of trust given by respondents to doctors and the low level of trust to the two most frequently used information sources, social platform and Web sites, health officials can collaborate frequently with associations of medical doctors, and invite them to help propagating official information in more sociable channels. This strategy is deemed more acceptable by the community than relying solely on the official channel, given only 16% of respondents rates official Web sites as reliable or very reliable. Our results also shortlisted information preferred by the community among an upsurge of disease-related information during the early stage (Appendix Table 4).

Fourth, our results pinpoint the drivers for greater level of adoption of social-distancing precautionary measures. In line with literatures that being female and an elevated anxiety level prompted compliance of precautionary measures (13,14), we also identified similar association in this survey (Table 2, (<https://wwwnc.cdc.gov/EID/article/26/7/20-0500-T2.htm>)). Interestingly, specific to this COVID-19 epidemic, residents in the New Territories were more likely to comply with social-distancing precautionary measures than their counterparts in other areas of Hong Kong. Separating Hong Kong and the Guangdong Province are two busiest custom borders, Lo Wu and Lok Ma Chau, such that the residents in the New Territories may consider themselves at greater risk of infection. Those who claimed they understood COVID-19 were more likely to adopt preventive measures, suggesting mass promotion of knowledge about COVID-19 in the community can boost uptake of precautionary measures. On the other hand, the less propensity to adopt precautionary measures among individuals who left Hong Kong in the previous month or who regularly visited China. Some of these visits might be work- or family-related. Social interactions became difficult to avoid during travels such as the use of public transportation, going out, and going to crowded places (as mentioned above, the borders are always crowded except with border control). These findings reinforce the need for border screening and for promoting social hygiene amid of epidemic times.

Fifth, this local study has profound implication to overseas countries undergoing the initial phase of the COVID-19 epidemic. The WHO European region has been accumulating

COVID-19 cases, but in only 4 days (22–25 February 2020), the number of laboratory-confirmed cases in Italy has risen from 9 to 229 (15–18). Recently on 24 February 2020, the Ministry of Health announced the first COVID-19 case in Iraq. The presence of initial cases, aligning with the human-to-human (19) and asymptomatic (20) transmission, suggest that many countries may experience the initial phase of the COVID-19 epidemic soon. Results of this survey serve as a reference for overseas health officials to better prepare their containment strategies and handle the potential mass scares in their community.

Strengths and Limitations

This study has two strengths. First, it started within 36 hours after the detection of first local cases. This early start enables timely assessment of the community responses such that there is sufficient gap period to inform intervention policies. Second, our recruitment method, online survey via dissemination by DCCA councilors, is the first of its kinds to capture responses during public holidays while maintaining good geographic representation. The COVID-19 epidemic was amid of the Chinese New Year holidays and a series of large-scale social-distancing interventions enacted by Hong Kong government, particularly the home-office arrangement for employees. Therefore, the conventional random digit dialing approach adopted in the past local outbreaks (13,21,22) was not possible. And the involvement of all 452 DCCA councilors allows a thorough representation of every district in Hong Kong in the absence of a universal email database.

This study has two limitations. First, with an online approach, responses of those without internet access, particularly the oldest age group (55 years or above), were under-represented. Despite this, online surveys were the only feasible means of data collection during outbreak times. Second, this survey was conducted during the early phase that temporal variations of responses are not captured as the epidemic progresses. However, contact information were collected from this study cohort and follow-up surveys will be carried out as the disease progresses.

References

1. Leung CM, Wing YK, Kwong PK, Lo A, Shum K. Validation of the Chinese-Cantonese version of the hospital anxiety and depression scale and comparison with the Hamilton Rating Scale of

- Depression. *Acta Psychiatr Scand*. 1999;100:456–61. [PubMed](#) <https://doi.org/10.1111/j.1600-0447.1999.tb10897.x>
2. Bjelland I, Dahl AA, Haug TT, Neckelmann D. The validity of the Hospital Anxiety and Depression Scale. An updated literature review. *J Psychosom Res*. 2002;52:69–77. [PubMed](#) [https://doi.org/10.1016/S0022-3999\(01\)00296-3](https://doi.org/10.1016/S0022-3999(01)00296-3)
 3. Snaith RP. The Hospital Anxiety And Depression Scale. *Health Qual Life Outcomes*. 2003;1:29. [PubMed](#) <https://doi.org/10.1186/1477-7525-1-29>
 4. Bocéréan C, Dupret E. A validation study of the Hospital Anxiety and Depression Scale (HADS) in a large sample of French employees. *BMC Psychiatry*. 2014;14:354. [PubMed](#) <https://doi.org/10.1186/s12888-014-0354-0>
 5. Djukanovic I, Carlsson J, Årestedt K. Is the Hospital Anxiety and Depression Scale (HADS) a valid measure in a general population 65-80 years old? A psychometric evaluation study. *Health Qual Life Outcomes*. 2017;15:193. [PubMed](#) <https://doi.org/10.1186/s12955-017-0759-9>
 6. Iani L, Lauriola M, Costantini M. A confirmatory bifactor analysis of the Hospital Anxiety and Depression Scale in an Italian community sample. *Health Qual Life Outcomes*. 2014;12:84. [PubMed](#) <https://doi.org/10.1186/1477-7525-12-84>
 7. Shek DT. The Chinese version of the State-Trait Anxiety Inventory: its relationship to different measures of psychological well-being. *J Clin Psychol*. 1993;49:349–58. [PubMed](#) [https://doi.org/10.1002/1097-4679\(199305\)49:3<349::AID-JCLP2270490308>3.0.CO;2-J](https://doi.org/10.1002/1097-4679(199305)49:3<349::AID-JCLP2270490308>3.0.CO;2-J)
 8. Spielberger CD, Gorsuch RL, Lushene R, Vagg PR, Jacobs GA. *Manual for the State-Trait Anxiety Inventory*. Palo Alto, California: Consulting Psychologists Press; 1983.
 9. El-Toukhy S. Parsing susceptibility and severity dimensions of health risk perceptions. *J Health Commun*. 2015;20:499–511. [PubMed](#) <https://doi.org/10.1080/10810730.2014.989342>
 10. Cohen J. *Statistical power analysis for the behavioral sciences*. 2nd ed. Hillsdale, N.J.: L. Erlbaum Associates; 1988.
 11. Hong Kong Centre for Health Protection. Latest situation of cases of COVID-19. 2020 [cited 2020 Mar 21]. https://www.chp.gov.hk/files/pdf/local_situation_covid19_en.pdf
 12. Herrera-Diestra JL, Meyers LA. Local risk perception enhances epidemic control. *PLoS One*. 2019;14:e0225576. [PubMed](#) <https://doi.org/10.1371/journal.pone.0225576>

13. Leung GM, Lam TH, Ho LM, Ho SY, Chan BH, Wong IO, et al. The impact of community psychological responses on outbreak control for severe acute respiratory syndrome in Hong Kong. *J Epidemiol Community Health*. 2003;57:857–63. [PubMed](#) <https://doi.org/10.1136/jech.57.11.857>
14. Bults M, Beaujean DJ, de Zwart O, Kok G, van Empelen P, van Steenberghe JE, et al. Perceived risk, anxiety, and behavioural responses of the general public during the early phase of the Influenza A (H1N1) pandemic in the Netherlands: results of three consecutive online surveys. *BMC Public Health*. 2011;11:2. [PubMed](#) <https://doi.org/10.1186/1471-2458-11-2>
15. World Health Organization. Coronavirus disease (COVID-2019) Situation Report - 33. 2020 [cited 2020 Mar 25]. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200222-sitrep-33-covid-19.pdf?sfvrsn=c9585c8f_4
16. World Health Organization. Coronavirus disease (COVID-2019) Situation Report - 34. 2020 [cited 2020 Mar 25]. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200223-sitrep-34-covid-19.pdf?sfvrsn=44ff8fd3_2
17. World Health Organization. Coronavirus disease (COVID-2019) Situation Report - 35. 2020 [cited 2020 Mar 25]. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200224-sitrep-35-covid-19.pdf?sfvrsn=1ac4218d_2
18. World Health Organization. Coronavirus disease (COVID-2019) Situation Report - 36. 2020 [cited 2020 Mar 25]. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200225-sitrep-36-covid-19.pdf?sfvrsn=2791b4e0_2
19. Paules CI, Marston HD, Fauci AS. Coronavirus Infections—More Than Just the Common Cold. *JAMA*. 2020;323:707; Epub ahead of print. [PubMed](#) <https://doi.org/10.1001/jama.2020.0757>
20. Bai Y, Yao L, Wei T, Tian F, Jin DY, Chen L, et al. Presumed Asymptomatic Carrier Transmission of COVID-19. *JAMA*. 2020; Epub ahead of print. [PubMed](#) <https://doi.org/10.1001/jama.2020.2565>
21. Lau JT, Yang X, Tsui H, Kim JH. Monitoring community responses to the SARS epidemic in Hong Kong: from day 10 to day 62. *J Epidemiol Community Health*. 2003;57:864–70. [PubMed](#) <https://doi.org/10.1136/jech.57.11.864>
22. Cowling BJ, Ng DM, Ip DK, Liao Q, Lam WW, Wu JT, et al. Community psychological and behavioral responses through the first wave of the 2009 influenza A(H1N1) pandemic in Hong Kong. *J Infect Dis*. 2010;202:867–76. [PubMed](#) <https://doi.org/10.1086/655811>

Appendix Table 1. Respondent characteristics

Characteristics	No. respondents (%), n = 1715	Effect Size*
Sex		0.27
Male	539 (31)	
Female	1176 (69)	
Age group, y		0.82
18–24	441 (26)	
25–34	558 (33)	
35–44	381 (22)	
45–54	197 (11)	
≥55	138 (8)	
Education attainment		1.14†
Lower secondary or below	58 (3)	
Higher secondary	302 (18)	
Diploma	274 (16)	
Degree or above	1081 (63)	
Living district		0.27
Hong Kong Island	307 (18)	
Kowloon West	128 (7)	
Kowloon East	268 (16)	
New Territories West	471 (27)	
New Territories East	541 (32)	
Employment status		0.57†‡
Employee	1106 (64)	
Employer	62 (4)	
Housekeeper	135 (8)	
Student	285 (17)	
Retired	46 (3)	
Unemployed	81 (5)	
Monthly household income (HKD)		Nil§
≤10,000	104 (6)	
10,001–20,000	277 (16)	
20,001–30,000	297 (17)	
30,001–40,000	233 (14)	
40,001–60,000	290 (17)	
>60,000	257 (15)	
Not disclosed	257 (15)	

*Cohen's w effect size.

†Data for 15 y old or above is extracted from 2016 by-census for comparison.

‡The "unemployed" category is excluded from comparison as it is unavailable from 2016 by-census.

§The "monthly household income" category is excluded from comparison as it is unavailable from 2016 by-census.

Appendix Table 2. Background health conditions and travel history of respondents

Characteristics	No. respondents (%), n = 1715
Self-perceived health status	
Very good / good	1331 (78)
Fair	352 (21)
Very bad / bad	32 (2)
Presence of chronic conditions	
Yes	192 (11)
No	1523 (89)
Medical consultation in the past 14 d*	
Yes	293 (17)
No	1422 (83)
Presence of respiratory symptoms in the past 14 d	
Yes	423 (25)
No	1292 (75)
Leave Hong Kong in the previous month	
Yes†	408 (24)
No	1307 (76)
Regular visitors to the Mainland China	
Yes‡	46 (3)
No	1669 (97)

*Both Chinese and Western medical consultations are included.

†Multiple destinations are allowed. Number of respondents (out of 408) who indicated travel outside Hong Kong in the previous month: outside China (294), China - Guangdong province (96), China - other province (13), Macau (29).


‡Number of respondents (out of 46) who indicated regular visit to the Mainland China: daily (4), weekly (7), monthly (21), quarterly (4), and at most quarterly (10).

Appendix Table 3. Comparison of disease severity


Diseases	Very bad	Bad	Neutral	Not bad	Not bad at all
Emerging infectious disease					
COVID-19	1545 (90)	150 (9)	15 (1)	0 (0)	1 (0)
Existing infectious diseases					
SARS	1551 (91)	133 (8)	21 (1)	2 (0)	4 (0)
2009 pandemic influenza	604 (35)	889 (52)	172 (10)	40 (2)	6 (0)
Seasonal influenza	191 (11)	948 (55)	311 (18)	251 (15)	10 (1)
Noncommunicable diseases					
Diabetes	659 (39)	804 (47)	188 (11)	51 (3)	9 (1)
Cancer	1432 (84)	215 (13)	45 (3)	11 (1)	8 (0)
Heart disease	1123 (66)	502 (29)	66 (4)	17 (1)	3 (0)
Acquired immune deficiency syndrome	1354 (79)	257 (15)	69 (4)	22 (1)	9 (1)

Appendix Table 4. Information wanted by the respondents

Information you want to receive about COVID-19	No. (%), n = 1639
Distribution of cases	1506 (92)
Number of people infected	1497 (91)
Interventions of Hong Kong government	1450 (88)
Preventive measures	1424 (87)
Disease progression	1327 (81)
Symptoms/how to know if one is infected	1310 (80)
Interventions of international organizations	1182 (72)
What to do if infected	1087 (66)
Impact on risk groups	1073 (65)
Risks and consequences	1061 (65)
Interventions of Chinese government	1010 (62)



Appendix Figure 1. Time trend of HADS-A score.



Appendix Figure 2. Information reliability and the access to information of COVID-19.