

Precise Species Identification by Whole-Genome Sequencing of *Enterobacter* Bloodstream Infection, China

Wenjing Wu,¹ Li Wei,¹ Yu Feng, Yi Xie, Zhiyong Zong

The clinical importance of *Enterobacter* spp. remains unclear because phenotype-based *Enterobacter* species identification is unreliable. We performed a genomic study on 48 cases of *Enterobacter*-caused bloodstream infection by using in silico DNA–DNA hybridization to identify precise species. Strains belonged to 12 species; *Enterobacter xiangfangensis* (n = 21) and an unnamed species (taxon 1, n = 8) were dominant. Most (63.5%) *Enterobacter* strains (n = 349) with genomes in GenBank from human blood are *E. xiangfangensis*; taxon 1 (19.8%) was next most common. *E. xiangfangensis* and taxon 1 were associated with increased deaths (20.7% vs. 15.8%), lengthier hospitalizations (median 31 d vs. 19.5 d), and higher resistance to aztreonam, cefepime, ceftriaxone, piperacillin-tazobactam, and tobramycin. Strains belonged to 37 sequence types (STs); ST171 (*E. xiangfangensis*) was most common (n = 6). Four ST171 strains belonged to a defined clone. Precise species identification has greater implications for epidemiology and infection control than treatment.

Enterobacter spp. belongs to the family *Enterobacteriaceae* and is a common pathogen in a variety of infections, such as bloodstream and intraabdominal infections, most of which are healthcare associated (1). *Enterobacter* spp. is the third most common human pathogen, after *Escherichia coli* and *Klebsiella pneumoniae*, and is therefore of clinical importance (1). *Enterobacter* consists of several closely related species (1) that cannot typically be identified precisely by common phenotypic tests. The taxonomy of *Enterobacter* is complicated by the reassignment to other genera of some species that formerly belonged to the *Enterobacter* genus. For example, *E. aerogenes* has been moved to genus *Klebsiella*

(2), *E. agglomerans* to genus *Pantoea* (3), and *E. sakazakii* to genus *Cronobacter* (4). Currently, 14 *Enterobacter* spp. with validly published names exist, and 3 additional *Enterobacter* spp. have tentative species designations awaiting validation under the rules of the International Code of Nomenclature of Bacteria (Bacteriological Code) (Appendix 1 Table 1, <https://wwwnc.cdc.gov/EID/article/27/1/19-0154-App1.pdf>).

Several *Enterobacter* spp., such as *E. asburiae*, *E. cloacae*, and *E. hormaechei*, cause infections in humans (1). *Enterobacter* strains extracted from clinical samples are usually reported as *E. cloacae*, and sometimes *E. asburiae*, *E. hormaechei*, or *E. kobei*, by automated microbial identification systems such as Vitek II (bioMérieux, <https://www.biomerieux.com>). However, such phenotype-based tests are unreliable for species identification of *Enterobacter* and can result in misidentification (1). For instance, all *Enterobacter* spp. have a positive reaction for β -galactosidase, arginine dihydrolase, citrate utilization, sucrose, amygdalin, arabinose, and D-glucose but are negative for lysine decarboxylase, H₂S production, urease activity, indole production, deaminase, and gelatinase (5–7). Differentiating *Enterobacter* spp. by biochemical reactions commonly used in clinical microbiology laboratories is therefore difficult. The differences in clinical importance of each *Enterobacter* species remain largely unknown because they are regularly misidentified in clinical microbiology laboratories.

Because of the substantial reduction in cost of whole-genome sequencing for bacterial strains, we are entering the era of genomic microbiology (8). Newly created methods can determine the overall nucleotide identities between genome sequences and therefore enable more precise species identification (9). Calculation of average nucleotide identity (ANI) between genomes is widely used for species

Author affiliations: West China Hospital, Sichuan University, Chengdu, China (W. Wu, L. Wei, Y. Feng, Y. Xie, Z. Zong); State Key Laboratory of Biotherapy, Chengdu (W. Wu, Y. Feng, Z. Zong)

DOI: <https://doi.org/10.3201/eid2701.190154>

¹These authors contributed equally to this article.

identification. It has been proposed that ANI >96% would guarantee species assignment, whereas ANI of <93% can be considered species differentiation (10). However, ANI values in the range of 93%–96% represent a vague zone in which the boundary of a species might fall (10). DNA–DNA hybridization (DDH) remains the standard for species identification, with a $\geq 70\%$ cutoff recommended to define a species. However, DDH is cumbersome, prone to fluctuation, and requires the availability of type strains. To overcome the shortcomings of DDH, in silico DDH (isDDH) mimics DDH by comparing genome sequences and can be a reliable and convenient tool for species assignment. To provide insight into the potential clinical importance of different *Enterobacter* spp., we performed a genomic study using isDDH to identify bloodstream infection (BSI)–causing *Enterobacter* strains to the species level.

Materials and Methods

Strain and Susceptibility Tests

We collected nonduplicate *Enterobacter* strains recovered from blood cultures during January 2016–June 2018 at West China Hospital of Sichuan University (Appendix 2 Table 1, <https://wwwnc.cdc.gov/EID/article/27/1/19-0154-App2.xlsx>). West China Hospital is a 5,000-bed major referral hospital in western China. Initial species identification and in vitro susceptibility testing were performed by using Vitek II. We determined MICs of colistin by using the broth microdilution method of the Clinical and Laboratory Standards Institute (CLSI) and interpreted susceptibility following CLSI guidelines (11). For colistin and tigecycline, no CLSI breakpoints are available, so we used breakpoint standards defined by the European Committee on Antimicrobial Susceptibility Testing (<https://www.eucast.org>). Multidrug resistance was defined based on the criteria for *Enterobacteriaceae* (12).

Patient Data

West China Hospital has a comprehensive hospital information system, which allowed us to retrieve patient data including age, sex, length of hospitalization, and clinical outcomes (death or discharge) from electronic medical records. One patient (with strain 090040) had an unusually long hospital stay (578 d) because of a medical dispute and was removed from our analysis of length of stay. According to social customs in China, dying at home is preferred over the hospital; it is likely many patients chose to stop treatment and return home if

treatment was not working and patients felt death was imminent. We categorized patients who chose to be discharged but were likely to die in the hospital (judged by the consensus of 2 physicians reviewing blind data) as patients with predicted death. BSI, the type of BSI (primary or secondary to infection of other sites), central line-associated BSI (CLABSI), and healthcare-associated infection were determined by using criteria established by the Centers for Disease Control and Prevention's National Healthcare Safety Network (13,14). We conducted the study in accordance with the amended Declaration of Helsinki. The Ethics Committee of West China Hospital approved the study and waived informed consent.

Short-Read Genome Sequencing, Analysis, and Precise Species Identification

All strains underwent whole-genome sequencing by using the HiSeq X10 platform (Illumina, <https://www.illumina.com>). Genomic DNA was prepared by using the QIAamp DNA mini kit (QIAGEN, <https://www.qiagen.com>). We used Unicycler version 0.4.3 (15), in the conservative mode for increased accuracy, to perform a de novo hybrid assembly. Precise species identification was established by determining the pairwise isDDH between the genome sequence of the query strain and those of type strains of *Enterobacter* spp., including the validly published species and the species awaiting validation (Appendix 1 Table 1). This process was performed by using the Genome-to-Genome Distance Calculator, formula 2 (16). A $\geq 70\%$ cutoff was applied to define a species. In addition, we determined the pairwise ANI of the genome sequence of the query strain and those of type strains of *Enterobacter* spp. (Appendix 1 Table 1) by using JSpecies software (<https://imedeia.uib-csic.es/jspecies>) with a >96% ANI cutoff to define a species (10). Sequence types (STs) were determined by using the genomic sequences to query the multilocus sequence typing database of *E. cloacae* (<https://pubmlst.org/ecloacae>). Antimicrobial resistance genes were identified from genome sequences by using the ABRicate program (<https://github.com/tseemann/abricate>) to query the ResFinder database (<https://genomicepidemiology.org>).

The genome sequences of all *Enterobacter* strains recovered from human blood ($n = 349$, Appendix 2 Table 2) were retrieved from GenBank (accessed 2018 Nov 1). These *Enterobacter* genomes were subjected to precise species identification by using the Genome-to-Genome Distance Calculator as described.

Clonal Relatedness on the Basis of Single-Nucleotide Polymorphisms

We performed single-nucleotide polymorphism (SNP) calling for genome sequences to untangle the clonal relatedness of ST171 strains and to investigate whether the ST171 strains in this study are clonally related to strains recovered elsewhere. All genome sequences of ST171 *Enterobacter* strains ($n = 102$), regardless of types of the host source (human, nonhuman, or unknown) and source (blood, nonblood, or unknown), were retrieved from GenBank. We used the complete chromosome sequence of ST171 strain 34798 (GenBank accession no. CP012165), which was recovered from a bile sample in the United States in 2011, as our reference for mapping. Genome sequences of the strains were mapped against the reference genome by using Snippy version 4.3.6 (<https://github.com/tseemann/snippy>) at default settings. The resulting core SNPs ($n = 1,918$) were concatenated and used to infer a phylogenomic tree by using RAxML version 8.2.12 (17) under the general time-reversible plus gamma model with a 1,000-bootstrap test.

Long-Read Genome Sequencing and Plasmid Analysis for ST171 Strains

We determined plasmid replicons of all ST171 strains in this study by using PlasmidFinder version 2.0 (<https://cge.cbs.dtu.dk/services/PlasmidFinder>). The first *bla*_{NDM-5}-harboring ST171 strain (090011) in this study and the only *bla*_{NDM-1}-harboring ST171 strain (045001) were selected for whole-genome sequencing by using the long-read MinION sequencer (Nanopore, <https://nanoporetech.com>) to obtain complete chromosomal and plasmid sequences. De novo hybrid assembly of short Illumina reads and long MinION reads was performed by using Unicycler version 0.4.3 (15) in conservative mode for increased accuracy. Complete circular contigs were then corrected by using Pilon version 1.22 (18) with Illumina reads for several rounds until no further improvements were reported. Short reads of the remaining four *bla*_{NDM-5}-harboring ST171 strains (090022, 090023, 090055, and 090059) were mapped against the *bla*_{NDM-5}-carrying plasmid (designated pNDM5_090011) of strain 090011 by using BWA version 0.7.17 (H. Ling, unpub. data, <https://arxiv.org/abs/1303.3997>) at default settings.

Statistical Analysis

Continuous variables were presented as median and interquartile range and were compared by using rank-sum test. We used the Pearson χ^2 test, Yates correction for continuity, or Fisher exact test to compare disparities between different groups

for categorical variables. Pearson χ^2 test was used when sample size (n) was ≥ 40 and theoretical frequency (T) ≥ 5 , Yates correction for continuity when $n \geq 40$ and $1 \leq T < 5$, and Fisher exact test when $n < 40$ or $T < 1$. We used SPSS Statistics 21.0 (IBM Inc., <https://www.ibm.com>) to perform statistical analyses. All p values were 2-tailed, and $p < 0.05$ was considered statistically significant. We used PASS version 11.0 (NCSS, <https://www.ncss.com>) to calculate statistical power after using the Wilcoxon test to conduct nonparametric adjustment.

Draft genome sequences of the strains in this study were deposited into GenBank (accession numbers in Appendix 2 Table 1). The complete genome sequence of strain 090011 was deposited into GenBank under accession nos. CP036310–2. and the sequence for strain 045001 was deposited under accession nos. CP043382–5.

Results

Precise Species Identification of *Enterobacter* Strains

A total of 48 nonduplicated *Enterobacter* strains were recovered from blood during our 2.5-year study period and were collected for study (Table 1; Appendix 2 Table 1). Whole-genome sequencing results, isDDH, and ANI values of these strains are summarized in Appendix 2 Table 1. The 48 strains were identified as *E. cloacae* ($n = 42$), *E. asburiae* ($n = 3$), and *E. kobei* ($n = 3$) by Vitek II. However, precise species identification on the basis of isDDH revealed that the most common species was actually *E. xiangfangensis* ($n = 21$) (Table 2); the next most common was an *Enterobacter* sp. ($n = 8$) that has no assigned species name but was previously known as *Enterobacter* cluster III, as defined by Hoffman et al. (6). This species is most closely related to *E. xiangfangensis* with a 66.6% isDDH value and is temporarily assigned taxon 1 here (Appendix 2 Table 2). The remaining strains were assigned to 10 *Enterobacter* species: *E. bugandensis* ($n = 4$), *E. cloacae* ($n = 3$), *E. asburiae* ($n = 2$), *E. hormaechei* ($n = 2$), *E. huaxiensis* ($n = 2$), *E. roggenkampii* ($n = 2$), *E. chuandensis* ($n = 1$), *E. ludwigii* ($n = 1$), *E. sichuanensis* ($n = 1$), and an unnamed *Enterobacter* sp. ($n = 1$) (Table 2). The unnamed species is most closely related to *E. roggenkampii* with a 65.4% isDDH value (Appendix 1 Table 2) and is temporarily assigned taxon 2 here.

Of 349 *Enterobacter* strains in GenBank that were recovered from human blood, most (221, 63.3%) are also *E. xiangfangensis*; taxon 1 is the second most common species with 69 strains (19.8%) (Table 2). The remaining 59 strains (16.9%) belong to 14 species: *E. asburiae* ($n = 14$ [4.0%]), *E. kobei* ($n = 10$ [2.9%]), *E. bugandensis* ($n = 7$ [2.0%]), *E. roggenkampii* ($n = 7$ [2.0%]),

Table 1. *Enterobacter* strains in genomic study of *Enterobacter* bloodstream infection, China*

Strain	Date	Species, by isDDH	ST†	Carbapenemase	BSI types				Hospitalization, d		Death
					P	S	CLA	HA	Before BSI	After BSI	
090034	201706	<i>Enterobacter asburiae</i>	N12		+				0	27	
090058	201805	<i>E. asburiae</i>	879		+		+	+	22	18	
090031	201711	<i>E. bugandensis</i>	N10		+				1	0	+
090283	201610	<i>E. bugandensis</i>	N16			+		+	7	6	
090210	201607	<i>E. bugandensis</i>	499			+			1	16	
090029	201709	<i>E. bugandensis</i>	718	NDM-5	+			+	71	16	+
090028	201708	<i>E. chuandaensis</i>	N9			+			1	11	
090005	201706	<i>E. cloacae</i>	1	NDM-1	+		+	+	29	11	
090016	201703	<i>E. cloacae</i>	519			+		+	17	24	
090014	201712	<i>E. cloacae</i>	922			+		+	0	10	
090027	201704	<i>E. hormaechei</i>	528			+		+	24	38	
090003	201705	<i>E. hormaechei</i>	696		+			+	0	15	
045002	201609	<i>E. huaxiensis</i>	N1		+				0	7	
090008	201709	<i>E. huaxiensis</i>	N1		+			+	3	23	
090017	201702	<i>E. ludwigii</i>	12			+		+	6	8	
045158	201607	<i>E. roggenkampii</i>	N2		+			+	3	7	
090037	201608	<i>E. roggenkampii</i>	984		+		+	+	14	8	
090032	201712	<i>E. sichuanensis</i>	N11		+		+	+	6	16	+
090004	201706	<i>E. xiangfangensis</i>	N3		+			+	6	13	
090006	201707	<i>E. xiangfangensis</i>	N4		+			+	12	8	
090007	201707	<i>E. xiangfangensis</i>	N5		+			+	24	19	
090012	201711	<i>E. xiangfangensis</i>	N6			+			0	3	
090018	201705	<i>E. xiangfangensis</i>	N7		+		+	+	12	14	+
090020	201712	<i>E. xiangfangensis</i>	N8		+		+	+	4	60	
090057	201804	<i>E. xiangfangensis</i>	N15	NDM-1	+		+	+	36	43	
090015	201712	<i>E. xiangfangensis</i>	50		+				0	2	
090026	201704	<i>E. xiangfangensis</i>	50		+			+	11	52	
090035	201712	<i>E. xiangfangensis</i>	127		+			+	13	20	
045001	201801	<i>E. xiangfangensis</i>	171	NDM-1	+		+	+	8	33	
090011	201710	<i>E. xiangfangensis</i>	171	NDM-5	+		+	+	7	14	+
090022	201802	<i>E. xiangfangensis</i>	171	NDM-5	+		+	+	40	20	
090023	201802	<i>E. xiangfangensis</i>	171	NDM-5	+		+	+	26	19	
090055	201806	<i>E. xiangfangensis</i>	171	NDM-5		+		+	14	14	
090059	201805	<i>E. xiangfangensis</i>	171	NDM-5	+		+	+	17	31	
090043	201612	<i>E. xiangfangensis</i>	337			+		+	11	4	+
090013	201711	<i>E. xiangfangensis</i>	418	NDM-5	+		+	+	28	10	+
090038	201609	<i>E. xiangfangensis</i>	418		+			+	3	33	
090060	201806	<i>E. xiangfangensis</i>	550		+		+	+	20	8	+
090042	201611	<i>E. xiangfangensis</i>	828		+			+	5	3	
090036	201605	Taxon 1	N13		+		+	+	13	22	
090019	201706	Taxon 1	78		+			+	16	62	
090030	201709	Taxon 1	78		+		+	+	0	15	
090039	201610	Taxon 1	78			+		+	12	8	+
090021	201801	Taxon 1	97		+		+	+	5	28	
090009	201709	Taxon 1	104			+		+	5	66	
090033	201704	Taxon 1	316		+			+	10	3	
090056	201803	Taxon 1	568		+			+	11	7	
090040	201611	Taxon 2	N14		+		+	+	208	370	

*BSI, bloodstream infection; CLA, central line-associated; HA, healthcare associated; isDDH, in silico DNA-DNA hybridization; P, primary; S, secondary; ST, sequence type.

†There are 16 new sequence types, which are temporarily assigned N1–N16 (Appendix 1 Table 3, <https://wwwnc.cdc.gov/EID/article/27/1/19-0154-App1.pdf>).

E. ludwigii (n = 6 [1.7%]), *E. cloacae* (n = 4 [1.1%]), taxon 1 (n = 2), *E. chengduensis* (n = 1), *E. mori* (n = 1), *E. sichuanensis* (n = 1), and 4 species without assigned species names (n = 1 or 2 for each species, 6 in total; Table 2). The 4 unnamed species were assigned taxon 3–6 (Table 2); the closest species of taxon 3–6 are listed in Appendix 1 Table 2 and are also shown in a phylogenomic tree in Appendix 1 Figure 1. *E. xiangfangensis* and taxon 1 are closely related as shown by their phylogenetic position in the phylogenomic tree of *Enterobacter* spp.

and by their common 66.6% isDDH value (close to the 70% cutoff to define a species). We therefore combined the 2 species in the following analysis.

BSI Types and Characteristics

Most of the 48 BSI cases (n = 36, 75%) were primary BSIs, including 16 cases of CLABSI. BSIs in the remaining 12 cases were secondary; original sources were intraabdominal infection (n = 5), cholangitis (n = 3), urinary tract infection (n = 2), wound infection

(n = 1), and gastrointestinal tract infection (n = 1). Most (n = 41, 85.4%) BSIs caused by *Enterobacter* spp. were healthcare-associated infections. *E. xiangfangensis* and taxon 1 were more likely to cause primary BSI (82.8% vs. 63.2%), CLABSI (27.9% vs. 17.2%), and healthcare-associated BSI (93.1% vs. 73.7%) than were other *Enterobacter* spp. However, the differences were not statistically significant (Table 3).

Two patients who had *Enterobacter* BSIs (1 *E. xiangfangensis* and 1 taxon 1) died in the hospital. In addition, 7 patients with *Enterobacter* BSIs (4 *E. xiangfangensis*, 2 *E. bugandensis*, and 1 *E. sichuanensis*) did not respond to treatment and were discharged in critical condition. These 7 case-patients were categorized as patients in whom death was predicted. The death rate for *Enterobacter* BSI was 18.8% (9/48); the death rate (20.7% [6/29]) of BSI caused by *E. xiangfangensis* or taxon 1 was not statistically different (15.8% [3/19] p>0.05) (Table 3) from that of BSI caused by other *Enterobacter* spp. Of note, 2 of the 4 patients with *E. bugandensis*-caused BSI had poor outcomes (predicted death) (Table 1). BSIs caused by *E. xiangfangensis* or taxon 1 were more common in younger patients and resulted in lengthier overall hospitalizations (median 33 vs. 19.5 d; p>0.05) (Table 3) than BSIs caused by other *Enterobacter* spp. This difference was largely because of the prolonged length of stay (median 11 d vs. 4.5 d; p>0.05) before the episode of BSI.

Antimicrobial Susceptibility and Antimicrobial Resistance Genes

The antimicrobial susceptibility and antimicrobial resistance gene repertoire of the 48 *Enterobacter* strains are shown in Appendix 2 Table 1. *E. xiangfangensis* and taxon 1 had substantially higher rates of resistance to aztreonam (48.3% vs. 10.5%), cefepime (41.4% vs. 10.5%), ceftriaxone (58.6% vs. 15.8%), piperacillin/tazobactam (41.4% vs. 10.5%), and tobramycin (44.8%

Table 2. Proportion of *Enterobacter* species recovered from blood in genomic study of *Enterobacter* bloodstream infection, China

Species*	No. (%)	
	Strains from blood in this study	Strains from blood in GenBank
<i>Enterobacter xiangfangensis</i>	21 (43.8)	221 (63.3)
Taxon 1	8 (16.7)	69 (19.8)
<i>E. bugandensis</i>	4 (8.3)	7 (2.0)
<i>E. cloacae</i>	3 (6.3)	4 (1.1)
<i>E. asburiae</i>	2 (4.2)	14 (4.0)
<i>E. roggkampii</i>	2 (4.2)	7 (2.0)
<i>E. hormaechei</i>	2 (4.2)	0
<i>E. huaxiensis</i>	2 (4.2)	0
<i>E. ludwigii</i>	1 (2.1)	6 (1.7)
<i>E. sichuanensis</i>	1 (2.1)	1 (0.3)
<i>E. chuandaensis</i>	1 (2.1)	0
Taxon 2	1 (2.1)	2 (0.6)
<i>E. kobei</i>	0	10 (2.9)
Taxon 3	0	2 (0.6)
Taxon 4	0	2 (0.6)
Taxon 5	0	1 (0.3)
Taxon 6	0	1 (0.3)
<i>E. chengduensis</i>	0	1 (0.3)
<i>E. mori</i>	0	1 (0.3)
Total	48 (100.0)	349 (100.0)

*Taxons 1–6 represent 6 *Enterobacter* spp. without assigned names. Their most closely related *Enterobacter* spp. are listed in Appendix 1 Table 2 and shown in Appendix 1 Figure 1 (<https://wwwnc.cdc.gov/EID/article/27/1/19-0154-App1.pdf>).

vs. 5.3%) (Table 4) and were substantially more likely to be multidrug resistant (55.2% vs. 10.6%) (Table 3). There were 10 carbapenem-resistant strains (8 *E. xiangfangensis*, 1 *E. bugandensis*, and 1 *E. cloacae*), all of which carried a *bla*_{NDM} gene (*bla*_{NDM-5}, n = 7; *bla*_{NDM-1}, n = 3) (Table 1; Appendix 2 Table 1), and they belonged to 5 STs (ST171 [n = 6], ST718 [n = 1], ST1 [n = 1], ST418 [n = 1], and a new ST [n = 1]). No carbapenemase genes were identified in carbapenem-susceptible strains.

Sequence Types and Clonal Relatedness

The 48 *Enterobacter* strains belonged to 37 STs (Table 1; Appendix 1 Table 3) but only ST171 (*E. xiangfangensis*)

Table 3. Patient demographics, types of bloodstream infection, and outcomes in patients with bloodstream infection caused by *E. xiangfangensis* plus taxon 1 and other *Enterobacter* species, China*

Characteristic	<i>E. xiangfangensis</i> and taxon 1, n = 29	Other species, n = 19	χ ²	p value	Power†
Age, y, median (IQR)	15 (8–32)	52 (16–71)	–	0.958	0.085
Male sex	19 (65.5)	14 (73.7)	0.356	0.551	0.084
MDR	16 (55.2)	2 (10.6)	9.763	0.002	–
Primary BSI	24 (82.8)	12 (63.2)	1.423	0.233	0.317
CLABSI	11 (27.9)	5 (17.2)	0.697	0.404	0.115
HA BSI	27 (93.1)	14 (73.7)	2.091	0.148	0.430
Deaths	6 (20.7)	3 (15.8)	0.176	0.675	0.053
Total time hospitalized, d, median (IQR)‡	33 (19–47)	20 (2–40)	–	0.098	0.265
Time hospitalized before BSI onset, d, median (IQR)‡	11 (5–7)	5 (1–18)	–	0.154	0.070
Time hospitalized after BSI onset, d, median (IQR)	15 (8–32)	13 (8–19)	–	0.357	0.384

*Values are no. (%) except as indicated. Bold indicates significance. BSI, bloodstream infection; CLABSI, central line-associated bloodstream infection; HA BSI, healthcare-associated bloodstream infection; IQR, interquartile range; MDR, multidrug resistance; –, not calculated.

†Statistical power was calculated for parameters without statistical significance (p>0.05).

‡One patient belonging to the other species group had an unusually lengthy hospitalization (578 d) because of a medical dispute and was therefore removed from the analysis of hospitalization time.

and ST78 (taxon 1) contained ≥ 3 strains (6 for ST171 and 3 for ST78). We performed analysis of clonal relatedness based on SNPs for ST78 and ST171 strains. In the 3 ST78 strains, there were 306 to 1,052 SNPs difference, suggesting no recent shared origins (Appendix 1 Table 4). The 6 ST171 strains were all resistant to carbapenems and carried bla_{NDM-5} (n = 5) or bla_{NDM-1} (n = 1). Among the 6 ST171 strains, 4 (all carrying bla_{NDM-5}) had 0–2 SNPs difference (Table 5) and were recovered from patients in the same ward (cardiac surgery). One patient infected with strain 090011 died but the remaining 3 patients recovered.

The remaining 2 ST171 strains, 045001 (carrying bla_{NDM-1}) and 090055 (carrying bla_{NDM-5}), were 81–82 SNPs different from the 4 previously mentioned strains and were 38 SNPs different from each other. These 2 strains were recovered from patients in 2 different wards (medical and respiratory intensive care units). The 6 ST171 strains isolated in our study formed a phylogenetic cluster with strain CCBH10892, which was isolated from a rectal swab sample in Brazil in 2012 (GenBank accession no. JSBO00000000), and strain EC_849, which was isolated from a sputum sample in South Africa in 2012 (GenBank accession no. LRIZ00000000) (Appendix 1 Figure 2). The cluster contained 98 to 107 SNPs difference (Table 5; Appendix 1 Figure 2). Of note, both CCBH10892 and EC_849 carried bla_{NDM-1} . By contrast, the 6 strains were >300 SNPs different from other ST171 strains with genome sequences deposited in GenBank (Appendix 2 Table 3).

Plasmid Analysis of ST171 Strains

The complete genome sequences of bla_{NDM-5} -harboring strain 090011 and the bla_{NDM-1} -harboring strain 045001 were obtained. Strain 090011 had a 4.64-Mb circular chromosome and 2 plasmids (a 102.5-kb

plasmid containing IncFIA, IncFIB, and IncR replicons and a 46.1-kb plasmid containing an IncX3 replicon) (Appendix 1 Table 5). The bla_{NDM-5} gene in strain 090011 was carried on the 46.1-kb IncX3 plasmid, designated pNDM5_090011. The short reads of the remaining 4 bla_{NDM-5} -harboring strains were then mapped against pNDM5_090011. The 4 strains had contigs showing 100% coverage and 100% identity with pNDM5_090011, suggesting a common plasmid in all 5 bla_{NDM-5} -harboring ST171 strains in our study. Strain 045001 had a 4.70-Mb circular chromosome and 3 plasmids (an 85.7-kb IncFII plasmid, a 78.2-kb plasmid, and a 2.5-kb plasmid) (Appendix 1 Table 5). The replicon type of the latter 2 plasmids could not be determined by the current replicon-typing scheme. The bla_{NDM-1} gene in strain 045001 was carried on the 85.7-kb IncFII plasmid.

Discussion

Although genome sequences deposited in GenBank might be biased in sampling, they can provide complementary information on the species distribution of *Enterobacter* in cases of BSI. Examination of our set of strains and those available in GenBank demonstrates that a variety of *Enterobacter* spp. can cause BSI, but most BSI-causing *Enterobacter* strains belong to either *E. xiangfangensis* or, less commonly, taxon 1. *E. xiangfangensis* and taxon 1 are closely related, with a 66.6% isDDH value (near the 70% cutoff to define a species). Why the 2 species account for most *Enterobacter* BSIs, however, remains unknown. Although the colonization of the human gastrointestinal tract by *Enterobacter* has not been investigated to the level of precise species identification, *E. xiangfangensis* and taxon 1 could be the most common *Enterobacter* species colonizing there, which warrants further study.

Table 4. Antimicrobial resistance rates in *E. xiangfangensis* plus taxon 1 and other *Enterobacter* spp. in genomic study of *Enterobacter* bloodstream infection, China*

Antimicrobial agent	No. (%)		χ^2	p value	Power†
	<i>E. xiangfangensis</i> + taxon 1, n = 29	Other species, n = 19			
Amikacin	2 (6.9)	0	–	0.512	ND
Gentamicin	9 (31.0)	1 (5.3)	3.192	0.074	0.600
Tobramycin	13 (44.8)	1 (5.3)	8.698	0.003	–
Aztreonam	14 (48.3)	2 (10.5)	7.361	0.007	0.829
Cefepime	12 (41.4)	2 (10.5)	5.289	0.021	–
Ceftriaxone	17 (58.6)	3 (15.8)	8.664	0.003	–
Imipenem	8 (27.6)	2 (10.5)	1.123	0.289	0.258
Piperacillin/tazobactam	12 (41.4)	2 (10.5)	5.289	0.021	–
Ciprofloxacin	11 (37.9)	0	7.326	0.007	ND
Levofloxacin	11 (37.9)	0	7.326	0.007	ND
Colistin	10 (34.5)	6 (31.6)	0.044	0.835	0.052
Tigecycline	5 (17.2)	0	2.043	0.153	ND
Trimethoprim/sulfamethoxazole	12 (41.4)	0	8.392	0.004	ND

*Bold indicates significance. ND, not determined; –, not calculated.

†Statistical power was calculated for parameters without statistical significance (p>0.05) but could not be calculated for any parameters being 0.

Table 5. Single-nucleotide polymorphisms between the 6 ST171 strains in genomic study of *Enterobacter* bloodstream infection, China, and strain EC_849 from South Africa and strain CCBH10892 from Brazil*

Strain	090011	090022	090023	090059	045001	090055	EC_849	CCBH10892
090011	–	1	0	1	81	81	106	102
090022	1	–	1	2	82	82	107	103
090023	0	1	–	1	81	81	106	102
090059	1	2	1	–	82	82	107	103
045001	81	82	81	82	–	38	103	99
090055	81	82	81	82	38	–	103	99
EC_849	106	107	106	107	103	103	–	98
CCBH10892	102	103	102	103	99	99	98	–

*–, not calculated.

Alternatively, *E. xiangfangensis* and taxon 1 could be more pathogenic than other *Enterobacter* spp., which also requires further study.

BSIs caused by *E. xiangfangensis* and taxon 1 were more likely to occur in younger patients and result in longer overall hospital stays, although the differences in length of hospitalization between the 2 groups were not statistically significant ($p > 0.05$). This finding might be because of the relatively small sample size (power < 0.8) (Table 3). Resistance rates to certain antimicrobial agents, including aztreonam, cefepime, ceftriaxone, piperacillin/tazobactam, and tobramycin and prevalence of multidrug resistance in *E. xiangfangensis* and taxon 1 were substantially higher than in other *Enterobacter* spp. This difference suggests that the identification of *Enterobacter* strains to precise species level also has implications in options of antimicrobial treatment. Although BSI caused by *E. xiangfangensis* and taxon 1 was not associated with higher death rates, 2 of the 4 patients with *E. bugandensis* had poor outcomes (predicted death). *E. bugandensis* has been reported to be a highly pathogenic species associated with life-threatening BSIs and sepsis (19). The virulence of this species warrants further study.

As previously noted, 4 *bla*_{NDM-5}-harboring ST171 strains had 0–2 SNPs difference (Table 5) and were taken from patients in the same ward (cardiac surgery). The first patient with BSI caused by a strain belonging to the clone (090011) had *Enterobacter* BSI before being transferred to West China Hospital from another facility. Although the particular *Enterobacter* strain from the first hospital was not available for analysis, it is very likely that 090011 was introduced to West China Hospital by transfer of this patient. The next 3 cases were acquired in West China Hospital, highlighting both interhospital and intrahospital transmission of a common strain. BSI in 2 of the 3 cases was CLABSI. These findings suggest that the 4 strains belong to a common clone that caused a cluster of BSI cases. In the hospital ward, central lines were commonly used for drawing blood but were not properly decontaminated after each use; in addition,

healthcare workers were observed by anonymous interns to have low compliance (24.8%) with hand hygiene standards established by the World Health Organization. There have been no further *Enterobacter* BSIs after restricting access to central lines and promoting hand hygiene among healthcare workers, which resulted in the compliance rate increasing to 45.5%.

The remaining 2 ST171 strains, 045001 (harboring *bla*_{NDM-1}) and 090055 (harboring *bla*_{NDM-5}), belong to 2 clones (38 SNPs between each other) which differed from the aforementioned clone by 81–82 SNPs. The relatively low number of SNPs among the 3 ST171 clones also suggests recent divergence within the lineage. In addition, the 6 ST171 strains in this study were clustered together with 2 *bla*_{NDM-1}-harboring strains, strain CCBH10892 isolated from Brazil in 2012, and strain EC_849 from South Africa in 2012, with 98 to 107 SNPs, but had > 300 SNPs with other ST171 strains that had genome sequences deposited in GenBank. This finding suggests that the 6 strains identified in this study, CCBH10892, and EC_849 represent a subclade of ST171, which carries *bla*_{NDM'} has an international distribution, and might have emerged within the past 10 years. In addition, although 090055 and the other 4 *bla*_{NDM-5}-harboring strains (090011, 090022, 090023, and 090059) belonged to 2 different clones, they had the same IncX3 plasmid carrying *bla*_{NDM-5'}, which suggests that the spread of *bla*_{NDM-5} in the hospital was both clonal (vertical) and plasmidborne (horizontal).

The association of ST171 *E. xiangfangensis* with outbreaks is not rare; repeated reports from different geographic locations have demonstrated the same association (20–24). This association suggests that ST171 is a lineage of *Enterobacter*, which might be well adapted to causing infections in healthcare settings. Previous reports have demonstrated that ST171 *E. xiangfangensis* is a high-risk clone mediating the spread of carbapenem resistance (21,23). Its emergence was initially documented in 2015 and 2016 by 2 studies in the United States (24,25). Subsequent

studies have revealed the international distribution of ST171 (21–23). Almost all carbapenem-resistant ST171 *E. xiangfangensis* strains have *bla*_{KPC} (21), and only a small number of strains carry *bla*_{NDM} instead (21,26). In this study, we identified in-hospital transmission of carbapenem-resistant ST171 strains, which carried *bla*_{NDM-5} rather than *bla*_{KPC} as seen in previous studies (21,23). The ability to acquire different carbapenemase genes and its adaptability to healthcare settings might be major drivers in the emergence of ST171, which warrants further study.

Our investigation demonstrates the value of whole-genome sequencing for precise species identification. However, this study has several limitations. First, because it is a single site study, the application of our findings could be limited. However, we analyzed genomes available in GenBank to provide the most comprehensive information possible. Second, the relatively small sample size in this study might not have adequate power to examine statistical significance in BSI type and patient outcomes. However, this study provided useful information on the clinical importance of *E. xiangfangensis* and its closely related taxon 1. Larger-scale studies are warranted.

In conclusion, most *Enterobacter* strains recovered from human blood in China were not *E. cloacae* but *E. xiangfangensis*. Most *Enterobacter* BSI cases in our study were healthcare-associated and primary infections. *E. xiangfangensis* ST171 is a major lineage of carbapenem-resistant *Enterobacter*, has an intercontinental distribution, is usually healthcare associated, and carries *bla*_{NDM} rather than *bla*_{KPC} in China. Precise species identification of *Enterobacter* has clinical importance in antimicrobial therapy and infection control.

Addendum: Since submission and acceptance of this manuscript, the taxonomy of *Enterobacter* has been substantially updated. The updated *Enterobacter* taxonomy is available at <https://doi.org/10.1128/mSystems.00527-20>. The update of new taxa identified in this study is shown in Appendix 3 Table (<https://wwwnc.cdc.gov/EID/article/27/1/19-0154-App3.pdf>).

Acknowledgments

We are grateful for Alan McNally for his proofreading and helpful suggestions.

The work was supported by grants from the National Natural Science Foundation of China (project no. 81772233, 81661130159 and 81861138055), West China Hospital of Sichuan University (1.3.5 project for disciplines of excellence, project no. ZYYC08006) and the Newton Advanced Fellowship, Royal Society, UK (NA150363).

About the Author

Dr. Wu is a scientist in the Department of Laboratory Medicine, West China Second Hospital, Sichuan University. Her major research interests are the epidemiology and clonal background of carbapenemase-producing *Enterobacter* spp.

References

1. Mezzatesta ML, Gona F, Stefani S. *Enterobacter cloacae* complex: clinical impact and emerging antibiotic resistance. *Future Microbiol.* 2012;7:887–902. <https://doi.org/10.2217/fmb.12.61>
2. Tindall BJ, Sutton G, Garrity GM. *Enterobacter aerogenes* Hormaeche and Edwards 1960 (approved lists 1980) and *Klebsiella mobilis* Bascomb et al. 1971 (approved lists 1980) share the same nomenclatural type (ATCC 13048) on the approved lists and are homotypic synonyms, with consequences for the name *Klebsiella mobilis* Bascomb et al. 1971 (approved lists 1980). *Int J Syst Evol Microbiol.* 2017;67:502–4. <https://doi.org/10.1099/ijsem.0.001572>
3. Gavini R, Mergaert J, Beji A, Mielcarek C, Izard D, Kersters K, et al. Transfer of *Enterobacter agglomerans* (Beijerinck 1888) Ewing and Fife 1972 to *Pantoea* gen. nov. as *Pantoea agglomerans* comb. nov. and description of *Pantoea dispersa* sp. nov. *Int J Syst Evol Microbiol.* 1989;39:337–45.
4. Iversen C, Mullane N, McCardell B, Tall BD, Lehner A, Fanning S, et al. *Cronobacter* gen. nov., a new genus to accommodate the biogroups of *Enterobacter sakazakii*, and proposal of *Cronobacter sakazakii* gen. nov., comb. nov., *Cronobacter malonicus* sp. nov., *Cronobacter turicensis* sp. nov., *Cronobacter muytjensii* sp. nov., *Cronobacter dublinensis* sp. nov., *Cronobacter* genomospecies 1, and of three subspecies, *Cronobacter dublinensis* subsp. *dublinensis* subsp. nov., *Cronobacter dublinensis* subsp. *lausannensis* subsp. nov. and *Cronobacter dublinensis* subsp. *lactaridi* subsp. nov. *Int J Syst Evol Microbiol.* 2008;58:1442–7. <https://doi.org/10.1099/ijms.0.65577-0>
5. Hoffmann H, Roggenkamp A. Population genetics of the nomenspecies *Enterobacter cloacae*. *Appl Environ Microbiol.* 2003;69:5306–18. <https://doi.org/10.1128/AEM.69.9.5306-5318.2003>
6. Hoffmann H, Stindl S, Stumpf A, Mehlen A, Monget D, Heesemann J, et al. Description of *Enterobacter ludwigii* sp. nov., a novel *Enterobacter* species of clinical relevance. *Syst Appl Microbiol.* 2005;28:206–12. <https://doi.org/10.1016/j.syapm.2004.12.009>
7. Wu W, Feng Y, Zong Z. *Enterobacter sichuanensis* sp. nov., recovered from human urine. *Int J Syst Evol Microbiol.* 2018;68:3922–7. <https://doi.org/10.1099/ijsem.0.003089>
8. Balloux F, Brønstad Brynildsrud O, van Dorp L, Shaw LP, Chen H, Harris KA, et al. From theory to practice: translating whole-genome sequencing (WGS) into the clinic. *Trends Microbiol.* 2018;26:1035–48. <https://doi.org/10.1016/j.tim.2018.08.004>
9. Besser J, Carleton HA, Gerner-Smidt P, Lindsey RL, Trees E. Next-generation sequencing technologies and their application to the study and control of bacterial infections. *Clin Microbiol Infect.* 2018;24:335–41. <https://doi.org/10.1016/j.cmi.2017.10.013>
10. Rosselló-Móra R, Amann R. Past and future species definitions for *Bacteria* and *Archaea*. *Syst Appl Microbiol.* 2015;38:209–16. <https://doi.org/10.1016/j.syapm.2015.02.001>

11. Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing; twenty-seventh informational supplement. M100-S27. Wayne (PA): The Institute; 2017.
12. Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, et al. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect*. 2012;18:268–81. <https://doi.org/10.1111/j.1469-0691.2011.03570.x>
13. Centers for Disease Control and Prevention. National Healthcare Safety Network FAQs: bloodstream infection (BSI) events. 2018 [cited 2018 Nov 1]. <https://www.cdc.gov/nhsn/faqs/faq-bsi.html>
14. Centers for Disease Control and Prevention. Identifying healthcare-associated infections (HAI) for NHSN surveillance. 2018 [cited 2018 Nov 1]. https://www.cdc.gov/nhsn/pdfs/pscmanual/2psc_identifyinghais_nhsncurrent.pdf
15. Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol*. 2017;13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>
16. Meier-Kolthoff JP, Auch AF, Klenk HP, Göker M. Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC Bioinformatics*. 2013;14:60. <https://doi.org/10.1186/1471-2105-14-60>
17. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*. 2014;30:1312–3. <https://doi.org/10.1093/bioinformatics/btu033>
18. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al. Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS One*. 2014;9:e112963. <https://doi.org/10.1371/journal.pone.0112963>
19. Pati NB, Dojjad SP, Schultze T, Mannala GK, Yao Y, Jaiswal S, et al. *Enterobacter bugandensis*: a novel enterobacterial species associated with severe clinical infection. *Sci Rep*. 2018;8:5392. <https://doi.org/10.1038/s41598-018-23069-z>
20. Hawken SE, Washer LL, Williams CL, Newton DW, Snitkin ES. Genomic investigation of a putative endoscope-associated carbapenem-resistant *Enterobacter cloacae* outbreak reveals a wide diversity of circulating strains and resistance mutations. *Clin Infect Dis*. 2018;66:460–3. <https://doi.org/10.1093/cid/cix934>
21. Chavda KD, Chen L, Fouts DE, Sutton G, Brinkac L, Jenkins SG, et al. Comprehensive genome analysis of carbapenemase-producing *Enterobacter* spp.: new insights into phylogeny, population structure, and resistance mechanisms. *MBio*. 2016;7:02093–16. <https://doi.org/10.1128/mBio.02093-16>
22. Harada K, Shimizu T, Mukai Y, Kuwajima K, Sato T, Kajino A, et al. Phenotypic and molecular characterization of antimicrobial resistance in *Enterobacter* spp. isolates from companion animals in Japan. *PLoS One*. 2017;12:e0174178. <https://doi.org/10.1371/journal.pone.0174178>
23. Gomez-Simmonds A, Annavaiah MK, Wang Z, Macesic N, Hu Y, Giddins MJ, et al. Genomic and geographic context for the evolution of high-risk carbapenem-resistant *Enterobacter cloacae* complex clones ST171 and ST78. *MBio*. 2018;9:00542–18. <https://doi.org/10.1128/mBio.00542-18>
24. Hargreaves ML, Shaw KM, Dobbins G, Snippes Vagnone PM, Harper JE, Boxrud D, et al. Clonal dissemination of *Enterobacter cloacae* harboring *bla*_{KPC-3} in the upper midwestern United States. *Antimicrob Agents Chemother*. 2015;59:7723–34. <https://doi.org/10.1128/AAC.01291-15>
25. Gomez-Simmonds A, Hu Y, Sullivan SB, Wang Z, Whittier S, Uhlemann AC. Evidence from a New York City hospital of rising incidence of genetically diverse carbapenem-resistant *Enterobacter cloacae* and dominance of ST171, 2007–14. *J Antimicrob Chemother*. 2016;71:2351–3. <https://doi.org/10.1093/jac/dkw132>
26. Peirano G, Matsumura Y, Adams MD, Bradford P, Motyl M, Chen L, et al. Genomic epidemiology of global carbapenemase-producing *Enterobacter* spp., 2008–2014. *Emerg Infect Dis*. 2018;24:1010–9. <https://doi.org/10.3201/eid2406.171648>

Address for correspondence: Zhiyong Zong, Center of Infectious Diseases, West China Hospital (Huaxi), Guoxuexiang 37, Chengdu 610041, China; email: zongzhiy@scu.edu.cn

Precise Species Identification by Whole-Genome Sequencing of *Enterobacter* Bloodstream Infection, China

Appendix 1

Appendix 1 Table 1. Classification and nomenclature of the genus *Enterobacter* as of December 2018*

Species	Hoffman cluster	Reference	Type strain	GenBank accession no. or current species name
Species name for validation (n = 14)				
<i>Enterobacter asburiae</i>	I	(1)	JCM 6051	CP011863
<i>E. cancerogenus</i>	-	(2)	ATCC 35316	ERR1854846
<i>E. chuandaensis</i>	-	(3)	090028 ^T	QZCS00000000
<i>E. cloacae</i>	XI		ATCC 13047	ERR1854846
<i>E. bugandensis</i>	IX	(4)	EB-247	FYBI00000000
<i>E. hormaechei</i>	VII	(5)	ATCC 49162	MKEQ00000000
<i>E. huaxiensis</i>	-	(3)	090008 ^T	QZCT00000000
<i>E. kobei</i>	II	(6)	ATCC BAA-260	CP017181
<i>E. ludwigii</i>	V	(7)	EN-119	CP017279
<i>E. mori</i>	-	(8)	LMG 25706	AEXB00000000
<i>E. soli</i>	-	(9)	ATCC BAA-2102	LXES00000000
<i>E. tabaci</i>	-	(10)	YIM Hb-3	N/A
<i>E. xiangfangensis</i> [†]	VI	(11)	LMG 27195	CP017183
<i>E. sichuanensis</i>	-	(12)	WCHECI1597	POVL00000000
Species in doubt (n = 2)				
<i>E. muelleri</i> [‡]	-	(13)	JM-458	FXLQ00000000
<i>E. siamensis</i> [§]	-	(14)	C2361	N/A
Species name awaiting validation (n = 3)				
<i>E. timonensis</i>	-	(15)	mt20	FCOP00000000
<i>E. chengduensis</i>	-	(16)	WCHECI-C4	MTSO00000000
<i>E. roggenkampii</i>	IV	(17)	DSM16690	CP017184
Species listed in LPSN but moved out of <i>E.</i> (n = 20)				

Species	Hoffman			GenBank accession no.
	cluster	Reference	Type strain	or current species name
<i>E. aerogenes</i>		(18)	ATCC 13048	<i>Klebsiella aerogenes</i>
<i>E. agglomerans</i>		(19)	ATCC 27155	<i>Pantoea agglomerans</i>
<i>E. amnigenus</i>		(20)	ATCC 33072	<i>Lelliottia amnigena</i>
<i>E. arachidis</i>		(20)	KCTC 22375	<i>Kosakonia arachidis</i>
<i>E. cowanii</i>		(20)	CCUG 45998	<i>Kosakonia cowanii</i>
<i>E. gergoviae</i>		(20)	ATCC 33028	<i>Pluralibacter gergoviae</i>
<i>E. helveticus</i>		(20)	JCM 16470	<i>Cronobacter helveticus</i>
<i>E. intermedius</i>		(21)	ATCC 33110	<i>Kluyvera intermedia</i>
<i>E. massiliensis</i>		(22)	JC163	<i>Metakosakonia massiliensis</i>
<i>E. nimipressuralis</i>	X	(20)	CIP 104980	<i>Lelliottia nimipressuralis</i>
<i>E. oryzae</i>		(20)	LMG 24251	<i>Kosakonia oryzae</i>
<i>E. oryzendophyticus</i>		(23)	LMG 26432	<i>Kosakonia oryzendophytica</i>
<i>E. oryziphilus</i>		(23)	LMG 26429	<i>Kosakonia oryziphila</i>
<i>E. pulveris</i>		(20)	DSM 19144	<i>Cronobacter pulveris</i>
<i>E. pyrinus</i>		(20)	ATCC 49851	<i>Pluralibacter pyrinus</i>
<i>E. radicincitans</i>		(20)	CIP 108468	<i>Kosakonia radicincitans</i>
<i>E. sacchari</i>		(11)	CGMCC 1.12102	<i>Kosakonia sacchari</i>
<i>E. sakazakii</i>		(24)	ATCC 29544	<i>Cronobacter sakazakii</i>
<i>E. taylorae</i>		(25)	ATCC 35317	<i>Enterobacter cancerogenus</i>
<i>E. turicensis</i>		(20)	DSM 18397	<i>Cronobacter zurichensis</i>

*LPSN, The list of Prokaryotic Names with Standing in Nomenclature.

†The species status of *E. xiangfangensis* has been doubted previously and it has been proposed as a subspecies of *E. hormaechei* rather than a valid species (17,26). However, its type strain has only 94.48% ANI and 60.0% isDDH with *E. hormaechei* type strain ATCC 49162^T (GenBank accession no. MKEQ00000000). Therefore, *E. xiangfangensis* and *E. hormaechei* are clearly 2 different species.

‡*E. muelleri* is a later synonym of *E. asburiae*.

§It has been proposed to reject *E. siamensis* because the 16S rRNA sequence of its type strain available in collections does not match its record in GenBank (27).

Appendix 1 Table 2. The 6 unnamed *Enterobacter* spp. identified in genomic study of *Enterobacter* bloodstream infection, China*

Species				
assignment	Representative strain	Genome accession no.	Closest species	isDDH (%)†
Taxon 1	DSM 14563‡	CP017186	<i>E. xiangfangensis</i>	66.6
Taxon 2	e362	FKDT00000000	<i>E. roggenkampii</i>	65.4
Taxon 3	e773	FKGE00000000	<i>E. asburiae</i>	65.3
Taxon 4	e2032	FKBK00000000	<i>E. asburiae</i>	52.0
Taxon 5	e483	FKEG01000000	<i>E. asburiae</i>	49.6
Taxon 6	153C2	QMCQ01000000	<i>E. xiangfangensis</i>	52.8

*isDDH, in silico DNA–DNA hybridization.

†isDDH values between the representative strain and the type strain of closest species.

‡Strain DSM 14563 has been proposed as the type strain of *E. hormaechei* subspecies *Hoffmannii*. However, the strain has only 94.13% ANI and 58.0% isDDH with *E. hormaechei* type strain ATCC 49162T (GenBank accession no. MKEQ00000000). It is clear that the *E. hormaechei* subspecies *hoffmannii* is actually not a subspecies of *E. hormaechei* but rather represents a new, unnamed *Enterobacter* species. In this study, we temporarily designated the species taxon 1 for simplicity.

Appendix 1 Table 3. Profiles of sequence types in genomic study of *Enterobacter* bloodstream infection, China*

ST	Species	<i>dnaA</i>	<i>fusA</i>	<i>gyrB</i>	<i>leuS</i>	<i>pyrG</i>	<i>rplB</i>	<i>rpoB</i>	Closest ST†
									(no. of allele difference)
1	<i>E. cloacae</i>	1	1	1	1	1	1	1	
12	<i>E. ludwigii</i>	13	2	45	24	52	2	14	
50	<i>E. xiangfangensis</i>	4	4	4	6	37	4	25	
78	Taxon 1	8	9	6	9	9	6	8	
97	Taxon 1	59	9	62	9	62	25	6	
104	Taxon 1	59	40	76	9	70	6	6	
127	<i>E. xiangfangensis</i>	46	20	74	44	45	24	6	
171	<i>E. xiangfangensis</i>	49	21	19	44	45	12	32	
316	Taxon 1	59	88	82	9	67	6	6	
337	<i>E. xiangfangensis</i>	67	21	9	129	45	12	32	
418	<i>E. xiangfangensis</i>	53	35	154	44	45	4	6	
499	<i>E. bugandensis</i>	164	18	183	200	120	8	29	
519	<i>E. cloacae</i>	1	107	158	1	168	36	1	
528	<i>E. hormaechei</i>	95	56	112	116	104	4	63	
550	<i>E. xiangfangensis</i>	179	4	4	6	112	4	6	
568	Taxon 1	189	9	12	9	67	6	6	
696	<i>E. hormaechei</i>	225	140	93	268	224	109	141	
718	<i>E. bugandensis</i>	140	18	248	31	230	8	29	
828	<i>E. xiangfangensis</i>	9	4	14	61	257	4	9	
879	<i>E. asburiae</i>	152	15	102	15	101	11	133	

ST	Species	<i>dnaA</i>	<i>fusA</i>	<i>gyrB</i>	<i>leuS</i>	<i>pyrG</i>	<i>rplB</i>	<i>rpoB</i>	Closest ST† (no. of allele difference)
922	<i>E. cloacae</i>	169	107	61	168	36	77	1	
984	<i>E. roggenkampii</i>	65	57	49	94	49	12	47	
N1	<i>E. huaxiensis</i>	n1	n1	n1	n1	n1	n1	n1	None
N2	<i>E. roggenkampii</i>	191	n2	254	193	49	12	26	613 (3)
N3	<i>E. xiangfangensis</i>	n2	20	148	44	45	4	6	886/916/986 (1)
N4	<i>E. xiangfangensis</i>	4	4	15	4	11	30	6	111/981 (2)
N5	<i>E. xiangfangensis</i>	58	22	14	6	39	4	9	79 (1)
N6	<i>E. xiangfangensis</i>	58	41	14	6	69	4	n2	106 (1)
N7	<i>E. xiangfangensis</i>	178	4	4	6	92	4	6	542 (1)
N8	<i>E. xiangfangensis</i>	4	37	4	6	42	4	6	329 (1)
N9	<i>E. chuandaensis</i>	n3	n3	n2	n2	n2	n2	120	573/944 (6)
N10	<i>E. bugandensis</i>	309	18	n3	n3	34	8	n3	1084 (3)
N11	<i>E. sichuanensis</i>	n4	98	170	n4	n3	68	n4	472/607/738/847 (4)
N12	<i>E. asburiae</i>	n5	15	n4	124	n4	11	68	319 (3)
N13	Taxon 1	59	9	n5	n5	79	37	n5	157/419/792 (4)
N14	Taxon 2	151	108	n6	n6	n5	14	93	474 (3)
N15	<i>E. xiangfangensis</i>	n6	69	19	44	64	4	32	270 (2)
N16	<i>E. bugandensis</i>	140	18	n7	31	230	8	29	718 (1)

*N1 to N16 are new sequence types. New alleles are temporarily assigned n1 to n6. ST, sequence type.

†For new sequence types only.

Appendix 1 Table 4. Single nucleotide polymorphisms between the 3 ST78 strains in genomic study of *Enterobacter* bloodstream infection, China*

Strain	090039	090030	090019
090039	–	1,052	814
090030	1,052	-	306
090019	814	306	–

*The 3 genomes were mapped against the complete chromosome

sequence of ST78 strain AR_0050 (GenBank accession no. CP021896)

by using Parsnp version 1.2 and alignment was obtained by using Harvest

(28).

Appendix 1 Table 5. Complete genome and antimicrobial resistance genes of strain 090011 and strain 045001 in genomic study of *Enterobacter* bloodstream infection, China*

Strain	Size, bp	Replicon type, Inc	Genes mediating resistance to							
			β -lactam	Aminoglycoside	Fluoroquinolone	Fosfomycin	Rifampin	Sulfonamid e	Tetracycline	Trimethoprim
090011										
Chromosome	4,639,926	-	<i>bla</i> _{ACT-7}				<i>fosA</i>			
pNDM5_090011	46,161	X3	<i>bla</i> _{NDM-5}							
pCTXM65_090011	102,543	FIA, FIB, R	<i>bla</i> _{CTX-M-65} , <i>bla</i> _{TEM-1B}	<i>aadA16</i> , <i>aac(6')-Ib-cr</i>	<i>aac(6')-Ib-cr</i>			<i>arr-3</i>	<i>sul1</i>	<i>tet(A)</i> <i>dfrA27</i>
045001										
Chromosome	4,698,270	-	<i>bla</i> _{CTX-M-15} , <i>bla</i> _{OXA-1} , <i>bla</i> _{ACT-7}	<i>aac(3)-IIa</i> , <i>aac(6')-Ib-cr</i>	<i>aac(6')-Ib-cr</i> , <i>qnrB1</i>		<i>fosA</i>			
pNDM1_045001	85,718	FII	<i>bla</i> _{NDM-1} , <i>bla</i> _{TEM-1}	<i>rmtB</i>						<i>tet(A)</i> <i>dfrA14</i>
p1_045001	78,247	ND								
p2_045001	2,496	ND								

*ND, undetermined.

References


1. Brenner DJ, McWhorter AC, Kai A, Steigerwalt AG, Farmer JJ III. *Enterobacter asburiae* sp. nov., a new species found in clinical specimens, and reassignment of *Erwinia dissolvens* and *Erwinia nimipressuralis* to the genus *Enterobacter* as *Enterobacter dissolvens* comb. nov. and *Enterobacter nimipressuralis* comb. nov. J Clin Microbiol. 1986;23:1114–20. [PubMed](#)
<https://doi.org/10.1128/JCM.23.6.1114-1120.1986>
2. Dickey RS, Zumoff CH. Emended description of *Enterobacter cancerogenus* comb. nov. (formerly *Erwinia cancerogena*). Int J Syst Bacteriol. 1988;38:371–4. <https://doi.org/10.1099/00207713-38-4-371>
3. Wu W, Wei L, Feng Y, Kang M, Zong Z. *Enterobacter huaxiensis* sp. nov. and *Enterobacter chuandaensis* sp. nov. recovered from human blood. Int J Syst Evol Microbiol. 2018;69:708–14 In press. [PubMed](#) <https://doi.org/10.1099/ijsem.0.003207>
4. Doijad S, Imirzalioglu C, Yao Y, Pati NB, Falgenhauer L, Hain T, et al. *Enterobacter bugandensis* sp. nov., isolated from neonatal blood. Int J Syst Evol Microbiol. 2016;66:968–74. [PubMed](#)
<https://doi.org/10.1099/ijsem.0.000821>
5. O'Hara CM, Steigerwalt AG, Hill BC, Farmer JJ III, Fanning GR, Brenner DJ. *Enterobacter hormaechei*, a new species of the family *Enterobacteriaceae* formerly known as enteric group 75. J Clin Microbiol. 1989;27:2046–9. [PubMed](#) <https://doi.org/10.1128/JCM.27.9.2046-2049.1989>
6. Kosako Y, Tamura K, Sakazaki R, Miki K. *Enterobacter kobei* sp. nov., a new species of the family *Enterobacteriaceae* resembling *Enterobacter cloacae*. Curr Microbiol. 1996;33:261–5. [PubMed](#)
<https://doi.org/10.1007/s002849900110>
7. Hoffmann H, Stindl S, Stumpf A, Mehlen A, Monget D, Heesemann J, et al. Description of *Enterobacter ludwigii* sp. nov., a novel *Enterobacter* species of clinical relevance. Syst Appl Microbiol. 2005;28:206–12. [PubMed](#) <https://doi.org/10.1016/j.syapm.2004.12.009>

8. Zhu B, Lou MM, Xie GL, Wang GF, Zhou Q, Wang F, et al. *Enterobacter mori* sp. nov., associated with bacterial wilt on *Morus alba* L. Int J Syst Evol Microbiol. 2011;61:2769–74. [PubMed](#)
<https://doi.org/10.1099/ijs.0.028613-0>
9. Manter DK, Hunter WJ, Vivanco JM. *Enterobacter soli* sp. nov.: a lignin-degrading γ -proteobacteria isolated from soil. Curr Microbiol. 2011;62:1044–9. [PubMed](#) <https://doi.org/10.1007/s00284-010-9809-9>
10. Duan YQ, Zhou XK, Di-Yan L, Li QQ, Dang LZ, Zhang YG, et al. *Enterobacter tabaci* sp. nov., a novel member of the genus *Enterobacter* isolated from a tobacco stem. Antonie van Leeuwenhoek. 2015;108:1161–9. [PubMed](#) <https://doi.org/10.1007/s10482-015-0569-1>
11. Gu CT, Li CY, Yang LJ, Huo GC. *Enterobacter xiangfangensis* sp. nov., isolated from Chinese traditional sourdough, and reclassification of *Enterobacter sacchari* Zhu et al. 2013 as *Kosakonia sacchari* comb. nov. Int J Syst Evol Microbiol. 2014;64:2650–6. [PubMed](#)
<https://doi.org/10.1099/ijs.0.064709-0>
12. Wu W, Feng Y, Zong Z. *Enterobacter sichuanensis* sp. nov., recovered from human urine. Int J Syst Evol Microbiol. 2018;68:3922–7. [PubMed](#) <https://doi.org/10.1099/ijsem.0.003089>
13. Kämpfer P, McInroy JA, Glaeser SP. *Enterobacter muelleri* sp. nov., isolated from the rhizosphere of *Zea mays*. Int J Syst Evol Microbiol. 2015;65:4093–9. [PubMed](#)
<https://doi.org/10.1099/ijsem.0.000547>
14. Khunthongpan S, Bourneow C, H-Kittikun A, Tanasupawat S, Benjakul S, Sumpavapol P. *Enterobacter siamensis* sp. nov., a transglutaminase-producing bacterium isolated from seafood processing wastewater in Thailand. J Gen Appl Microbiol. 2013;59:135–40. [PubMed](#)
<https://doi.org/10.2323/jgam.59.135>
15. Tidjani Alou M, Cadoret F, Brah S, Diallo A, Sokhna C, Mehrej V, et al. ‘*Khelaifiella massiliensis*’, ‘*Niameybacter massiliensis*’, ‘*Brachybacterium massiliense*’, ‘*Enterobacter timonensis*’, ‘*Massilibacillus massiliensis*’, new bacterial species and genera isolated from the gut microbiota


- of healthy infants. *New Microbes New Infect.* 2017;19:1–7. [PubMed](#)
<https://doi.org/10.1016/j.nmni.2017.02.002>
16. Wu W, Feng Y, Zong Z. Characterization of a strain representing a new *Enterobacter* species, *Enterobacter chengduensis* sp. nov. *Antonie van Leeuwenhoek.* 2019;112:491–500. [PubMed](#)
<https://doi.org/10.1007/s10482-018-1180-z>
17. Granger G, Sutton, Lauren M, Brinkac, Thomas H, Clarke, Fouts DE. *Enterobacter hormaechei* subsp. *hoffmannii* subsp. nov., *Enterobacter hormaechei* subsp. *xiangfangensis* comb. nov., *Enterobacter roggenkampii* sp. nov., and *Enterobacter muelleri* is a later heterotypic synonym of *Enterobacter asburiae* based on computational analysis of sequenced *Enterobacter* genomes. *F1000 Res.* 2018;7:521. <https://doi.org/10.12688/f1000research.14566.1>
18. Tindall BJ, Sutton G, Garrity GM. *Enterobacter aerogenes* Hormaeche and Edwards 1960 (approved lists 1980) and *Klebsiella mobilis* Bascomb et al. 1971 (approved lists 1980) share the same nomenclatural type (ATCC 13048) on the approved lists and are homotypic synonyms, with consequences for the name *Klebsiella mobilis* Bascomb et al. 1971 (approved lists 1980). *Int J Syst Evol Microbiol.* 2017;67:502–4. [PubMed](#) <https://doi.org/10.1099/ijsem.0.001572>
19. Edwards J, Baque E, Tang J. Proposal to acknowledge Beijerinck as the original author of the species *Pantoea agglomerans*. Request for an opinion. *Int J Syst Evol Microbiol.* 2004;54:2437. [PubMed](#)
<https://doi.org/10.1099/ijms.0.02955-0>
20. Brady C, Cleenwerck I, Venter S, Coutinho T, De Vos P. Taxonomic evaluation of the genus *Enterobacter* based on multilocus sequence analysis (MLSA): proposal to reclassify *E. nimipressuralis* and *E. amnigenus* into *Lelliottia* gen. nov. as *Lelliottia nimipressuralis* comb. nov. and *Lelliottia amnigena* comb. nov., respectively, *E. gergoviae* and *E. pyrinus* into *Pluralibacter* gen. nov. as *Pluralibacter gergoviae* comb. nov. and *Pluralibacter pyrinus* comb. nov., respectively, *E. cowanii*, *E. radicincitans*, *E. oryzae* and *E. arachidis* into *Kosakonia* gen. nov. as *Kosakonia cowanii* comb. nov., *Kosakonia radicincitans* comb. nov., *Kosakonia oryzae*

- comb. nov. and *Kosakonia arachidis* comb. nov., respectively, and *E. turicensis*, *E. helveticus* and *E. pulveris* into *Cronobacter* as *Cronobacter zurichensis* nom. nov., *Cronobacter helveticus* comb. nov. and *Cronobacter pulveris* comb. nov., respectively, and emended description of the genera *Enterobacter* and *Cronobacter*. Syst Appl Microbiol. 2013;36:309–19. [PubMed](#)
<https://doi.org/10.1016/j.syapm.2013.03.005>
21. Pavan ME, Franco RJ, Rodriguez JM, Gadaleta P, Abbott SL, Janda JM, et al. Phylogenetic relationships of the genus *Kluyvera*: transfer of *Enterobacter intermedius* Izard et al. 1980 to the genus *Kluyvera* as *Kluyvera intermedia* comb. nov. and reclassification of *Kluyvera cochleae* as a later synonym of *K. intermedia*. Int J Syst Evol Microbiol. 2005;55:437–42. [PubMed](#)
<https://doi.org/10.1099/ijs.0.63071-0>
22. Alnajar S, Gupta RS. Phylogenomics and comparative genomic studies delineate six main clades within the family *Enterobacteriaceae* and support the reclassification of several polyphyletic members of the family. Infect Genet Evol. 2017;54:108–27. [PubMed](#)
<https://doi.org/10.1016/j.meegid.2017.06.024>
23. Di Pilato V, Arena F, Tascini C, Cannatelli A, Henrici De Angelis L, Fortunato S, et al. *mcr-1.2*, a new *mcr* variant carried on a transferable plasmid from a colistin-resistant KPC carbapenemase-producing *Klebsiella pneumoniae* strain of sequence type 512. Antimicrob Agents Chemother. 2016;60:5612–5. [PubMed](#) <https://doi.org/10.1128/AAC.01075-16>
24. Iversen C, Lehner A, Mullane N, Bidlas E, Cleenwerck I, Marugg J, et al. The taxonomy of *Enterobacter sakazakii*: proposal of a new genus *Cronobacter* gen. nov. and descriptions of *Cronobacter sakazakii* comb. nov. *Cronobacter sakazakii* subsp. *sakazakii*, comb. nov., *Cronobacter sakazakii* subsp. *malonaticus* subsp. nov., *Cronobacter turicensis* sp. nov., *Cronobacter muytjensii* sp. nov., *Cronobacter dublinensis* sp. nov. and *Cronobacter* genomospecies 1. BMC Evol Biol. 2007;7:64. [PubMed](#) <https://doi.org/10.1186/1471-2148-7-64>

25. Schönheyder HC, Jensen KT, Frederiksen W. Taxonomic notes: synonymy of *Enterobacter cancerogenus* (Urosević 1966) Dickey and Zumoff 1988 and *Enterobacter taylorae* Farmer et al. 1985 and resolution of an ambiguity in the biochemical profile. *Int J Syst Bacteriol.* 1994;44:586–7. [PubMed https://doi.org/10.1099/00207713-44-3-586](https://doi.org/10.1099/00207713-44-3-586)
26. Chavda KD, Chen L, Fouts DE, Sutton G, Brinkac L, Jenkins SG, et al. Comprehensive genome analysis of carbapenemase-producing *Enterobacter* spp.: new insights into phylogeny, population structure, and resistance mechanisms. *MBio.* 2016;7:02093–16. [PubMed https://doi.org/10.1128/mBio.02093-16](https://doi.org/10.1128/mBio.02093-16)
27. Kämpfer P, Dojjad S, Chakraborty T, Glaeser SP. The status of the species *Enterobacter siamensis* Khunthongpan et al. 2014. Request for an Opinion. *Int J Syst Evol Microbiol.* 2016;66:524–5. [PubMed https://doi.org/10.1099/ijsem.0.000777](https://doi.org/10.1099/ijsem.0.000777)
28. Treangen TJ, Ondov BD, Koren S, Phillippy AM. The Harvest suite for rapid core-genome alignment and visualization of thousands of intraspecific microbial genomes. *Genome Biol.* 2014;15:524. [PubMed https://doi.org/10.1186/s13059-014-0524-x](https://doi.org/10.1186/s13059-014-0524-x)



Appendix 1 Figure 1. Phylogenomic tree of *Enterobacter* species. Phylogenomic tree based on the concatenated nucleotide sequence of the core-genes of type strains of species belonging to the genus *Enterobacter* (Appendix 1 Table 1) and representative strains of the 6 new species (Appendix 1 Table 2) as described previously (12). The species and strain name are shown and accession numbers are indicated in parentheses. The tree was inferred by using the maximum-likelihood algorithm. Bootstrap values >50% (based on 1,000 resamplings) are indicated by the dotted lines. Scale bar indicates 0.4 nt substitution per site.



Appendix 1 Figure 2. Phylogenomic tree of ST171 *Enterobacter* strains. Strains in this study are highlighted in red. The strain name, accession no., sample type, year and country of recovery, and carbapenemase genes are shown. Among 108 ST171 strains (102 from GenBank and 6 in this study), 2 strains, e1481 and e1486, had >20,000 single nucleotide polymorphisms compared with other strains (Appendix 2 Table 3), suggesting different origins. These 2 strains were therefore removed from the phylogenomic tree. NA, not available.

Precise Species Identification by Whole-Genome Sequencing of *Enterobacter* Bloodstream Infection, China

Appendix 3

The Update of Taxonomic Assignments of New Taxa Identified in this Study

Since the submission and acceptance of this manuscript, the taxonomy of *Enterobacter* has been substantially updated. The updated *Enterobacter* taxonomy is available at doi: 10.1128/mSystems.00527-20. The update of new taxa identified in this study is shown in the table below.

Appendix 3 Table. The update of taxonomic assignments of new taxons identified in genomic study of *Enterobacter* bloodstream infection

Taxonomic assignment in this study	Assignment in the updated taxonomy at doi: 10.1128/mSystems.00527-20
Taxon 1	<i>Enterobacter hoffmannii</i>
Taxon 2	<i>Enterobacter quasiroggenkampii</i>
Taxon 3	Taxon14
Taxon 4	Taxon 4
Taxon 5	Taxon 5
Taxon 6	<i>Enterobacter quasihormaechei</i>