RESEARCH LETTERS

prolonged winter break might have played a major role in reducing NPEV activity. However, considering the high contagiousness of NPEV, their activity was expected to peak after school reopening if no other interventions were implemented. The persistent low NPEV activity throughout the semester, which began in March 2020, indicated the effectiveness of other interventions.

Acknowledgments

We thank the Taiwan CDC for making their data publicly available. We also thank Kung-Yee Liang, Huey-Kang Sytwu, Shiow-Ing Wu, and Yu-Chieh Cheng for their valuable comments and support.

This project was supported by intramural grants from the National Health Research Institutes (grant nos. IV-109-PP-01, PH-109-PP-02, and PH-109-GP-02).

About the Author

Dr. Kuo is an attending physician and associate investigator at the National Institute of Infectious Diseases and Vaccinology, National Health Research Institutes, Taiwan. His primary research interest involves infectious diseases and clinical microbiology.

References

- Kuo SC, Shih SM, Chien LH, Hsiung CA. Collateral benefit of COVID-19 control measures on influenza activity, Taiwan. Emerg Infect Dis. 2020;26:1928–30. https://doi.org/10.3201/ eid2608.201192
- Galvin CJ, Li YJ, Malwade S, Syed-Abdul S. COVID-19 preventive measures showing an unintended decline in infectious diseases in Taiwan. Int J Infect Dis. 2020;98:18–20. https://doi.org/10.1016/j.ijid.2020.06.062
- Choe YJ, Lee JK. The impact of social distancing on the transmission of influenza virus, South Korea, 2020. Osong Public Health Res Perspect. 2020;11:91–2. https://doi.org/10.24171/j.phrp.2020.11.3.07
- Sakamoto H, Ishikane M, Ueda P. Seasonal influenza activity during the SARS-CoV-2 outbreak in Japan. JAMA. 2020;323:1969–71. https://doi.org/10.1001/jama.2020.6173
- Owino CO, Chu JJH. Recent advances on the role of host factors during non-poliovirus enteroviral infections. J Biomed Sci. 2019;26:47. https://doi.org/10.1186/ s12929-019-0540-y
- Jian SW, Chen CM, Lee CY, Liu DP. Real-time surveillance of infectious diseases: Taiwan's experience. Health Secur. 2017;15:144–53. https://doi.org/10.1089/hs.2016.0107

Address for correspondence: Wei J. Chen, Center for

Neuropsychiatric Research, National Health Research Institutes, No. 35 Keyan Rd, Zhunan, Miaoli 350, Taiwan; email: wjchen@nhri.edu. tw; or Shu-Chen Kuo, National Institute of Infectious Diseases and Vaccinology, National Health Research Institutes, No. 35 Keyan Rd, Zhunan, Miaoli 350, Taiwan; email: sckuo@nhri.edu.tw

Absence of SARS-CoV-2 Transmission from Children in Isolation to Guardians, South Korea

Eun Joo Lee, Dong Hyun Kim, Sung Hee Chang, Sun Bok Suh, Jina Lee, Hyunju Lee, Mi Seon Han

Author affiliations: Seongnam Citizens Medical Center, Seongnam, South Korea (E.J. Lee); Inha University Hospital, Incheon, South Korea (D.H. Kim); Seoul Medical Center, Seoul, South Korea (S.H. Chang); Busan Medical Center, Busan, South Korea (S.B. Suh); Asan Medical Center, University of Ulsan College of Medicine, Seoul (J. Lee); Seoul National University Bundang Hospital, Seoul National University College of Medicine, Seongnam (H. Lee); Seoul Metropolitan Government-Seoul National University Boramae Medical Center, Seoul (M.S. Han)

DOI: https://doi.org/10.3201/eid2701.203450

We explored transmission of severe acute respiratory syndrome coronavirus 2 among 12 children and their uninfected guardians in hospital isolation rooms in South Korea. We found that, even with close frequent contact, guardians who used appropriate personal protective equipment were not infected by children with diagnosed coronavirus disease.

Coronavirus disease (COVID-19) in children is known to occur mainly from family clusters (1). However, children can be the only infected members in a household, especially when COVID-19 is contracted from relatives or teachers. Such situations raise concerns about isolation because little information is available on transmission of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), which causes COVID-19, during colocation of young children with their uninfected guardians. Although children generally are asymptomatic or have mild symptoms, they could be infective (1,2). We explored whether SARS-CoV-2 was transmitted from children to their uninfected guardians in a hospital isolation setting.

During February 18–June 7, 2020, we analyzed all children <19 years of age with COVID-19 and their uninfected guardians who were isolated together in 7 hospitals in South Korea. The infected children were encouraged to wear face masks. The guardians were advised to wear personal protective equipment (PPE), but the degree of PPE varied among hospitals. Adherence to PPE was monitored by the medical staff; compliance was judged as good when PPE was worn most of the time, fair for frequent adherence, and poor when PPE was worn for less than half of the observed time. Children's isolation was lifted when 2 consecutive negative respiratory samples were obtained ≥24 hours apart. To ascertain secondary transmission, guardians' respiratory samples were tested for SARS-CoV-2 if symptoms developed, before their child's isolation was lifted, and 2 weeks after the end of isolation. This study was approved by the institutional review board of each hospital and written informed consent was waived.

Among 94 children with COVID-19 isolated in 7 hospitals, 12 children were isolated with a single uninfected guardian (Table). The median age of the patients was 6 years (range 2 months–11 years), and children were isolated for a median of 17 days (range 7–37 days). Most (7/12) children were asymptomatic, 4 had fever or respiratory symptoms, and 1 had pneumonia. Only 4 children cooperated well with wearing face masks.

The guardians included 10 mothers, 1 father, and 1 uncle; all complied with wearing PPE (Table). Most (10/12) guardians wore gloves and masks, either KF94 masks, which filter \approx 94% of particles of 0.4 µm in size, or N95 masks which filter \approx 95% of particles of 0.3 µm in size; 7 also wore gowns or coveralls. One guardian used a surgical mask and 1 guardian wore a KF80 mask, which filters \approx 80% of particles of 0.6 µm, and gloves. Most (10/12) guardians had frequent close contact, but 2 children kept a distance of \geq 1 m from their guardians during isolation. None of the guardians were SARS-CoV-2-positive during the study. For comparison, we also analyzed 2 cases in which adults with COVID-19 were isolated with their uninfected children because no other caregivers were available (Appendix Table, https:// wwwnc.cdc.gov/EID/article/27/1/20-3450-App1. pdf). The adult patients always wore face masks, but the 2 children never wore PPE and always had physical contact with their parents. However, the children did not become infected. The infected parents' adherence to the use of masks likely aided in curbing SARS-CoV-2 transmission to their uninfected children by reducing virus particles in respiratory droplets (3).

Appropriate use of PPE, especially face masks, might have protected uninfected guardians in our study. Previous reports have emphasized the use of face masks to prevent SARS-CoV-2 transmission in healthcare and community settings (4). Considering the decreased risk for virus transmission noted with PPE, guardians should be counseled on the proper use of PPE when in isolation with infected children.

We did not observe SARS-CoV-2 transmission from children to guardians in isolation settings in which close proximity would seem to increase transmission risk. Recent studies have suggested that children are not the main drivers of the COVID-19 pandemic, although the reasons remain unclear (5). A large study on contacts of COVID-19 case-patients in South Korea observed that household transmission was lowest when the index case-patient was 0–9 years of age (6). Among pediatric cases, the secondary

 Table. Clinical characteristics and infection control measures of 12 children with COVID-19 and their uninfected guardians sharing hospital isolation rooms, South Korea, February 18–June 7, 2020*

						Patient n	10.					
Characteristics	1	2	3	4	5	6	7	8	9	10	11	12
Patient sex	F	F	М	F	М	М	М	М	F	М	М	М
Patient age, y	6	9	8	9	11	8	0	6	6	5	4	4
Days of isolation	7	37	17	9	15	21	19	12	16	31	30	17
Symptoms	None	None	Mild FV, C	None	None	None	FV, C, S	None	None	FV, ST	FV, V	C, S
Pneumonia	Ν	Ν	Y	Ν	Ν	Ν	Ν	Ν	Ν	N	Ν	N
Face mask†	KF94	Surgical	Surgical	Surgical	Surgical	Surgical	None	KF94	KF94	KF80	None	KF94
Compliance	Good	Good	Poor	Good	Good	Poor	NA	Fair	Fair	Fair	NA	Poor
Guardian	Mother	Mother	Mother	Mother	Mother	Mother	Father	Mother	Uncle	Mother	Mother	Mother
Symptoms‡	None	None	None	None	None	None	None	None	None	None	None	C, S, H
Face mask	KF94	KF94	N95	N95	N95	N95	KF94	KF94	KF94	KF80	Surgical	KF94
Goggles	N	Y	Y	Y	Y	Y	Y	Ν	Ν	N	N	N
Gloves	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Ν	Y
Gown/coverall	Y/N	Y/N	N/Y	N/Y	N/Y	N/Y	N/Y	N/N	N/N	N/N	N/N	Y/N
Contact with patient§	<u>></u> 1 m	Close	Close	Close	Close	Close	Close	Close	<u>></u> 1 m	Close	Close	Close

*Guardians were with the children for the duration of their isolation. COVID-19, coronavirus disease; C, cough; FV, fever; H, headache; NA, not applicable; S, sputum; ST, sore throat; V, vomiting.

†Compliance was deemed good when masks were worn most of the time, fair for frequent adherence, and poor when worn <50% of observed time. A KF94 mask filters ≈94% of particles of 0.4 µm in size, an N95 mask filters ≈95% of particles of 0.3 µm in size, and a KF80 mask filters ≈80% of particles of 0.6 µm. ‡None of the guardians tested positive for COVID-19.

§Guardians for patients 1 and 9 maintained ≥1-meter distance from patients, but other guardians had frequent close contact with patients.

RESEARCH LETTERS

attack rate from children to household members was estimated to be only 0.5% (7). Reduced transmission from children in households was also reported in Switzerland and China and in educational settings in Australia (8–10).

This study is limited by its small sample size, which limits the ability to generalize its results. Moreover, we did not assess the patients' viral load, which could indirectly reflect the infectivity of the children, nor did we assess patient serology, which could further ascertain their infection status. Despite these limitations, our study provides information on SARS-CoV-2 transmission from children to guardians in isolation rooms. Additional assessments of the transmissibility of SARS-CoV-2 by children and the role of PPE in preventing infection could provide guidance during the ongoing pandemic. Nonetheless, our study adds to growing evidence that young children are less likely to contribute to the spread of COVID-19 among their adult guardians.

About the Author

Dr. E.J. Lee is a pediatrician at the Seongnam Citizens Medical Center, Seongnam, Korea. Her clinical expertise is in the field of pediatric gastroenterology, and her primary research focuses on gut microbiome in children.

References

- Han MS, Choi EH, Chang SH, Jin B-L, Lee EJ, Kim BN, et al. Clinical characteristics and viral RNA detection in children with COVID-19 in the Republic of Korea. JAMA Pediatr. 2020 Aug 28 [Epub ahead of print]. https://doi.org/10.1001/ jamapediatrics.2020.3988
- Furukawa NW, Brooks JT, Sobel J. Evidence supporting transmission of severe acute respiratory syndrome coronavirus 2 while presymptomatic or asymptomatic. Emerg Infect Dis. 2020 May 4 [Epub ahead of print]. https://doi.org/10.3201/eid2607.201595
- Leung NHL, Chu DKW, Shiu EYC, Chan KH, McDevitt JJ, Hau BJP, et al. Respiratory virus shedding in exhaled breath and efficacy of face masks. Nat Med. 2020;26:676–80. https://doi.org/10.1038/s41591-020-0843-2
- Chu DK, Akl EA, Duda S, Solo K, Yaacoub S, Schünemann HJ, et al.; COVID-19 Systematic Urgent Review Group Effort (SURGE) study authors. Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis. Lancet. 2020;395:1973–87. https://doi.org/10.1016/S0140-6736(20)31142-9
- Lee B, Raszka WV Jr. COVID-19 transmission and children: the child is not to blame. Pediatrics. 2020;146:e2020004879. https://doi.org/10.1542/peds.2020-004879
- Park YJ, Choe YJ, Park O, Park SY, Kim YM, Kim J, et al.; COVID-19 National Emergency Response Center, Epidemiology and Case Management Team. Contact tracing during coronavirus disease outbreak, South Korea, 2020. Emerg Infect Dis. 2020;26:2465–8. https://doi.org/10.3201/ eid2610.201315

- Kim J, Choe YJ, Lee J, Park YJ, Park O, Han MS, et al. Role of children in household transmission of COVID-19. Arch Dis Child. 2020 [Epub ahead of print]. https://doi.org/10.1136/ archdischild-2020-319910
- Posfay-Barbe KM, Wagner N, Gauthey M, Moussaoui D, Loevy N, Diana A, et al. COVID-19 in children and the dynamics of infection in families. Pediatrics. 2020;146:e20201576. https://doi.org/10.1542/peds.2020-1576
- Jiehao C, Jin X, Daojiong L, Zhi Y, Lei X, Zhenghai Q, et al. A case series of children with 2019 novel coronavirus infection: clinical and epidemiological features. Clin Infect Dis. 2020;71:1547–51. https://doi.org/10.1093/cid/ciaa198
- Macartney K, Quinn HE, Pillsbury AJ, Koirala A, Deng L, Winkler N, et al.; NSW COVID-19 Schools Study Team. Transmission of SARS-CoV-2 in Australian educational settings: a prospective cohort study. Lancet Child Adolesc Health. 2020;4:807–16. https://doi.org/10.1016/ S2352-4642(20)30251-0

Address for correspondence: Mi Seon Han, Department of Pediatrics, Seoul Metropolitan Government-Seoul National University Boramae Medical Center, 20 Boramae-ro 5-gil, Dongjak-gu, Seoul 07061, South Korea; email: msh0827@snu.ac.kr

Superspreading Event of SARS-CoV-2 Infection at a Bar, Ho Chi Minh City, Vietnam

Nguyen Van Vinh Chau, Nguyen Thi Thu Hong, Nghiem My Ngoc, Tran Tan Thanh, Phan Nguyen Quoc Khanh, Lam Anh Nguyet, Le Nguyen Truc Nhu, Nguyen Thi Han Ny, Dinh Nguyen Huy Man, Vu Thi Ty Hang, Nguyen Thanh Phong, Nguyen Thi Hong Que, Pham Thi Tuyen, Tran Nguyen Hoang Tu, Tran Tinh Hien, Ngo Ngoc Quang Minh, Le Manh Hung, Nguyen Thanh Truong, Lam Minh Yen, H. Rogier van Doorn, Nguyen Thanh Dung, Guy Thwaites, Nguyen Tri Dung, Le Van Tan, for the OUCRU COVID-19 research group¹

¹Members of the group are listed in the Appendix (https://wwwnc. cdc.gov/EID/article/27/1/20-3480-App1.pdf).

Absence of SARS-CoV-2 Transmission from Children in Isolation to Guardians, South Korea

Appendix

Appendix Table. Clinical characteristics and infection control measures of 2 adults with COVID-19 and their uninfected children who shared isolation rooms, South Korea, February 18–June 7, 2020*

Characteristics	Adult patient 1	Adult patient 2				
Sex	F	Μ				
Age, y	33	41				
Days of isolation	25	20				
Symptoms	Anosmia	Fever, sore throat, myalgia, cough,				
		sputum, rhinorrhea				
Pneumonia	No	Yes				
Face mask†	KF94	Surgical				
Compliance	Good	Good				
Child	Son	Son				
Age, mo.	23	17				
Contact with patient	Frequent close contact	Frequent close contact				
Symptoms‡	Mild fever	Cough, rhinorrhea				
Face mask	-	-				
Goggles	_	-				
Gloves	-	-				
Gown/coverall	-/-	-/-				

*COVID-19, coronavirus disease; -, not used.

†A KF94 mask filters $\approx\!\!94\%$ of particles of 0.4 μm in size.

‡Neither child tested positive for COVID-19.