

Acinetobacter baumannii among Patients Receiving Glucocorticoid Aerosol Therapy during Invasive Mechanical Ventilation, China

Wenchao Zhang,¹ Mei Yin,¹ Wei Li, Nana Xu, Haining Lu, Weidong Qin, Hui Han, Chen Li, Dawei Wu, Hao Wang



In support of improving patient care, this activity has been planned and implemented by Medscape, LLC and Emerging Infectious Diseases. Medscape, LLC is jointly accredited with commendation by the Accreditation Council for Continuing Medical Education (ACCME), the Accreditation Council for Pharmacy Education (ACPE), and the American Nurses Credentialing Center (ANCC), to provide continuing education for the healthcare team.

Medscape, LLC designates this Journal-based CME activity for a maximum of 1.00 **AMA PRA Category 1 Credit(s)**[™]. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

Successful completion of this CME activity, which includes participation in the evaluation component, enables the participant to earn up to 1.0 MOC points in the American Board of Internal Medicine's (ABIM) Maintenance of Certification (MOC) program. Participants will earn MOC points equivalent to the amount of CME credits claimed for the activity. It is the CME activity provider's responsibility to submit participant completion information to ACCME for the purpose of granting ABIM MOC credit.

All other clinicians completing this activity will be issued a certificate of participation. To participate in this journal CME activity: (1) review the learning objectives and author disclosures; (2) study the education content; (3) take the post-test with a 75% minimum passing score and complete the evaluation at <http://www.medscape.org/journal/eid>; and (4) view/print certificate. For CME questions, see page XXX.

Release date: November 17, 2022; Expiration date: November 17, 2023

Learning Objectives

Upon completion of this activity, participants will be able to:

- Distinguish the percentage of patients receiving invasive mechanical ventilation (IMV) who had a positive culture for *Acinetobacter baumannii* (AB)
- Assess the role of aerosol inhalation in the isolation of AB in the current study
- Analyze risk factors for the isolation of AB in the current study
- Evaluate the effects of aerosol inhalation and AB on the risk for mortality among patients receiving IMV

CME Editor

Amy J. Guinn, BA, MA, Technical Writer/Editor, Emerging Infectious Diseases. *Disclosure: Amy J. Guinn, BA, MA, has no relevant financial relationships.*

CME Author

Charles P. Vega, MD, Health Sciences Clinical Professor of Family Medicine, University of California, Irvine School of Medicine, Irvine, California. *Disclosure: Charles P. Vega, MD, has the following relevant financial relationships: served as an advisor or consultant for GlaxoSmithKline; Johnson & Johnson Pharmaceutical Research & Development, L.L.C.*

Authors

Wenchao Zhang, MD, PhD; Mei Yin, MD, PhD; Wei Li, MD, PhD; Nana Xu, MD; Haining Lu, MD; Weidong Qin, MD, PhD; Hui Han, MD; Chen Li, MD; Dawei Wu, MD; Hao Wang, MD, PhD.

Author affiliations: Qilu Hospital of Shandong University, Jinan, China (W. Zhang, M. Yin, W. Li, N. Xu, W. Qin, H. Han, C. Li, H. Wang); Qingdao Branch, Qilu Hospital of Shandong University, Qingdao, China (H. Lu, D. Wu)

DOI: <https://doi.org/10.3201/eid2812.220347>

¹These authors contributed equally to this article.

Acinetobacter baumannii is a nosocomial pathogen associated with severe illness and death. Glucocorticoid aerosol is a common inhalation therapy in patients receiving invasive mechanical ventilation. We conducted a prospective cohort study to analyze the association between glucocorticoid aerosol therapy and *A. baumannii* isolation from ventilator patients in China. Of 497 enrolled patients, 262 (52.7%) received glucocorticoid aerosol, and *A. baumannii* was isolated from 159 (32.0%). Glucocorticoid aerosol therapy was an independent risk factor for *A. baumannii* isolation (hazard ratio 1.5, 95% CI 1.02–2.28; $p = 0.038$). Patients receiving glucocorticoid aerosol had a higher cumulative hazard for *A. baumannii* isolation and analysis showed that glucocorticoid aerosol therapy increased *A. baumannii* isolation in most subpopulations. Glucocorticoid aerosol was not a direct risk factor for 30-day mortality, but *A. baumannii* isolation was independently associated with 30-day mortality in ventilator patients. Physicians should consider potential *A. baumannii* infection when prescribing glucocorticoid aerosol therapy.

Acinetobacter baumannii, a gram-negative coccobacillus, is a major nosocomial pathogen worldwide. *A. baumannii* is particularly challenging in intensive care units (ICUs). According to the Extended Prevalence of Infection in Intensive Care study, aimed at providing information on the prevalence of infection in ICUs worldwide, *Acinetobacter* spp. constituted 8.8% of all culture-positive ICU infections in 2007 (1), which increased to 11.4% in 2017 (2). However, infection rates differed markedly, ranging from 1.0% in North America to 25.6% in Asia and the Middle East and 22.9% in eastern Europe (2). Patients on invasive mechanical ventilation are particularly vulnerable to *A. baumannii* infection and colonization due to airway barrier destruction and bacterial virulence factors such as motility, epithelial adherence, and biofilm formation that enable *A. baumannii* colonization in the airways (3,4). *A. baumannii* in patient airways is associated with longer hospitalization, higher medical expenses, and increased mortality rates (5–7). Identifying risk factors for *A. baumannii* infection is crucial for implementing preventive measures and decreasing overall illness and death.

Aerosol inhalation is widely used in patients requiring mechanical ventilation. Glucocorticoids are frequently administered during aerosol therapy, especially in China (8–10). Compared with systemic application, aerosol therapy has several advantages, including targeted delivery to the lungs, faster response, and fewer systemic side effects (11,12). However, the aerosols and droplets generated during aerosol inhalation can become sources of respiratory pathogens (13), and inhaled glucocorticoids might

suppress pulmonary immunity (14), which could increase the opportunity for nosocomial acquisition. Inhaled corticosteroids are associated with an increased risk for pneumonia in patients with chronic obstructive pulmonary disease (COPD) (15). However, the effects of glucocorticoid aerosol inhalation on nosocomial infection risk has not been clearly elucidated.

Glucocorticoid aerosol therapy is mainly indicated for patients with asthma, COPD (16), acute respiratory distress syndrome (ARDS) (17), and some pathophysiological conditions, such as airway hyperresponsiveness (18), hyperinflammation, and mucosal edema (19). In the past decade, use of glucocorticoid aerosol therapy has increased in hospitals in China; on average, >40% of patients on mechanical ventilation receive this therapy (9). In addition, a market analysis determined that aerosolized glucocorticoid sales in China were almost 3-fold higher in 2018 than in 2012 (20).

Although epidemiology has demonstrated a slow increase in *A. baumannii* infection globally over the past decade (1,2), the increase in *A. baumannii* incidence in China appears to have outpaced increases in other regions worldwide (21–23). According to the China Antimicrobial Surveillance Network (CHINET), a national surveillance of the trends of bacterial strains isolated from the major hospitals in China, the number of *Acinetobacter* spp. strains increased by 2.7-fold in 2018 compared with 2012 (23,24). Previously, we reported a marked increase in the incidence of *A. baumannii*-related bloodstream infections and incidence of pneumonia-related *A. baumannii* infections in ICUs in China that were 3.2-fold higher during 2017–2018 than during 2011–2012 (25). *A. baumannii* was the most frequent bacterial isolate in ventilator-associated pneumonia in China, and rates were 35.7%–52.7% (26,27). Furthermore, the incidence of the drug-resistant phenotype of *A. baumannii* is high. According to CHINET reports, carbapenem-resistant *A. baumannii* strains increased from 31% in 2005 to 66.7% in 2014 (28), then to ≈80% in 2018 (29). We previously reported that carbapenem resistance rates in ICUs in China increased from 25% during 2011–2012 to 95.7% during 2017–2018 (25). A multicenter study of ICUs in China reported that multidrug-resistant (MDR) *A. baumannii* was detected in 40% of all cases (30).

We hypothesized that increased use of glucocorticoid aerosol therapy might contribute to increased *A. baumannii* incidence. Therefore, we performed a prospective cohort study of critically ill patients receiving invasive mechanical ventilation in China to determine whether use of aerosolized glucocorticoid increased the risk for *A. baumannii* isolation.

Methods

Study Design and Patients

During January 2018–August 2019, we conducted a prospective cohort study at 3 adult ICUs in 2 hospitals in Shandong Province, China: Qilu Hospital of Shandong University in Jinan and Qingdao Branch of Qilu Hospital in Qingdao. We enrolled patients on their first day of invasive mechanical ventilation in the ICU and obtained written informed consent for all patients. We divided the patients into 3 groups on the basis of their treatment: no aerosol inhalation therapy, glucocorticoid aerosol therapy, and aerosol inhalation without glucocorticoid. Within 48 hours of patient enrollment, we collected secretion samples from the lower respiratory tract by transtracheal aspiration for microbial culture; thereafter, we collected samples 3 times per week until we obtained an *A. baumannii*-positive culture. We followed patients for 30 days after enrollment. If the patient was hospitalized for >3 weeks, we reduced the culture frequency to once a week. We excluded patients who received invasive mechanical ventilation for <48 hours; received aerosol inhalation or glucocorticoid aerosol for <48 hours after enrollment and before *A. baumannii*-positive culture; were <18 years of age; were assumed to have *A. baumannii* infection or colonization at baseline because they were *A. baumannii*-positive before enrollment or within the first 48 hours of enrollment; or had been exposed to ≥ 1 of the following *A. baumannii* risk factors before enrollment: antimicrobial drugs for ≥ 7 days, invasive mechanical ventilation for ≥ 5 days, or vasopressor for ≥ 3 days. We also excluded patients who lacked follow-up data or had incomplete information. The Institutional Ethics Committee of Qilu Hospital of Shandong University approved our study.

Microbiology

We performed microbial cultures according to standard procedures. In brief, we incubated respiratory samples on MacConkey agar plates at 5% CO₂ and 35°C for 48 h. We identified *A. baumannii*, a gram-negative, nonfermentative, and oxidase-negative coccobacillus, by using the VITEK 2 compact system and GN ID card (bioMérieux, <https://www.biomerieux.com>). We used *Escherichia coli* (ATCC accession no. 25922) and *Pseudomonas aeruginosa* (ATCC accession no. 27853) as quality controls.

Definitions and Data Collection

We defined no aerosol inhalation as patients who did not receive aerosolized medications during the

study. We defined glucocorticoid aerosol therapy as patients who received aerosolized glucocorticoids for ≥ 48 hours after enrollment and before *A. baumannii* isolation, with or without nonglucocorticoid aerosolized medications for any duration. We defined aerosol inhalation without glucocorticoid as patients who received only aerosolized nonglucocorticoid medications, such as bronchodilators and expectorants, for ≥ 48 hours after enrollment and before *A. baumannii* isolation. We excluded all other conditions.

The primary endpoint was *A. baumannii* isolation, which we defined as *A. baumannii*-positive culture from the lower respiratory tract samples collected during the ICU stay. Negative outcomes were no *A. baumannii* isolation before death, ICU discharge, or end of follow-up period. We recorded the time-to-event, which we defined as number of days from enrollment to *A. baumannii* isolation.

We collected baseline information at ICU admission, including age, sex, history of smoking and surgeries, underlying conditions, past inhaled steroids for chronic conditions, and Charlson comorbidity index. We used the Acute Physiology and Chronic Health Evaluation II (APACHE II) score to assess illness severity. We also recorded other possible *A. baumannii* risk factors, such as use of broad-spectrum antimicrobial drugs, invasive mechanical ventilation, urethral catheter placement, vasopressor treatment, renal dialysis, and length of ICU stay. In addition, we recorded indications for glucocorticoid aerosol therapy by reviewing patients' medical records. We reviewed patients' clinical data to determine *A. baumannii* isolation status as infection, colonization, or undefined.

Statistical Analyses

We expressed continuous variables as median and interquartile range (IQR) or mean and SD and categorical variables as number and percentage. We used univariate and multivariate Cox proportional hazards regression and hazard ratio (HR) and 95% CI to assess risk factors for *A. baumannii* isolation and 30-day mortality. We performed propensity score matching analysis to reduce the imbalance between the glucocorticoid aerosol therapy and nonglucocorticoid groups. We included all possible covariables (i.e., demographics, background history, underlying conditions, and disease severity) in the propensity score matching. We calculated propensity scores by using a logistic regression model. We applied a 1:1 nearest neighbor matching algorithm with a caliper of 0.02 and without replacement. We assessed balance of variables in both groups by standardized differences.

We analyzed *A. baumannii* isolation in complete cases and the propensity-matched cohort. We used Kaplan-Meier curves to visually compare cumulative hazards for *A. baumannii* isolation among the 3 groups, which we evaluated by using a log-rank test. To assess the consistency of glucocorticoid therapy in terms of its effect on *A. baumannii* isolation from prespecified subgroups with different characteristics, we applied Cox proportional hazards model with Efron's method for handling ties and used forest plots for HRs and 95% CIs. We assessed heterogeneity of efficacy of glucocorticoid therapy on *A. baumannii* isolation in subgroups by using an interaction test, expressed p values for interaction, and considered $p < 0.05$ statistically significant. We performed all analyses by using SPSS Statistics 16.0 (IBM, <https://www.ibm.com>) and R version 3.0 (R Foundation for Statistical Computing, <https://www.r-project.org>).

Results

Participant Characteristics

We enrolled 671 patients from 3 ICUs and excluded 174 patients. The final cohort consisted of 497 patients: 137 (27.6%) received no aerosol inhalation, 262 (52.7%) received glucocorticoid aerosol, and 98 (19.7%) received aerosol inhalation without glucocorticoid (Figure 1). We isolated *A. baumannii* from 159 (32.0%) patients. The median patient age was 60.1 (IQR 49–73) years, and 67.8% were male. The median length

of ICU stay was 15 (IQR 7–23) days. Besides *A. baumannii*, the 3 other bacteria commonly isolated were *Klebsiella pneumoniae* ($n = 38$, 7.6%), *P. aeruginosa* ($n = 27$, 5.4%), and *E. coli* ($n = 15$, 3.0%) (Appendix Table 1, <https://wwwnc.cdc.gov/EID/article/28/12/22-0347-App1.pdf>). Most (22.5%) study patients received glucocorticoid therapy for ARDS and for asthma or COPD (21.3%) (Appendix Table 2).

Risk Factors for A. baumannii Isolation

We performed univariate Cox regression analysis of risk factors for *A. baumannii* isolation (Table 1). Compared with no aerosol inhalation, glucocorticoid aerosol therapy had a statistically significant effect on *A. baumannii* isolation (HR 1.860, 95% CI 1.264–2.738; $p = 0.002$). Aerosol inhalation without glucocorticoid was not a risk factor for *A. baumannii* ($p > 0.05$). Other candidate risk factors were cardiovascular diseases, chronic renal insufficiency, COPD or asthma, current or former smoking history, use of broad-spectrum antimicrobial drugs for ≥ 7 days, invasive mechanical ventilation for ≥ 5 days, vasopressor treatment, renal dialysis for ≥ 3 days, and APACHE II score.

To assess whether glucocorticoid aerosol therapy was an independent risk factor for *A. baumannii* isolation, we established 2 models using multivariate Cox regression analysis in complete cases. Model 1 included all variables, and model 2 only included variables with $p < 0.1$ in the univariate analysis. Glucocorticoid aerosol was an independent risk factor for *A. baumannii*




Figure 1. Flowchart for enrolling participants in a study of *Acinetobacter baumannii* among patients receiving glucocorticoid aerosol therapy during invasive mechanical ventilation, China. ICU, intensive care units.

Table 1. Univariate analysis of risk factors for *Acinetobacter baumannii* among patients during invasive mechanical ventilation, China*

Variables	Overall, n =	<i>Acinetobacter baumannii</i> isolated		p value	Hazard ratio (95% CI)
	497	Yes, n = 159	No, n = 338		
Median age, y (IQR)	60.1 (49–73)	61 (50–74)	59.6 (49–73)	0.444	1.004 (0.994–1.013)
Sex, no. (%)					
F	160 (32.2)	44 (27.7)	116 (34.3)	Referent	
M	337 (67.8)	115 (72.3)	222 (65.7)	0.139	0.769 (0.544–1.089)
Mean Charlson comorbidity index, (SD)	4.26 (2.12)	4.40 (2.14)	4.20 (2.11)	0.293	1.038 (0.968–1.113)
Underlying conditions, no. (%)					
Cardiovascular diseases	200 (40.2)	79 (49.7)	121 (35.8)	0.003	1.596 (1.169–2.178)
Chronic renal insufficiency	179 (36.0)	71 (44.7)	108 (32.0)	0.011	1.504 (1.200–2.056)
COPD and asthma	176 (35.4)	71 (44.7)	105 (31.1)	0.005	1.570 (1.148–2.146)
Type 2 diabetes mellitus	116 (23.3)	46 (28.9)	70 (20.7)	0.057	1.395 (0.990–1.965)
Solid tumor	100 (20.1)	36 (22.6)	64 (18.9)	0.363	1.188 (0.820–1.723)
Hematologic malignancy	31 (6.2)	8 (5.0)	23 (6.8)	0.497	0.781 (0.384–1.591)
Past inhaled steroids use for chronic conditions	47 (9.5)	17 (10.7)	30 (8.9)	0.450	1.214 (0.734–2.007)
Current or former smoker	187 (37.6)	74 (46.5)	113 (33.4)	0.005	1.565 (1.146–2.138)
Postoperative admission	142 (28.6)	38 (23.9)	104 (30.8)	0.134	0.757 (0.526–1.090)
Treatment, no. (%)					
No aerosol inhalation	137 (27.6)	33 (20.8)	104 (30.8)	Referent	
Glucocorticoid aerosol inhalation	262 (52.7)	107 (67.3)	155 (45.9)	0.002	1.860 (1.264–2.738)
Aerosol inhalation without glucocorticoid	98 (19.7)	19 (11.9)	79 (23.4)	0.337	0.760 (0.433–1.332)
Broad-spectrum antimicrobial drugs, ≥ 7 d	417 (83.9)	157 (98.7)	260 (76.9)	<0.001	9.539 (4.595–18.795)
Invasive mechanical ventilation, ≥ 5 d	221 (44.5)	112 (70.4)	109 (32.2)	<0.001	3.452 (2.453–4.858)
Urethral catheter placement, ≥ 3 d	493 (99.2)	158 (99.4)	335 (99.1)	0.875	1.171 (0.164–8.361)
Vasopressor treatment, ≥ 3 d	75 (15.1)	42 (26.4)	33 (9.8)	<0.001	2.634 (1.850–3.750)
Renal dialysis, ≥ 3 d	84 (16.9)	34 (21.4)	50 (14.8)	0.063	1.432 (0.980–2.093)
APACHE II score, mean (SD)	18.18 (6.03)	18.98 (6.44)	17.80 (5.80)	0.053	1.026 (1.000–1.053)
Median length of ICU stay, d (IQR)	15 (7–23)	20 (10–28)	13 (6–20)	0.057	1.005 (1.000–1.010)

*APACHE II, Acute Physiology and Chronic Health Evaluation II; COPD, chronic obstructive pulmonary disease; ICU, intensive care unit; IQR, interquartile range.

isolation in both model 1 (HR 1.499, 95% CI 1.001–2.246; $p = 0.049$) (Appendix Table 3) and model 2 (HR 1.528, 95% CI 1.024–2.278; $p = 0.038$) (Table 2). Cardiovascular diseases, prolonged use of broad-spectrum antimicrobial drugs, invasive mechanical ventilation, and vasopressor treatment were other independent risk factors for *A. baumannii* (Table 2; Appendix Table 3). As a whole variable, aerosol inhalation had no effect on *A. baumannii* isolation (Appendix Table 4).

In the propensity-matched cohort, the possible glucocorticoid-related covariables were balanced in both

groups (Appendix Table 5). Univariate and multivariate Cox regression analyses also indicated that glucocorticoid aerosol therapy was an independent risk factor for *A. baumannii* isolation (Table 3; Appendix Table 6). In an independent model that included indications for glucocorticoid aerosol therapy, risk factors for *A. baumannii* isolation were glucocorticoid aerosol treatments for COPD or asthma and for ARDS (Appendix Table 7).

Log-rank analysis showed that the difference among the groups was statistically significant ($p < 0.001$). The cumulative hazard for *A. baumannii*

Table 2. Multivariate analysis of risk factors for *Acinetobacter baumannii* among patients during invasive mechanical ventilation, China*

Variables	p value	Hazard ratio (95% CI)
Underlying conditions		
Cardiovascular diseases	0.054	1.394 (0.994–1.955)
Chronic renal insufficiency	0.730	0.937 (0.648–1.356)
COPD and asthma	0.132	1.299 (0.924–1.825)
Type 2 diabetes mellitus	0.325	1.197 (0.837–1.714)
Current or former smoker	0.098	1.307 (0.951–1.797)
Treatment		
No aerosol inhalation	Referent	
Glucocorticoid aerosol inhalation	0.038	1.528 (1.024–2.278)
Aerosol inhalation without glucocorticoid	0.524	0.829 (0.467–1.475)
Broad-spectrum antimicrobial drugs, ≥ 7 d	0.001	7.238 (2.758–15.788)
Invasive mechanical ventilation, ≥ 5 d	0.001	2.381 (1.664–3.405)
Vasopressor treatment, ≥ 3 d	<0.001	2.060 (1.402–3.028)
Renal dialysis, ≥ 3 d	0.841	1.046 (0.675–1.620)
APACHE II score	0.586	0.992 (0.965–1.020)

*Results are from model 2; only variables with $p < 0.1$ in univariate analysis were included. APACHE II, Acute Physiology and Chronic Health Evaluation II; COPD, chronic obstructive pulmonary disease.

Table 3. Multivariate analysis of risk factors for *Acinetobacter baumannii* among propensity-matched patient cohort during invasive mechanical ventilation, China*

Variables	p value	Hazard ratio (95% CI)
Underlying conditions		
Cardiovascular diseases	0.117	1.361 (0.926–2.001)
Chronic renal insufficiency	0.800	1.052 (0.712–1.554)
Type 2 diabetes mellitus	0.243	1.271 (0.850–1.899)
Current or former smoker	0.051	1.442 (0.998–2.083)
Treatment		
Glucocorticoid aerosol inhalation	0.032	1.489 (1.036–2.141)
Broad-spectrum antimicrobial drugs, ≥ 7 d	0.004	6.315 (2.543–13.921)
Invasive mechanical ventilation, ≥ 5 d	<0.001	2.388 (1.614–3.534)
Vasopressor treatment, ≥ 3 d	0.501	1.188 (0.719–1.963)
APACHE II score	0.363	1.014 (0.984–1.045)

*Only variables with $p < 0.1$ in univariate analysis of the propensity-matched cohort were included. APACHE II, Acute Physiology and Chronic Health Evaluation II; COPD, chronic obstructive pulmonary disease.

isolation was significantly higher in the glucocorticoid aerosol group compared with the no aerosol inhalation (HR 1.871; 95% CI 1.206–2.772; $p < 0.001$) and aerosol inhalation without glucocorticoid (HR 2.316; 95% CI 1.482–3.620; $p = 0.002$) groups (Figure 2).

Effect of Glucocorticoid Aerosol Therapy on *A. baumannii* Isolation among Subgroups

We divided patients into subgroups to evaluate the contribution of glucocorticoid aerosol to *A. baumannii* isolation from different subpopulations. We found glucocorticoid aerosol was a promoting factor for *A. baumannii* isolation from most subpopulations, except patients with type 2 diabetes mellitus, hematologic malignancy, antimicrobial drug use for *A. baumannii*, and short ICU stays ($p > 0.05$) (Figure 3). We noted no statistically significant interactions between most prespecified subgroups defined by demographics, medical history, underlying conditions, APACHE II score, treatment measures, and length of ICU stay (interaction $p > 0.05$). The favorable effect of glucocorticoid aerosol on *A. baumannii* isolation was relatively greater in the subgroup of patients with longer vasopressor treatment (interaction $p = 0.006$) (Figure 3).

Association between Glucocorticoid Aerosol Therapy and Clinical Prognosis

We performed univariate and multivariate Cox regression analyses to evaluate the possible risk factors for 30-day mortality in critically ill patients on ventilators. We found glucocorticoid aerosol was not a risk factor for 30-day mortality in those patients, but *A. baumannii* isolation was independently associated with 30-day mortality (HR 1.824, 95% CI 1.317–2.104; $p = 0.045$) (Appendix Tables 8, 9). A further separate analysis of *A. baumannii* isolation status showed that *A. baumannii* infection was independently associated with 30-day mortality (HR 2.759, 95% CI 1.575–4.833; $p = 0.012$) (Appendix Tables 8, 10).

Discussion

In this study, we assessed the effect of the commonly used glucocorticoid aerosol therapy on the frequency of *A. baumannii*-positive cultures from lower respiratory tract samples in 3 ICUs in China. After controlling for other variables, our findings showed that glucocorticoid aerosol increased the risk for *A. baumannii* isolation from critically ill patients on invasive mechanical ventilation.

A. baumannii is ubiquitous in nature and is becoming more frequent in hospitals. In our study, 32% of patients acquired *A. baumannii* during the 30-day follow-up period. Over the past 2 decades, several




Figure 2. Kaplan-Meier curves of cumulative hazards of different aerosol inhalation treatments on *Acinetobacter baumannii* isolation among patients during invasive mechanical ventilation, China. We used log-rank analysis to compare hazard ratios over time among all groups ($p < 0.001$), glucocorticoid aerosol therapy group with the no aerosol inhalation group ($p < 0.001$), and the glucocorticoid aerosol therapy group with the aerosol inhalation without glucocorticoid group ($p = 0.002$).




Figure 3. Effects of glucocorticoid aerosol therapy on *Acinetobacter baumannii* isolation during invasive mechanical ventilation among prespecified patient subgroups, China. We applied Cox proportional hazards model with Efron's method for handling ties between groups to assess favorable effects of glucocorticoid aerosol on *A. baumannii* isolation. Black horizontal marks indicate hazard ratios, and error bars indicate 95% CIs. APACHE II, Acute Physiology and Chronic Health Evaluation II; COPD, chronic obstructive pulmonary disease; ICU, intensive care unit. *Limited numbers for analysis. †For interaction, p = 0.006.

studies have attempted to characterize and identify risk factors for *A. baumannii* colonization or infection. Invasive operations, such as endotracheal mechanical ventilation, inserted invasive devices, ICU stays, recent surgery, use of broad-spectrum antimicrobial drugs, ineffective antimicrobial therapy, and septic shock at diagnosis, are risk factors for MDR *A. baumannii* colonization or infection and for death (5,31–33).

We determined that prolonged use of broad-spectrum antimicrobial drugs, invasive mechanical ventilation, and vasopressor treatment were independent risk factors for *A. baumannii* isolation from ventilated patients, which is consistent with previous studies (5,31–33). A previous study reported that cardiovascular organ failure was an independent risk factor associated with *A. baumannii* bloodstream infection (34). Of note, model 1 of our study showed that cardiovascular disease also was an independent risk factor for *A. baumannii* isolation. Another population-based study reported that patients with chronic heart failure had a markedly increased risk for hospitalization with pneumonia (35), indicating a possible correlation between cardiovascular diseases and pneumonia. However, the specific underlying mechanisms by which cardiovascular disease promotes *A. baumannii* isolation remain unknown.

In critically ill patients undergoing mechanical ventilation, aerosol inhalation is a common intervention for treating various pulmonary diseases. An international survey demonstrated that 99% of 611 ICUs from 70 countries reported using aerosol therapy during mechanical ventilation, including noninvasive ventilation, and the most frequently delivered drugs were bronchodilators and steroids (36). A web-based survey involving 447 hospitals in mainland China recorded a high proportion of aerosol therapy in both invasive (90.8%) and noninvasive (91.3%) mechanical ventilation; bronchodilators (64.8%) and topical corticosteroids (43.4%) were the most commonly used drugs (9). Aerosol inhalation is aimed at reversing bronchoconstriction, decreasing the work of breathing, relieving dyspnea, modifying the inflammatory response (19,37), ameliorating lung injury (38), and reducing the rate of exacerbation in both asthma and COPD. However, in patients with COPD and asthma, inhaled corticosteroids are associated with an increased risk for upper respiratory tract infections (39,40), pneumonia, and lower respiratory tract infections (15,41).

Ventilated patients are already vulnerable to pneumonia. Therefore, evaluating whether commonly used aerosol therapy increases the risk for

nosocomial pneumonia is crucial, especially when inhaled with glucocorticoids. Because *A. baumannii*-related pneumonia is associated with severe illness and death, we chose *A. baumannii* isolation as an outcome and explored its relationship with glucocorticoid aerosol therapy. Our study showed that glucocorticoid aerosol therapy was an independent risk factor for *A. baumannii* isolation from patients on ventilators. Compared with no aerosol inhalation, glucocorticoid aerosol inhalation increased the risk for *A. baumannii* by ≈ 1.5 times. Although further analysis revealed that glucocorticoid aerosol was not directly associated with 30-day mortality, it still might contribute to poor clinical prognosis due to its effect on *A. baumannii* isolation. As we described, *A. baumannii*, especially MDR *A. baumannii* pneumonia, is well recognized as a risk factor for death. In this study, we also found *A. baumannii* isolation was an independent risk factor for 30-day mortality in patients receiving invasive mechanical ventilation. Because glucocorticoid aerosol heightened the likelihood of acquiring *A. baumannii*, it might exert a secondary effect, death among *A. baumannii*-infected patients. Thus, further investigation in a much larger patient population could describe a downstream mortality effect of glucocorticoid aerosol therapy.

When we included glucocorticoid aerosol indications in multivariate analysis, we found COPD and asthma and possible ARDS were independent risk factors for *A. baumannii* isolation. Because these structural or underlying lung diseases and severe acute lung injury necessitate longer duration of mechanical ventilation, our results were compatible with previously described risk factors for *A. baumannii* infection. In contrast to glucocorticoid aerosol, we did not detect an association between aerosol inhalation without glucocorticoid and *A. baumannii*. Because both therapies generate aerosols, our previous concern that aerosols were a source of *A. baumannii* acquisition might not be reasonable.

Reasons why glucocorticoids increase the risk for *A. baumannii* isolation remain elusive. Previously considered sterile, healthy lungs harbor complex and dynamic microbiota communities (42). Pulmonary diseases, such as COPD (43), asthma (44), lung cancer (45), and ARDS (46), cause considerable alteration of lung microbiota. Pneumonia pathogenesis involves an abrupt and emergent disruption in the complex homeostasis of the lung microbial ecosystem (47). A recent study reported that inhaled corticosteroids altered the lung microbiota in both COPD patients and mouse models and impaired bacterial control in models with *Streptococcus pneumoniae* infection (14).

Antimicrobial peptides, also known as host defense peptides, are short and generally positively charged peptides that participate in the regulation of the host's antibacterial actions and immune defense (48). Another study observed that cathelicidin, an antimicrobial peptide, was impaired by inhaled corticosteroids among COPD patients by increasing protease cathepsin D, thereby promoting the proliferation of *Streptococcus* (14). In a bacterial 16S rRNA gene sequencing and host transcriptomic analysis, another study reported that as COPD severity increased, the airway microbiome becomes associated with decreased abundance of *Prevotella* bacteria in concert with downregulation of genes promoting epithelial defense associated with inhaled corticosteroid use (49). Evidence also suggests that in asthma, inhaled corticosteroids can alter the relative abundance of genera in airway microbiome (50). Therefore, inhaled corticosteroids play a primary role in lung microbiota disruption and host-defense suppression, which could explain why glucocorticoid aerosol contributed to the increased risk for *A. baumannii* isolation from patients on ventilators.

Clinicians should individualize patient care and manage treatments on the basis of subgroup analysis results. Our results showed that, in most subpopulations, regardless of the presence or absence of the pre-specified characteristics, patients were at higher risk for *A. baumannii* isolation when receiving glucocorticoid aerosol. Based on our findings, we recommend that intensive care teams more carefully consider the risk of widespread usage of glucocorticoid aerosol in patients with invasive mechanical ventilation. Because glucocorticoid aerosol therapy had a much greater favorable effect on *A. baumannii* isolation in the subgroup of patients on vasopressors for ≥ 3 days, clinicians should be particularly cautious about giving glucocorticoid aerosol to patients on long vasopressor treatments. Because patients with diabetes, hematologic malignancy, and shorter ICU stays were at relatively lower risk for glucocorticoid aerosol-associated *A. baumannii* acquisition, glucocorticoid aerosol could be considered when appropriate for indications in these patients. Altogether, our study suggests ICU teams need to identify the specific patient subgroups that will truly benefit from glucocorticoid aerosol therapy rather than more generalized administration. Limiting glucocorticoid aerosol use might be considered as part of existing antimicrobial stewardship bundles. In addition, defining the duration of glucocorticoid aerosol therapy might help maximize benefit while reducing associated risk. Interventional studies exploring the effects of different glucocorticoid

aerosol therapy durations on the occurrence of various types of nosocomial pneumonia are needed.

The first limitation of our study is that, because it was an observational study, confounders that might influence the effect of glucocorticoid aerosol on *A. baumannii* isolation remain, even after adjusting by subgroup analysis, propensity score matching analysis, and multivariable Cox regression. A similar situation exists for the analysis of risk factors for death. Second, a potential time bias remains, in which patients might have been exposed to risk factors before study enrollment. However, we have excluded patients with prolonged exposure to several well-known risk factors to minimize the possible effect of time bias. Third, exclusion of certain cases might pose a potential selection bias, including survivorship bias. Future randomized controlled interventional studies are expected to confirm our findings and minimize the effects of confounders and the above biases. Fourth, the study included only 3 ICUs in China and did not focus on pathogens other than *A. baumannii*. Research involving more centers and more cases could explore whether glucocorticoid aerosol is a common risk factor in the overall nosocomial pneumonia risk.

In conclusion, we found glucocorticoid aerosol therapy was an independent risk factor for *A. baumannii* isolation from patients receiving invasive mechanical ventilation. Because of high mortality rates associated with *A. baumannii*-related nosocomial pneumonia, clinicians should carefully consider both the beneficial and harmful effects of glucocorticoid aerosol before administering this therapy.

Acknowledgments

We thank Oxford Science Editing Ltd. for language editing.

This work was supported by the National Natural Science Foundation of China (grant nos. 81873927, 82072231), Taishan Scholars Program of Shandong Province (award no. tsqn202103165), Clinical Research Center of Shandong University (grant no.2020SDUCRCC013), Natural Science Foundation of Shandong Province (grant no. ZR2020QH021), China Postdoctoral Science Foundation (grant no. 2018M632685), and the Project of Science and Technology of Qingdao People's Livelihood (grant no. 19-6-1-23-nsh).

About the Author

Dr. Zhang is an attending physician and a research scientist at the Department of Critical Care Medicine, Qilu Hospital of Shandong University, Jinan, China. Her research interest is infection and multiorgan failure related to critical illnesses.

References

- Vincent JL, Rello J, Marshall J, Silva E, Anzueto A, Martin CD, et al.; EPIC II Group of Investigators. International study of the prevalence and outcomes of infection in intensive care units. *JAMA*. 2009;302:2323–9. <https://doi.org/10.1001/jama.2009.1754>
- Vincent JL, Sakr Y, Singer M, Martin-Loeches I, Machado FR, Marshall JC, et al.; EPIC III Investigators. Prevalence and outcomes of infection among patients in intensive care units in 2017. *JAMA*. 2020;323:1478–87. <https://doi.org/10.1001/jama.2020.2717>
- Mea HJ, Yong PVC, Wong EH. An overview of *Acinetobacter baumannii* pathogenesis: motility, adherence and biofilm formation. *Microbiol Res*. 2021;247:126722. <https://doi.org/10.1016/j.micres.2021.126722>
- Lynch JP III, Zhanel GG, Clark NM. Infections due to *Acinetobacter baumannii* in the ICU: treatment options. *Semin Respir Crit Care Med*. 2017;38:311–25. <https://doi.org/10.1055/s-0037-1599225>
- Shi J, Sun T, Cui Y, Wang C, Wang F, Zhou Y, et al. Multidrug resistant and extensively drug resistant *Acinetobacter baumannii* hospital infection associated with high mortality: a retrospective study in the pediatric intensive care unit. *BMC Infect Dis*. 2020;20:597. <https://doi.org/10.1186/s12879-020-05321-y>
- Sunshine RH, Wright MO, Maragakis LL, Harris AD, Song X, Hebden J, et al. Multidrug-resistant *Acinetobacter* infection mortality rate and length of hospitalization. *Emerg Infect Dis*. 2007;13:97–103. <https://doi.org/10.3201/eid1301.060716>
- Blanco N, Harris AD, Rock C, Johnson JK, Pineles L, Bonomo RA, et al.; CDC Epicenters Program. Risk factors and outcomes associated with multidrug-resistant *Acinetobacter baumannii* upon intensive care unit admission. *Antimicrob Agents Chemother*. 2017;62:e01631-17. <https://doi.org/10.1128/AAC.01631-17>
- Zhang C, Mi J, Wang X, Lv S, Zhang Z, Nie Z, et al. Knowledge and current practices of ICU nurses regarding aerosol therapy for patients treated with invasive mechanical ventilation: a nationwide cross-sectional study. *J Clin Nurs*. 2021;30:3429–38. <https://doi.org/10.1111/jocn.15639>
- Zhang Z, Xu P, Fang Q, Ma P, Lin H, Fink JB, et al.; China Union of Respiratory Care (CURC). Practice pattern of aerosol therapy among patients undergoing mechanical ventilation in mainland China: a web-based survey involving 447 hospitals. *PLoS One*. 2019;14:e0221577. <https://doi.org/10.1371/journal.pone.0221577>
- Lyu S, Li J, Wu M, He D, Fu T, Ni F, et al.; Respiratory Care Committee in Chinese Thoracic Society. The use of aerosolized medications in adult intensive care unit patients: A prospective, multicenter, observational, cohort Study. *J Aerosol Med Pulm Drug Deliv*. 2021;34:383–91. <https://doi.org/10.1089/jamp.2021.0004>
- Lyu S, Li J, Yang L, Du X, Liu X, Chuan L, et al. The utilization of aerosol therapy in mechanical ventilation patients: a prospective multicenter observational cohort study and a review of the current evidence. *Ann Transl Med*. 2020;8:1071. <https://doi.org/10.21037/atm-20-1313>
- Dhand R. Inhaled drug therapy 2016: the year in review. *Respir Care*. 2017;62:978–96. <https://doi.org/10.4187/respcare.05624>
- Ari A. Practical strategies for a safe and effective delivery of aerosolized medications to patients with COVID-19. *Respir Med*. 2020;167:105987. <https://doi.org/10.1016/j.rmed.2020.105987>
- Singanayagam A, Glanville N, Cuthbertson L, Bartlett NW, Finney LJ, Turek E, et al. Inhaled corticosteroid suppression of cathelicidin drives dysbiosis and bacterial infection in chronic obstructive pulmonary disease. *Sci Transl Med*. 2019;11:eaav3879. <https://doi.org/10.1126/scitranslmed.aav3879>
- Ernst P, Gonzalez AV, Brassard P, Suissa S. Inhaled corticosteroid use in chronic obstructive pulmonary disease and the risk of hospitalization for pneumonia. *Am J Respir Crit Care Med*. 2007;176:162–6. <https://doi.org/10.1164/rccm.200611-1630OC>
- Sibila O, Soto-Gomez N, Restrepo MI. The risk and outcomes of pneumonia in patients on inhaled corticosteroids. *Pulm Pharmacol Ther*. 2015;32:130–6. <https://doi.org/10.1016/j.pupt.2015.04.001>
- Artigas A, Camprubi-Rimblas M, Tantinyà N, Bringué J, Guillamat-Prats R, Matthay MA. Inhalation therapies in acute respiratory distress syndrome. *Ann Transl Med*. 2017;5:293. <https://doi.org/10.21037/atm.2017.07.21>
- van Rensen EL, Straathof KC, Veselic-Charvat MA, Zwinderman AH, Bel EH, Sterk PJ. Effect of inhaled steroids on airway hyperresponsiveness, sputum eosinophils, and exhaled nitric oxide levels in patients with asthma. *Thorax*. 1999;54:403–8. <https://doi.org/10.1136/thx.54.5.403>
- Horvath G, Wanner A. Inhaled corticosteroids: effects on the airway vasculature in bronchial asthma. *Eur Respir J*. 2006;27:172–87. <https://doi.org/10.1183/09031936.06.00048605>
- Leading Industry Research Network. Analysis on market demand status and market development trend of budesonide inhalation preparations in China in 2020. 2020 Nov 23 [cited 2022 Jan 24]. <http://www.leadingir.com/datacenter/view/5683.html>
- Gales AC, Seifert H, Gur D, Castanheira M, Jones RN, Sader HS. Antimicrobial susceptibility of *Acinetobacter calcoaceticus*-*Acinetobacter baumannii* complex and *Stenotrophomonas maltophilia* clinical isolates: results from the SENTRY antimicrobial surveillance program (1997–2016). *Open Forum Infect Dis*. 2019;6(Suppl 1):S34–46. <https://doi.org/10.1093/ofid/ofy293>
- European Centre for Disease Prevention and Control. Surveillance of antimicrobial resistance in Europe 2018. Stockholm: The Centre; 2019.
- Hu F, Guo Y, Zhu D, Wang F, Jiang X, Xu Y, et al. CHINET surveillance of bacterial resistance in China: 2018 report. *Chin J Infect Chemother*. 2020;20:1–10.
- Wang F, Zhu D, Hu F, Jiang X, Hu Z, Li Q, et al. 2012 CHINET surveillance of bacterial resistance in China. *Chin J Infect Chemother*. 2013;13:321–30.
- Meng X, Fu J, Zheng Y, Qin W, Yang H, Cao D, et al. Ten-year changes in bloodstream infection with *Acinetobacter baumannii* complex in intensive care units in eastern China: a retrospective cohort study. *Front Med (Lausanne)*. 2021;8:715213. <https://doi.org/10.3389/fmed.2021.715213>
- Chung DR, Song JH, Kim SH, Thamlikitkul V, Huang SG, Wang H, et al.; Asian Network for Surveillance of Resistant Pathogens Study Group. High prevalence of multidrug-resistant nonfermenters in hospital-acquired pneumonia in Asia. *Am J Respir Crit Care Med*. 2011;184:1409–17. <https://doi.org/10.1164/rccm.201102-0349OC>
- Xie J, Yang Y, Huang Y, Kang Y, Xu Y, Ma X, et al. The current epidemiological landscape of ventilator-associated pneumonia in the intensive care unit: a multicenter prospective observational study in China. *Clin Infect Dis*. 2018;67(suppl_2):S153–61. <https://doi.org/10.1093/cid/ciy692>
- Hu FP, Guo Y, Zhu DM, Wang F, Jiang XF, Xu YC, et al. Resistance trends among clinical isolates in China reported from CHINET surveillance of bacterial resistance,

- 2005–2014. *Clin Microbiol Infect.* 2016;22(Suppl 1):S9–14. <https://doi.org/10.1016/j.cmi.2016.01.001>
29. Hu F, Guo Y, Yang Y, Zheng Y, Wu S, Jiang X, et al.; China Antimicrobial Surveillance Network (CHINET) Study Group. Resistance reported from China antimicrobial surveillance network (CHINET) in 2018. *Eur J Clin Microbiol Infect Dis.* 2019;38:2275–81. <https://doi.org/10.1007/s10096-019-03673-1>
 30. Huang H, Chen B, Liu G, Ran J, Lian X, Huang X, et al. A multi-center study on the risk factors of infection caused by multi-drug resistant *Acinetobacter baumannii*. *BMC Infect Dis.* 2018;18:11. <https://doi.org/10.1186/s12879-017-2932-5>
 31. Bassetti M, Righi E, Vena A, Graziano E, Russo A, Peghin M. Risk stratification and treatment of ICU-acquired pneumonia caused by multidrug-resistant/extensively drug-resistant/pandrug-resistant bacteria. *Curr Opin Crit Care.* 2018;24:385–93. <https://doi.org/10.1097/MCC.0000000000000534>
 32. Freire MP, de Oliveira Garcia D, Garcia CP, Campagnari Bueno MF, Camargo CH, Kono Magri ASG, et al. Bloodstream infection caused by extensively drug-resistant *Acinetobacter baumannii* in cancer patients: high mortality associated with delayed treatment rather than with the degree of neutropenia. *Clin Microbiol Infect.* 2016;22:352–8. <https://doi.org/10.1016/j.cmi.2015.12.010>
 33. Yamada K, Yanagihara K, Araki N, Harada Y, Morinaga Y, Akamatsu N, et al. Clinical characteristics of tertiary hospital patients from whom *Acinetobacter calcoaceticus*–*Acinetobacter baumannii* complex strains were isolated. *Intern Med.* 2012;51:51–7. <https://doi.org/10.2169/internalmedicine.51.6018>
 34. Jang TN, Lee SH, Huang CH, Lee CL, Chen WY. Risk factors and impact of nosocomial *Acinetobacter baumannii* bloodstream infections in the adult intensive care unit: a case-control study. *J Hosp Infect.* 2009;73:143–50. <https://doi.org/10.1016/j.jhin.2009.06.007>
 35. Mor A, Thomsen RW, Ulrichsen SP, Sørensen HT. Chronic heart failure and risk of hospitalization with pneumonia: a population-based study. *Eur J Intern Med.* 2013;24:349–53. <https://doi.org/10.1016/j.ejim.2013.02.013>
 36. Ehrmann S, Roche-Campo F, Sferrazza Papa GF, Isabay D, Brochard L, Apiou-Sbirlea G; REVA research network. Aerosol therapy during mechanical ventilation: an international survey. *Intensive Care Med.* 2013;39:1048–56. <https://doi.org/10.1007/s00134-013-2872-5>
 37. Hashemian SM, Mortaz E, Jamaati H, Bagheri L, Mohajerani SA, Garssen J, et al. Budesonide facilitates weaning from mechanical ventilation in difficult-to-wean very severe COPD patients: association with inflammatory mediators and cells. *J Crit Care.* 2018;44:161–7. <https://doi.org/10.1016/j.jcrc.2017.10.045>
 38. Ju YN, Yu KJ, Wang GN. Budesonide ameliorates lung injury induced by large volume ventilation. *BMC Pulm Med.* 2016;16:90. <https://doi.org/10.1186/s12890-016-0251-z>
 39. Yang M, Zhang Y, Chen H, Lin J, Zeng J, Xu Z. Inhaled corticosteroids and risk of upper respiratory tract infection in patients with asthma: a meta-analysis. *Infection.* 2019;47:377–85. <https://doi.org/10.1007/s15010-018-1229-y>
 40. Yang M, Chen H, Zhang Y, Du Y, Xu Y, Jiang P, et al. Long-term use of inhaled corticosteroids and risk of upper respiratory tract infection in chronic obstructive pulmonary disease: a meta-analysis. *Inhal Toxicol.* 2017;29:219–26. <https://doi.org/10.1080/08958378.2017.1346006>
 41. McKeever T, Harrison TW, Hubbard R, Shaw D. Inhaled corticosteroids and the risk of pneumonia in people with asthma: a case-control study. *Chest.* 2013;144:1788–94. <https://doi.org/10.1378/chest.13-0871>
 42. Wang J, Li F, Tian Z. Role of microbiota on lung homeostasis and diseases. *Sci China Life Sci.* 2017;60:1407–15. <https://doi.org/10.1007/s11427-017-9151-1>
 43. Sze MA, Dimitriu PA, Hayashi S, Elliott WM, McDonough JE, Gosselink JV, et al. The lung tissue microbiome in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* 2012;185:1073–80. <https://doi.org/10.1164/rccm.201111-2075OC>
 44. Hilty M, Burke C, Pedro H, Cardenas P, Bush A, Bossley C, et al. Disordered microbial communities in asthmatic airways. *PLoS One.* 2010;5:e8578. <https://doi.org/10.1371/journal.pone.0008578>
 45. Laroumagne S, Lepage B, Hermant C, Plat G, Phelippeau M, Bigay-Game L, et al. Bronchial colonisation in patients with lung cancer: a prospective study. *Eur Respir J.* 2013;42:220–9. <https://doi.org/10.1183/09031936.00062212>
 46. Dickson RP, Singer BH, Newstead MW, Falkowski NR, Erb-Downward JR, Standiford TJ, et al. Enrichment of the lung microbiome with gut bacteria in sepsis and the acute respiratory distress syndrome. *Nat Microbiol.* 2016;1:16113. <https://doi.org/10.1038/nmicrobiol.2016.113>
 47. Dickson RP, Erb-Downward JR, Huffnagle GB. Towards an ecology of the lung: new conceptual models of pulmonary microbiology and pneumonia pathogenesis. *Lancet Respir Med.* 2014;2:238–46. [https://doi.org/10.1016/S2213-2600\(14\)70028-1](https://doi.org/10.1016/S2213-2600(14)70028-1)
 48. Lai Y, Gallo RL. AMPed up immunity: how antimicrobial peptides have multiple roles in immune defense. *Trends Immunol.* 2009;30:131–41. <https://doi.org/10.1016/j.it.2008.12.003>
 49. Ramsheh MY, Haldar K, Esteve-Codina A, Purser LF, Richardson M, Müller-Quernheim J, et al. Lung microbiome composition and bronchial epithelial gene expression in patients with COPD versus healthy individuals: a bacterial 16S rRNA gene sequencing and host transcriptomic analysis. *Lancet Microbe.* 2021;2:e300–10. [https://doi.org/10.1016/S2666-5247\(21\)00035-5](https://doi.org/10.1016/S2666-5247(21)00035-5)
 50. Huang C, Yu Y, Du W, Liu Y, Dai R, Tang W, et al. Fungal and bacterial microbiome dysbiosis and imbalance of trans-kingdom network in asthma. *Clin Transl Allergy.* 2020;10:42. <https://doi.org/10.1186/s13601-020-00345-8>

Address for correspondence: Hao Wang, Department of Critical Care Medicine, Qilu Hospital of Shandong University, 107 Wenhua Rd, Jinan 250012, China; email: wanghao34@126.com

Acinetobacter baumannii among Patients Receiving Glucocorticoid Aerosol Therapy during Invasive Mechanical Ventilation, China

Appendix

Appendix Table 1. Other pathogens detected from MacConkey agar plates in a study of *Acinetobacter baumannii* among patients during invasive mechanical ventilation, China*

Bacteria	Total (%), n = 497	No. of isolates (%)		p value
		Glucocorticoid aerosol therapy, n = 262	No glucocorticoid aerosol therapy, n = 235	
<i>Klebsiella pneumoniae</i>	38 (7.6)	23 (8.8)	15 (6.4)	0.316
<i>Pseudomonas aeruginosa</i>	27 (5.4)	16 (6.1)	11 (4.7)	0.484
<i>Escherichia coli</i>	15 (3.0)	9 (3.4)	6 (2.6)	0.566
<i>Stenotrophomonas maltophilia</i>	7 (1.4)	4 (1.5)	3 (1.3)	0.885
<i>Burkholderia cepacia</i>	3 (0.6)	1 (0.4)	2 (0.9)	0.925
<i>Serratia marcescens</i>	2 (0.4)	1 (0.4)	1 (0.4)	0.527
Others	4 (0.8)	3 (1.1)	1 (0.4)	0.694

Appendix Table 2. Indications for glucocorticoid aerosol therapy in a study of *Acinetobacter baumannii* among patients during invasive mechanical ventilation, China*

Variables	Total, n = 262	<i>A. baumannii</i> isolated,		p value	Hazard ratio (95% CI)
		n = 107	No <i>A. baumannii</i> isolated, n = 155		
Asthma and COPD	106 (21.3)	41 (25.8)	65 (18.9)	0.087	1.400 (0.952–2.059)
ARDS	112 (22.5)	44 (27.7)	68 (20.1)	0.076	1.408 (0.965–2.052)
Other conditions†	30 (6.0)	14 (8.8)	16 (4.7)	0.112	1.592 (0.897–2.827)
Undetermined	14 (2.8)	8 (5.0)	6 (1.8)	0.115	1.801 (0.867–3.744)

*Data are expressed as no. (%). ARDS, acute respiratory distress syndrome; COPD, chronic obstructive pulmonary disease.

†Other conditions included recorded airway hyperresponsiveness, hyperinflammation or mucosal edema following diseases other than asthma, COPD, and ARDS.

Appendix Table 3. Multivariate analysis of risk factors for *Acinetobacter baumannii* isolation among patients during invasive mechanical ventilation, China*

Variables	p value	Hazard ratio (95% CI)
Age, y	0.987	1.000 (0.988–1.012)
Sex		
F	–	Referent
M	0.748	0.936 (0.624–1.403)
Charlson comorbidity index	0.115	0.920 (0.830–1.020)
Underlying conditions		
Cardiovascular diseases	0.031	1.530 (1.040–2.250)
Chronic renal insufficiency	0.703	1.080 (0.727–1.605)
COPD and asthma	0.082	1.393 (0.959–2.024)
Type 2 diabetes mellitus	0.108	1.379 (0.932–2.040)
Solid tumor	0.084	1.455 (0.950–2.229)
Hematologic malignancy	0.252	0.655 (0.317–1.352)
Past inhaled steroid usage for chronic conditions	0.654	0.868 (0.467–1.612)
Current or former smoker	0.191	1.273 (0.886–1.828)
Postoperative admission	0.581	0.899 (0.616–1.312)
Treatment		
No aerosol inhalation	–	Referent
Glucocorticoid aerosol therapy	0.049	1.499 (1.001–2.246)
Aerosol inhalation without glucocorticoid	0.571	0.846 (0.474–1.508)
Broad-spectrum antimicrobial drugs for ≥ 7 d	0.001	9.436 (2.803–18.661)
Invasive mechanical ventilation for ≥ 5 d	<0.001	2.325 (1.616–3.344)
Urethral catheter placement for ≥ 3 d	0.737	0.707 (0.093–5.352)
Vasopressor treatment for ≥ 3 d	<0.001	2.034 (1.379–3.001)
Renal dialysis for ≥ 3 d	0.746	1.077 (0.687–1.688)
APACHE II score	0.755	0.995 (0.967–1.025)

*Results are based on model 1 in which all variables are included. APACHE II, Acute Physiology and Chronic Health Evaluation II; COPD, chronic obstructive pulmonary disease; –, not applicable.

Appendix Table 4. Multivariate analysis of risk factors for *Acinetobacter baumannii* isolation among patients during invasive mechanical ventilation, China*

Variables	Model 1		Model 2	
	p value	Hazard ratio (95% CI)	p value	Hazard ratio (95% CI)
Age, y	0.947	1.000 (0.988–1.013)	ND	ND
Sex				
F	–	Referent	ND	ND
M	0.633	0.907 (0.608–1.354)	ND	ND
Charlson comorbidity index	0.086	0.913 (0.824–1.013)	ND	ND
Underlying conditions				
Cardiovascular diseases	0.036	1.510 (1.027–2.219)	0.060	1.386 (0.987–1.948)
Chronic renal insufficiency	0.609	1.110 (0.745–1.653)	0.782	0.949 (0.654–1.376)
COPD and asthma	0.087	1.387 (0.954–2.019)	0.121	1.309 (0.931–1.839)
Type 2 diabetes mellitus	0.109	1.379 (0.931–2.043)	0.358	1.183 (0.826–1.694)
Solid tumor	0.075	1.476 (0.962–2.266)	ND	ND
Hematologic malignancy	0.205	0.626 (0.303–1.292)	ND	ND
Past inhaled steroid usage for chronic conditions	0.824	0.932 (0.504–1.726)	ND	ND
Current or former smoker	0.215	1.255 (0.877–1.795)	0.107	1.299 (0.945–1.785)
Postoperative admission	0.462	0.868 (0.595–1.266)	ND	ND
Treatment				
No aerosol inhalation	–	Referent	–	Referent
Aerosol inhalation	0.106	1.391 (0.932–2.077)	0.089	1.413 (0.949–2.103)
Broad-spectrum antimicrobial drugs for ≥ 7 d	0.001	8.845 (2.903–18.339)	0.001	8.591 (2.844–17.246)
Invasive mechanical ventilation for ≥ 5 d	0.001	2.386 (1.659–3.431)	0.001	2.442 (1.707–3.494)
Urethral catheter placement for ≥ 3 d	0.671	0.645 (0.086–4.859)	ND	ND
Vasopressor treatment for ≥ 3 d	<0.001	2.141 (1.453–3.154)	<0.001	2.157 (1.468–3.169)
Renal dialysis for ≥ 3 d	0.614	1.122 (0.717–1.758)	0.726	1.081 (0.698–1.675)
APACHE II score	0.628	0.993 (0.964–1.022)	0.522	0.991 (0.964–1.019)

*We considered aerosol inhalation as a whole variable. Model 1 includes all variables; model 2 only includes variables for which $p < 0.1$ in univariate analysis. APACHE II, Acute Physiology and Chronic Health Evaluation II; COPD, chronic obstructive pulmonary disease; ND, not done; –, not applicable.

Appendix Table 5. Univariate analysis of patients with or without glucocorticoid aerosol therapy after propensity score adjustment in a study of *Acinetobacter baumannii* isolation among patients during invasive mechanical ventilation, China*

Variables	Glucocorticoid aerosol therapy, n = 199	No glucocorticoid aerosol therapy, n = 199	p value
Median age, y (IQR)	59.9 (49–72)	60.1 (49–72)	0.899
Sex, no. (%)			
F	61 (30.7)	64 (32.2)	–
M	138 (69.3)	135 (67.8)	0.527
Mean Charlson comorbidity index (SD)	4.18 (2.04)	4.23 (2.19)	0.795
Underlying conditions, no. (%)			
Cardiovascular diseases	79 (39.7)	79 (39.7)	1.000
Chronic renal insufficiency	73 (36.7)	68 (34.2)	0.600
COPD and asthma	65 (32.7)	68 (34.2)	0.750
Type 2 diabetes mellitus	46 (23.1)	48 (24.1)	0.813
Solid tumor	37 (18.6)	39 (19.6)	0.799
Hematologic malignancy	14 (7.0)	12 (6.0)	0.685
Past inhaled steroid usage for chronic conditions	17(8.5)	15 (7.5)	0.712
Current or former smoker	75 (37.7)	73 (36.7)	0.836
Postoperative admission	59 (29.6)	58 (29.1)	0.912
Treatment, no. (%)			
Broad-spectrum antimicrobial drugs for ≥ 7 d	170 (85.4)	167 (83.9)	0.676
Invasive mechanical ventilation for ≥ 5 d	84 (42.2)	82 (41.2)	0.839
Urethral catheter placement for ≥ 3 d	197 (99.0)	198 (99.5)	0.562
Vasopressor treatment for ≥ 3 d	23 (11.6)	25 (12.6)	0.758
Renal dialysis for ≥ 3 d	37 (18.6)	32 (16.1)	0.508
Mean APACHE II score (SD)	18.47 (6.23)	18.05 (5.72)	0.485
<i>Acinetobacter baumannii</i> isolated, no (%)	70 (35.2)	51 (25.6)	0.038
Median length of ICU stay, d (IQR)	21 (9–24)	18 (5–21)	0.262

*APACHE II, Acute Physiology and Chronic Health Evaluation II; COPD, chronic obstructive pulmonary disease; ICU, intensive care unit; IQR, interquartile range; SD, standard deviation; –, not applicable.

Appendix Table 6. Univariate analysis of risk factors for *Acinetobacter baumannii* isolation among patients during invasive mechanical ventilation, China*

Variables	No		p value	Hazard ratio (95% CI)
	<i>A. baumannii</i> isolated, n = 121	<i>A. baumannii</i> isolated, n = 277		
Median age, y (IQR)	59.2 (49–71)	60.4 (49–73)	0.460	0.996 (0.986–1.006)
Sex, no. (%)				
F	35 (28.9)	90 (32.5)	–	Referent
M	86 (71.1)	187 (67.5)	0.527	0.881 (0.595–1.305)
Mean Charlson comorbidity index (SD)	4.36 (2.06)	4.14 (2.14)	0.295	1.043 (0.964–1.129)
Underlying conditions, no. (%)				
Cardiovascular diseases	58 (47.9)	100 (36.1)	0.021	1.520 (1.064–2.172)
Chronic renal insufficiency	57 (47.1)	84 (30.3)	0.002	1.750 (1.224–2.501)
COPD and asthma	47 (38.8)	86 (31.0)	0.171	1.291 (0.896–1.861)
Type 2 diabetes mellitus	37 (30.6)	57 (20.6)	0.038	1.505 (1.022–2.215)
Solid tumor	25 (20.7)	51 (18.4)	0.563	0.878 (0.565–1.364)
Hematologic malignancy	6 (5.0)	20 (7.2)	0.450	0.729 (0.321–1.657)
Past inhaled steroid usage for chronic conditions	7 (5.8)	25 (9.0)	0.280	0.657 (0.306–1.409)
Current or former smoker	57 (47.1)	91 (32.9)	0.008	1.627 (1.138–2.325)
Postoperative admission	32 (26.4)	85 (30.7)	0.441	0.853 (0.570–1.278)
Treatment, no. (%)				
Glucocorticoid aerosol therapy	70 (57.9)	129 (46.6)	0.038	1.438 (1.003–2.061)
Broad-spectrum antimicrobial drugs for ≥ 7 d	120 (99.2)	217 (78.3)	0.015	7.454 (1.925–15.582)
Invasive mechanical ventilation for ≥ 5 d	80 (66.1)	86 (31.0)	<0.001	3.203 (1.927–5.323)
Urethral catheter placement for ≥ 3 d	120 (99.2)	275 (99.3)	0.472	0.485 (0.067–3.494)
Vasopressor treatment for ≥ 3 d	19 (15.7)	29 (10.5)	0.010	2.224 (1.215–4.069)
Renal dialysis for ≥ 3 d	29 (24.0)	40 (14.4)	0.112	1.559 (0.902–2.697)
Mean APACHE II score (SD)	19.62 (6.51)	17.66 (5.63)	0.002	1.056 (1.019–1.094)
Median length of ICU stay, d (IQR)	22 (10–28)	18 (6–21)	0.205	1.004 (0.998–1.011)

*We used a propensity-matched cohort for analyses. APACHE II, Acute Physiology and Chronic Health Evaluation II; COPD, chronic obstructive pulmonary disease; ICU, intensive care unit; IQR, interquartile range; SD, standard deviation; –, not applicable.

Appendix Table 7. Multivariate analysis of risk factors for *Acinetobacter baumannii* isolation among patients during invasive mechanical ventilation, China*

Variables	p value	Hazard ratio (95% CI)
Underlying conditions		
Cardiovascular diseases	0.010	1.530 (1.109–2.112)
Chronic renal insufficiency	0.852	0.967 (0.676–1.382)
Type 2 diabetes mellitus	0.654	1.084 (0.761–1.544)
Current or former smoker	0.099	1.292 (0.953–1.753)
Treatment		
No aerosol inhalation	–	Referent
Glucocorticoid aerosol for asthma and COPD	0.024	1.774 (1.077–2.920)
Glucocorticoid aerosol for ARDS	0.003	3.589 (1.955–6.589)
Glucocorticoid aerosol for other conditions or undetermined	0.060	1.705 (0.990–2.669)
Aerosol inhalation without glucocorticoid	0.170	0.690 (0.406–1.172)
Broad-spectrum antimicrobial drugs for ≥ 7 d	0.001	4.677 (2.036–10.743)
Invasive mechanical ventilation for ≥ 5 d	<0.001	2.247 (1.588–3.179)
Vasopressor treatment for ≥ 3 d	0.001	2.703 (1.798–4.066)
Renal dialysis for ≥ 3 d	0.643	1.107 (0.720–1.702)
APACHE II score	0.256	0.985 (0.959–1.011)

*Calculations include indications for glucocorticoid aerosol therapy and only include variables with $p < 0.1$ in univariate analysis. APACHE II, Acute Physiology and Chronic Health Evaluation II; ARDS, acute respiratory distress syndrome; COPD, chronic obstructive pulmonary disease; ICU, intensive care unit; –, not applicable.

Appendix Table 8. Univariate analysis of risk factors for 30-d mortality in a study of *Acinetobacter baumannii* isolation among patients during invasive mechanical ventilation, China*

Variables	Total, n = 497	Died, n = 100	Survived, n = 397	p value	Hazard ratio (95% CI)
Median age, y (IQR)	60.1 (49–73)	59 (48–76)	61 (49–73)	0.673	0.998 (0.986–1.009)
Sex, no. (%)					
F	160 (32.2)	34 (34.0)	126 (31.7)	–	Referent
M	337 (67.8)	66 (66.0)	271 (68.3)	0.716	1.080 (0.714–1.633)
Mean Charlson comorbidity index (SD)	4.26 (2.12)	4.58 (1.96)	4.18 (2.15)	0.082	1.075 (0.991–1.167)
Underlying conditions, no. (%)					
Cardiovascular diseases	200 (40.2)	46 (46.0)	154 (38.8)	0.098	1.329 (0.897–1.970)
Chronic renal insufficiency	179 (36.0)	51 (51.0)	128 (32.2)	0.001	2.004 (1.354–2.967)
COPD and asthma	176 (35.4)	39 (39.0)	137 (34.5)	0.331	1.220 (0.817–1.824)
Type 2 diabetes mellitus	116 (23.3)	25 (25.0)	91 (22.9)	0.521	1.160 (0.738–1.824)
Solid tumor	100 (20.1)	23 (23.0)	73 (18.4)	0.327	1.262 (0.792–2.011)
Hematologic malignancy	31 (6.1)	11 (10.4)	20 (5.0)	0.022	2.075 (1.109–3.884)
Past inhaled steroid usage for chronic conditions	47 (9.5)	13 (13.0)	34 (8.6)	0.153	1.530 (0.854–2.740)
Current or former smoker	187 (37.6)	37 (37.0)	150 (37.8)	0.839	0.959 (0.639–1.439)
Postoperative admission	142 (28.6)	21 (21.0)	121 (30.5)	0.100	0.649 (0.401–1.050)
Treatment, no. (%)					
No aerosol inhalation	137 (27.6)	33 (33.0)	104 (26.2)	–	Referent
Glucocorticoid aerosol therapy	262 (52.7)	49 (49.0)	213 (53.7)	0.087	0.753 (0.484–1.170)
Aerosol inhalation without glucocorticoid	98 (19.7)	18 (18.0)	80 (20.2)	0.389	0.777 (0.438–1.380)
Broad-spectrum antimicrobial drugs for ≥ 7 d	417 (83.9)	85 (85.0)	332 (83.6)	0.928	1.026 (0.592–1.778)
Invasive mechanical ventilation for ≥ 5 d	220 (44.3)	62 (62.0)	158 (39.8)	0.001	2.079 (1.388–3.114)
Urethral catheter placement for ≥ 3 d	493 (99.2)	99 (99.0)	394 (99.2)	0.701	0.950 (0.680–4.873)
Vasopressor treatment for ≥ 3 d	75 (15.1)	52 (52.0)	23 (5.8)	<0.001	8.988 (6.048–13.357)
Renal dialysis for ≥ 3 d	84 (16.9)	22 (22.0)	62 (15.6)	0.163	1.401 (0.873–2.248)
Mean APACHE II score (SD)	18.18 (6.03)	20.07 (7.10)	17.70 (5.64)	0.001	1.056 (1.025–1.089)
<i>A. baumannii</i> isolation, no. (%)	159 (32.0)	47 (47.0)	112 (28.2)	0.001	1.977 (1.335–2.929)
<i>A. baumannii</i> isolation status, no. (%)					
No <i>A. baumannii</i> isolation	338 (68.0)	53 (53.0)	285 (71.8)	–	Referent
<i>A. baumannii</i> colonization	96 (19.3)	21 (21.0)	75 (18.9)	0.251	1.228 (0.720–2.097)
<i>A. baumannii</i> infection	58 (11.7)	25 (25.0)	33 (8.3)	0.001	3.346 (2.115–5.293)
<i>A. baumannii</i> colonization or infection not defined	5 (1.0)	1 (1.0)	4 (1.0)	0.755	1.371 (0.190–9.911)
Antimicrobial drugs treatment for <i>A. baumannii</i>	65 (13.1)	18 (18.0)	47 (11.8)	0.102	1.565 (0.943–3.297)

*APACHE II, Acute Physiology and Chronic Health Evaluation II; ARDS, acute respiratory distress syndrome; COPD, chronic obstructive pulmonary disease; –, not applicable.

Appendix Table 9. Multivariate analysis of risk factors for 30-d mortality in a study of *Acinetobacter baumannii* isolation among patients during invasive mechanical ventilation, China*

Variables	p value	Hazard ratio (95% CI)
Charlson comorbidity index	0.380	0.953 (0.855–1.062)
Underlying conditions		
Cardiovascular diseases	0.491	1.164 (0.756–1.791)
Chronic renal insufficiency	0.420	1.208 (0.763–1.914)
Hematologic malignancy	0.154	1.616 (0.836–3.127)
Treatment		
No aerosol inhalation	–	Referent
Glucocorticoid aerosol therapy	0.102	0.750 (0.497–1.065)
Aerosol inhalation without glucocorticoid	0.388	1.308 (0.711–2.408)
Invasive mechanical ventilation for ≥ 5 d	0.386	1.226 (0.774–1.942)
Vasopressor treatment for ≥ 3 d	<0.001	8.366 (5.314–13.170)
APACHE II score	0.114	1.026 (0.994–1.060)
<i>A. baumannii</i> isolation	0.045	1.824 (1.317–2.104)

*Calculations include *A. baumannii* isolation as a whole variable and only variables with $p < 0.1$ in univariate analysis were included. APACHE II, Acute Physiology and Chronic Health Evaluation II; –, not applicable.

Appendix Table 10. Multivariate analysis of risk factors for 30-d mortality in a study of *Acinetobacter baumannii* isolation among patients during invasive mechanical ventilation, China*

Variables	p value	Hazard ratio (95% CI)
Charlson comorbidity index	0.490	0.962 (0.863–1.073)
Underlying conditions		
Cardiovascular diseases	0.941	1.016 (0.657–1.572)
Chronic renal insufficiency	0.491	1.176 (0.741–1.869)
Hematologic malignancy	0.132	1.657 (0.859–3.195)
Treatment		
No aerosol inhalation	–	Referent
Glucocorticoid aerosol therapy	0.106	0.747 (0.493–1.066)
Aerosol inhalation without glucocorticoid	0.404	1.296 (0.705–2.382)
Invasive mechanical ventilation for ≥ 5 d	0.729	1.088 (0.675–1.756)
Vasopressor treatment for ≥ 3 d	0.001	8.989 (5.720–14.126)
APACHE II score	0.148	1.023 (0.992–1.055)
<i>A. baumannii</i> isolation status		
No <i>A. baumannii</i> isolation	–	Referent
<i>A. baumannii</i> colonization	0.720	0.894 (0.484–1.650)
<i>A. baumannii</i> infection	0.012	2.759 (1.575–4.833)
<i>A. baumannii</i> colonization or infection not defined	0.488	0.494 (0.067–3.634)

* Calculations include four *A. baumannii* isolation status and only variables with $p < 0.1$ in univariate analysis were included. APACHE II, Acute Physiology and Chronic Health Evaluation II; –, not applicable.