References

- Corey L, Beyrer C, Cohen MS, Michael NL, Bedford T, Rolland M. SARS-CoV-2 variants in patients with immunosuppression. N Engl J Med. 2021;385:562–6. https://doi.org/10.1056/NEJMsb2104756
- Wang Y, Wang D, Zhang L, Sun W, Zhang Z, Chen W, et al. Intra-host variation and evolutionary dynamics of SARS-CoV-2 populations in COVID-19 patients. Genome Med. 2021;13:30. https://doi.org/10.1186/ s13073-021-00847-5
- 3. Khateeb D, Gabrieli T, Sofer B, Hattar A, Cordela S, Chaouat A, et al. SARS-CoV-2 variants with reduced infectivity and varied sensitivity to the BNT162b2 vaccine are developed during the course of infection. PLoS Pathog. 2022;18:e1010242. https://doi.org/10.1371/ journal.ppat.1010242
- Weigang S, Fuchs J, Zimmer G, Schnepf D, Kern L, Beer J, et al. Within-host evolution of SARS-CoV-2 in an immunosuppressed COVID-19 patient as a source of immune escape variants. Nat Commun. 2021;12:6405. https://doi.org/10.1038/s41467-021-26602-3
- Cele S, Karim F, Lustig G, San JE, Hermanus T, Tegally H, et al.; COMMIT-KZN Team. SARS-CoV-2 prolonged infection during advanced HIV disease evolves extensive immune escape. Cell Host Microbe. 2022;30:154–162.e5. https://doi.org/10.1016/j.chom.2022.01.005
- Choi B, Choudhary MC, Regan J, Sparks JA, Padera RF, Qiu X, et al. Persistence and evolution of SARS-CoV-2 in an immunocompromised host. N Engl J Med. 2020;383:2291–3. https://doi.org/10.1056/NEJMc2031364
- Pérez-Lago L, Aldámiz-Échevarría T, García-Martínez R, Pérez-Latorre L, Herranz M, Sola-Campoy PJ, et al.; on behalf of Gregorio Marañón Microbiology-Id Covid Study Group. Different within-host viral evolution dynamics in severely immunosuppressed cases with persistent SARS-CoV-2. Biomedicines. 2021;9:808. https://doi.org/10.3390/ biomedicines9070808
- Avanzato VA, Matson MJ, Seifert SN, Pryce R, Williamson BN, Anzick SL, et al. Case study: prolonged infectious SARS-CoV-2 shedding from an asymptomatic immunocompromised individual with cancer. Cell. 2020;183:1901–12.e9. https://doi.org/10.1016/ j.cell.2020.10.049
- Truffot A, Andréani J, Le Maréchal M, Caporossi A, Epaulard O, Germi R, et al. SARS-CoV-2 variants in immunocompromised patient given antibody monotherapy. Emerg Infect Dis. 2021;27:2725–8. https://doi.org/10.3201/ eid2710.211509

Address for correspondence: Slim Fourati, Henri Mondor Hospital, 51 av du Marechal de Lattre de Tassigny, Créteil 94010, France; email: slim.fourati@aphp.fr

Increased Stability of SARS-CoV-2 Omicron Variant over Ancestral Strain

Alex Wing Hong Chin, Alison Man Yuk Lai, Malik Peiris, Leo Lit Man Poon

Author affiliations: The University of Hong Kong, Hong Kong, China (A.W.H. Chin, A.M.Y. Lai, M. Peiris, L.L.M. Poon); Centre for Immunology & Infection, Hong Kong (A.W.H. Chin, M. Peiris, L.L.M. Poon)

DOI: https://doi.org/10.3201/eid2807.220428

As of April 2022, the Omicron BA.1 variant of concern of SARS-CoV-2 was spreading quickly around the world and outcompeting other circulating strains. We examined its stability on various surfaces and found that this Omicron variant is more stable than its ancestral strain on smooth and porous surfaces.

The Omicron SARS-CoV-2 variant of concern (VOC) is highly transmissible in humans. As of April 2022, it has outcompeted other known variants and dominated in different regions (1). Its spike protein has >30 mutations compared with the ancestral strain (2). A 2022 structural study indicates the Omicron spike protein is more stable than that of the ancestral strain (3); this finding prompted us to hypothesize that Omicron VOC is also more stable on different surfaces. We previously showed that the ancestral SARS-CoV-2 strain can still be infectious at room temperature for several days on smooth surfaces and several hours on porous surfaces (4).

We used previously described ancestral SARS-CoV-2 (PANGO lineage A) and Omicron VOC (PANGO lineage BA.1) in this study (5,6). We tested their stability on different surfaces using our previously described protocol (4,7). In brief, we applied a 5-µL droplet of each virus (10^7 50% tissue culture infectious dose [TCID₅₀]/mL) on different surfaces in triplicate. We incubated the treated surfaces at room temperature (21° C- 22° C) for different time points as indicated and then immersed them in viral transport medium for 30 min to recover the residual infectious virus. We titrated the recovered virus by TCID₅₀ assays using Vero E6 cells, as described (4,7).

Compared with the ancestral SARS-CoV-2, the Omicron BA.1 variant was more stable on all surfaces we studied (Table). On day 4 postinoculation, we recovered no infectious ancestral SARS-CoV-2 from stainless steel, polypropylene sheet, or 2 of 3 glass

RESEARCH LETTERS

Table. Stability of ancestral SARS-Cov-2 and of Officion variant of different surfaces								
		Ancestral SARS-CoV-2		Omicron variant				
	Incubation	Mean log ₁₀ (TCID ₅₀ /mL)	% Reduction	Mean log ₁₀ (TCID ₅₀ /mL)	% Reduction			
Material	time†	±SD‡	in viral titer	<u>+</u> SD‡	in viral titer			
Stainless steel	0	5.02 <u>+</u> 0.39	NA	5.35 <u>+</u> 0.18	NA			
	3 h	4.21 <u>+</u> 0.36	85.15	4.82 +0.23	69.78			
	6 h	3.73 <u>+</u> 0.10	95.80	4.62 <u>+</u> 0.31	79.86			
	1 d	2.99 <u>+</u> 0.17	99.21	4.65 <u>+</u> 0.17	80.28			
	2 d	2.08 <u>+</u> 0.11	99.91	4.51 <u>+</u> 0.15	85.82			
	4 d	§	>99.93	3.72 <u>+</u> 0.12	97.72			
	7 d	§	>99.93	3.58 <u>+</u> 0.30	98.19			
Polypropylene	0	4.85 <u>+</u> 0.23	NA	5.43 <u>+</u> 0.16	NA			
	3 h	4.12 <u>+</u> 0.19	81.72	4.65 <u>+</u> 0.34	81.27			
	6 h	3.53 <u>+</u> 0.15	95.43	4.33 <u>+</u> 0.14	92.34			
	1 d	3.13 <u>+</u> 0.34	97.86	4.45 <u>+</u> 0.23	89.25			
	2 d	2.01 <u>+</u> 0.10¶	>99.86	4.34 <u>+</u> 0.25	91.53			
	4 d	§	>99.88	3.97 <u>+</u> 0.19	96.48			
	7 d	§	>99.88	2.95 <u>+</u> 0.27	99.65			
Glass	0	5.10 <u>+</u> 0.24	NA	5.65 <u>+</u> 0.28	NA			
	3 h	4.26 <u>+</u> 0.05	86.79	4.90 <u>+</u> 0.15	83.62			
	6 h	3.69 <u>+</u> 0.11	96.42	4.52 <u>+</u> 0.13	93.20			
	1 d	2.83 <u>+</u> 0.13	99.49	4.20 <u>+</u> 0.01	96.84			
	2 d	2.14 <u>+</u> 0.13	99.90	4.43 <u>+</u> 0.29	93.87			
	4 d	1.96 <u>+</u> 0.00¶	>99.93	4.06 <u>+</u> 0.16	97.64			
	7 d	§	>99.93	3.76 <u>+</u> 0.10	98.83			
Tissue paper	0	4.70 <u>+</u> 0.22	NA	5.21 <u>+</u> 0.14	NA			
	5 min	3.85 <u>+</u> 0.28	84.98	4.64 <u>+</u> 0.70	53.94			
	15 min	2.12 <u>+</u> 0.14	99.75	3.72 <u>+</u> 1.22	72.99			
	30 min	§	>99.84	2.92 <u>+</u> 0.40	99.34			
	60 min	§	>99.84	§	>99.95			
Printing paper	0	5.21 <u>+</u> 0.00	NA	5.34 <u>+</u> 0.13	NA			
	5 min	2.69 <u>+</u> 0.16	99.68	3.26 <u>+</u> 0.42	98.91			
	15 min	§	>99.94	2.20 <u>+</u> 0.33¶	>99.91			
	30 min	§	>99.94	2.16 <u>+</u> 0.36¶	>99.92			
	60 min	§	>99.94	§	>99.96			

Table. Stability of ancestral SARS-CoV-2 and of Omicron variant on different surfaces*

*Tests were performed in triplicate. NA, not applicable; TCID₅₀, 50% tissue culture infectious dose.

†The samples were incubated at room temperature (21°C-22°C).

‡Vero E6 cells were used for titration of viable viruses.

SAll the triplicates were below detection limit of the TCID₅₀ assay.

¶One or two out of three replicates were below detection limit of the TCID₅₀ assay.

samples. We did not recover infectious virus from glass on day 7. In contrast, infectious Omicron variant was still recoverable from all treated surfaces on day 7 postincubation.

The stability of the Omicron variant was also higher than ancestral SARS-CoV-2 on porous surfaces, such as tissue paper and printing paper. On tissue paper, viable ancestral SARS-CoV-2 was no longer recoverable after a 30-minute incubation. However, we detected viable Omicron variant after a 30-minute incubation. On printing paper, we detected no infectious virus after a 15-minute incubation. In contrast, viable Omicron variant was recovered from 2 of 3 replicates after a 30-minute incubation.

To confirm our observations, we used transmembrane serine protease 2 (TMPRSS2)-expressing Vero E6 cells to titrate infectious virus particles recovered from treated stainless steel and printing paper (Appendix Table, https://wwwnc.cdc.gov/EID/ article/28/7/22-0428-App1.pdf). On stainless steel, infectious ancestral virus was undetectable on day 10 postincubation, whereas viable Omicron variant was still recoverable. Similarly, no infectious ancestral virus was detected on printing paper after a 30-minute incubation, whereas we detected viable Omicron variant in 1 out of 3 replicates. Although the virus could be trapped in the porous materials and inefficiently recovered, our findings confirm that Omicron variant is more stable than its ancestral strain on surfaces.

We noted that the cell line used for virus titration can affect our findings. It has been reported that Omicron variant is less dependent upon TMPRSS2 for cell entry (ϑ); therefore, we were not surprised that different cell lines led to different viral inactivation profiles. Nonetheless, results from both cell lines suggest that the Omicron variant is more stable than the ancestral strain. This observation is consistent with other recent findings (R. Hirose et al., unpub. data, https://www. biorxiv.org/content/10.1101/2022.01.18.476607v1). More evidence is needed to account for the increased transmissibility of Omicron variant. The virus's stability on surfaces may be one factor and should be taken

into consideration when recommending control measures against infection. A recent study revealed that an infectious dose as low as 10 TCID₅₀ units could infect >50% of human study participants (9). Our findings indicate that Omicron variant has an increased likelihood for transmission by the fomite route; they may also indicate that the enhanced stability deduced from structural studies (3) and now demonstrated on different surfaces may be relevant for droplet or aerosol transmission of SARS-CoV-2. Of interest, stability of avian influenza A(H5N1) viruses has been shown to have an association with transmissibility of avian influenza virus between mammals by the airborne route, although the mechanisms underlying this association are not fully understood (10). Further studies on the stability of Omicron variant and its emerging subvariants in droplets and aerosols are warranted.

One limitation of our study is that the experiments were conducted in a well-controlled laboratory environment. Variations in environmental conditions would affect the rate of viral inactivation. Therefore, the time required for virus inactivation that we demonstrated may not reflect all real-life scenarios. In addition, the components of the viral droplet medium applied in this study were different from those of the respiratory droplets, which could also affect the stability of the virus. Nonetheless, our findings demonstrate that the Omicron variant is more stable than the ancestral SARS-CoV-2 on different surfaces, a finding that may be relevant for determining recommendations for public health measures to limit virus transmission.

Acknowledgments

This work was supported by the Hong Kong Health and Medical Research Fund (no. COVID190116) and Theme-based Research Scheme of Research Grant Council (no. T11-705/21-N), and InnoHK grants for Centre for Immunology & Infection.

About the Author

Dr. Chin is currently a research assistant professor of the University of Hong Kong. His research interests include basic virology and pathogenesis of emerging respiratory viruses.

References

- World Health Organization. Coronavirus update 74: addressing the challenges of SARS-CoV-2 variants for public health. 2022 25 Feb [cited 2022 Apr 29]. https://www.who. int/publications/m/item/update-74-addressing-the-challenges-of-sars-cov-2-variants-for-public-health
- Wang L, Cheng G. Sequence analysis of the emerging SARS-CoV-2 variant Omicron in South Africa. J Med Virol. 2022;94:1728–33. https://doi.org/10.1002/jmv.27516
- Cui Z, Liu P, Wang N, Wang L, Fan K, Zhu Q, et al. Structural and functional characterizations of infectivity and immune evasion of SARS-CoV-2 Omicron. Cell. 2022;185:860–871.e13. https://doi.org/10.1016/ j.cell.2022.01.019
- Chin AWH, Chu JTS, Perera MRA, Hui KPY, Yen HL, Chan MCW, et al. Stability of SARS-CoV-2 in different environmental conditions. Lancet Microbe. 2020;1:e10. https://doi.org/10.1016/S2666-5247(20)30003-3
- Sia SF, Yan LM, Chin AWH, Fung K, Choy KT, Wong AYL, et al. Pathogenesis and transmission of SARS-CoV-2 in golden hamsters. Nature. 2020;583:834–8. https://doi.org/10.1038/s41586-020-2342-5
- Cheng SMS, Mok CKP, Leung YWY, Ng SS, Chan KCK, Ko FW, et al. Neutralizing antibodies against the SARS-CoV-2 Omicron variant BA.1 following homologous and heterologous CoronaVac or BNT162b2 vaccination. Nat Med. 2022;28:486–9. https://doi.org/10.1038/s41591-022-01704-7
- Behzadinasab S, Chin A, Hosseini M, Poon L, Ducker WA. A surface coating that rapidly inactivates SARS-CoV-2. ACS Appl Mater Interfaces. 2020;12:34723-7. https://doi.org/10.1021/acsami.0c11425
- Meng B, Abdullahi A, Ferreira IATM, Goonawardane N, Saito A, Kimura I, et al.; CITIID-NIHR BioResource COVID-19 Collaboration; Genotype to Phenotype Japan (G2P-Japan) Consortium; Ecuador-COVID19 Consortium. Altered TMPRSS2 usage by SARS-CoV-2 Omicron impacts infectivity and fusogenicity. Nature. 2022;603:706– 14. https://doi.org/10.1038/s41586-022-04474-x
- Killingley B, Mann AJ, Kalinova M, Boyers A, Goonawardane N, Zhou J, et al. Safety, tolerability and viral kinetics during SARS-CoV-2 human challenge in young adults. Nat Med. 2022. https://doi.org/10.1038/ s41591-022-01780-9
- Imai M, Herfst S, Sorrell EM, Schrauwen EJ, Linster M, De Graaf M, et al. Transmission of influenza A/H5N1 viruses in mammals. Virus Res. 2013;178:15–20. https://doi.org/10.1016/j.virusres.2013.07.017

Address for correspondence: Alex Wing Hong Chin, School of Public Health, The University of Hong Kong, Hong Kong; email: alexchin@hku.hk; Leo Lit Man Poon, School of Public Health, The University of Hong Kong, Hong Kong; email: llmpoon@hku.hk

Increased Stability of SARS-CoV-2 Omicron Variant over Ancestral Strain

Appendix

Appendix Table. Stability of the ancestral SARS-CoV-2 and Omicron variant on stainless steel and printing paper, titrated with TMPRSS2-expressing Vero-E6 cells*

		Ancestral SARS-CoV-2		Omicron variant	
	Time of	Mean	% Reduction in viral	Mean	% Reduction in
Materials	incubation†	log ₁₀ (TCID ₅₀ /mL)±SD‡	titer	log₁₀(TCID₅₀/mL <u>)+</u> SD‡	viral titer
Stainless steel	0	5.15 <u>+</u> 0.17	NA	5.45 <u>+</u> 0.08	NA
	6 h	4.48 <u>+</u> 0.21	77.86	4.82 <u>+</u> 0.13	76.64
	1 d	4.07 <u>+</u> 0.23	91.35	4.58 <u>+</u> 0.21	85.95
	2 d	3.75 <u>+</u> 0.11	96.14	4.48 <u>+</u> 0.32	87.78
	4 d	3.20 <u>+</u> 0.14	98.89	3.89 <u>+</u> 0.11	97.24
	7 d	2.67 <u>+</u> 0.12	99.68	3.37 <u>+</u> 0.29	99.04
	10 d	§	>99.94	2.55 <u>+</u> 0.08	99.88
Printing paper	0	4.32 <u>+</u> 0.04	NA	4.62 <u>+</u> 0.48	NA
	5 min	2.92 <u>+</u> 0.27	95.53	3.64 <u>+</u> 0.31	90.95
	15 min	2.09 <u>+</u> 0.12¶	>99.40	2.27 <u>+</u> 0.19	99.65
	30 min	§	>99.57	2.02 <u>+</u> 0.10¶	>99.81
	60 min	§	>99.57	§	>99.84

*Tests were performed in triplicate.NA, not applicable; TMPRSS2, transmembrane serine protease 2.

†The samples were incubated at room temperature (21-22°C).

‡TMPRSS2-expressing Vero-E6 cells were used for titration of viable viruses.

 $All the triplicates were below detection limit of the TCID_{50} assay.$

¶One or 2 out of 3 replicates were below detection limit of the TCID₅₀ assay.