

# Naturally Acquired Rabies in White-Eared Opossum, Brazil

Eduardo Ferreira-Machado,<sup>1</sup> Juliana Amorim Conselheiro,<sup>1</sup> Bruno Emerson Bernardes da Silva,<sup>1</sup> Patricia Sayuri Silvestre Matsumoto,<sup>1</sup> Cláudio Luiz Castagna,<sup>2</sup> Aline Nitsche,<sup>2</sup> Celino Simão de Lima,<sup>2</sup> Douglas Presotto,<sup>2</sup> Madelline Christie Rodrigues Nunes da Silva,<sup>2</sup> Ticiana Brasil Ervedosa,<sup>2</sup> Pedro Enrique Navas-Suárez,<sup>2</sup> Ísis Paixão de Jesus,<sup>2</sup> Julia de Carvalho,<sup>2</sup> Rodrigo Albergaria Ressio,<sup>2</sup> Cinthya dos Santos Cirqueira,<sup>2</sup> Gisely Toledo Barone,<sup>2</sup> Leila del Castillo Saad,<sup>2</sup> Paulo Eduardo Brandão,<sup>3</sup> José Luiz Catão-Dias,<sup>3</sup> Juliana Mariotti Guerra,<sup>3</sup> Natália Coelho Couto de Azevedo Fernandes<sup>3</sup>

Opossums are considered resistant to rabies. Nonhematophagous bats are reservoirs of rabies in urban areas of South America. We analyzed bats and opossums tested for rabies during 2021 in a highly urbanized city in Brazil to understand spillover in an urban setting. Wildlife surveillance is necessary to prevent rabies in humans and domestic animals.

Rabies is a viral zoonosis with high mortality rates caused by *Lyssavirus rabies* lineages (rabies virus, RABV) (1). Opossums of the genus *Didelphis* are marsupials widely distributed in the Americas, synanthropic in urban scenarios, and considered resistant to RABV (2). The main urban reservoirs of RABV in Brazil are nonhematophagous bats with distinct lineages and epidemiologic aspects (3). In 2021, passive surveillance programs detected an unusual case of rabies in a white-eared opossum (*D. albiventris*) by a RABV lineage of frugivorous bats of genus *Artibeus* spp. in Campinas, São Paulo state, Brazil, the 10th most urbanized city in the country (4). To elucidate the dynamics of this spillover, we describe the results of passive surveillance for rabies in bats and opossums in Campinas in 2021.

## The Study

In 2021, we tested samples of frozen brain tissue from 930 bats and 22 opossums for rabies by direct

fluorescent antibody test and confirmed infection by virus isolation in cell culture (5) in Campinas. Fixed formalin brain tissue fragments in 15 of these 22 opossums were analyzed by histopathology. In addition, for the opossum that tested positive for rabies, we performed reverse transcription PCR and subsequent phylogenetic analysis of the glycoprotein gene of RABV in the frozen brain tissue and conducted immunohistochemical analysis for rabies in fixed formalin tissues (cerebrum, cerebellum, heart, lungs, liver, spleen, kidney, and adrenal glands) (Appendix 1, <https://wwwnc.cdc.gov/EID/article/29/12/23-0373-App1.pdf>; Appendix 2, <https://wwwnc.cdc.gov/EID/article/29/12/23-0373-App2.xlsx>). Ethics approval was granted by the Ethics Committee in the Use of Animals of the School of Veterinary Medicine and Animal Science, University of São Paulo (approval no. 8227140222), according to the Ethical Principles in Animal Research.

Of the 22 opossums tested for rabies, 1 (4.5%) adult female white-eared opossum (*D. albiventris*) had a positive result. Death was caused by traumatic lesions in 10 (45.4%) opossums; 4 (18.2%) of those deaths were caused by interspecies interactions with dogs. Of the 15 opossums analyzed by histopathology, 14 (93.3%) were found in the urban zone, inside households in densely urbanized areas, or in residences on the

Author affiliations: Adolfo Lutz Institute, São Paulo, Brazil (E. Ferreira-Machado, T.B. Ervedosa, P.E. Navas-Suárez, I.P. de Jesus, J. de Carvalho, R.A. Ressio, C.S. Cirqueira, J.M. Guerra, N.C.C.A. Fernandes); University of São Paulo, São Paulo (E. Ferreira-Machado, J.L. Catão-Dias, L.C. Saad, P.E. Brandão, J.M. Guerra); Laboratory of Zoonoses and Vector-borne Diseases, Zoonoses Surveillance Division, Health Surveillance Coordination, São Paulo (J.A. Conselheiro, G.T. Barone); Surveillance Unit in Zoonoses, Campinas, Brazil

(B.E.B. Silva, C.L. Castagna, A. Nitsche, C.S. Lima); Saint Mary's University, Halifax, Nova Scotia, Canada (P.S.S. Matsumoto); Bosque dos Jequitibás Zoo Department of Parks and Gardens, Municipal Secretary of Public Services, Campinas (D. Presotto, M.C.R.N. Silva)

DOI: <https://doi.org/10.3201/eid2912.230373>

<sup>1</sup>These first authors contributed equally to this article.


<sup>2</sup>These senior authors contributed equally to this article.

<sup>3</sup>These authors were co-principal investigators.


outskirts of the city; death was caused by traumatic lesions in 10 (66.7%) opossums, 4 of those deaths were caused by interspecies interactions with dogs. On histopathologic examination, we observed no lesions in 8 opossums, hemodynamic lesions in 4, autolysis in 2, and mononuclear meningoencephalitis in the rabies-positive opossum (Appendix 1 Figure 1). In the rabies-positive opossum, RABV antigen was detected by immunohistochemistry in the cerebrum (Appendix 1 Figure 2), cerebellum, adrenal gland, liver, and heart. The RABV-positive opossum was found in a zoo located within a park in the urban center of Campinas and demonstrated signs of the paralytic form of RABV infection. Phylogenetic reconstruction demonstrated

that the RABV clustered within the frugivorous fruit-eating bats (*Artibeus* spp.) lineage circulating in Brazil (GenBank accession no. ON604858) (Figure 1).

During 2021, the frequency of rabies detected in bats was 3.2% (30/930). Among the rabies-positive bats, 17 (56.7%) were frugivorous species of fruit-eating bats (*Artibeus* spp.); the other 13 (43.4%) were insectivorous bats of *Eptesicus* spp., *Myotis* spp., and *Tadarida* spp. (Appendix 2 Table 3). In total, bats from those 4 genera represented 153 (16.4%) of the total bats investigated. Bats were tested periodically, and different seasonality peaks were noted in frugivorous and insectivorous bats (Appendix 1 Figures 5, 6). Rabies-positive bats were found in the urban perimeter of the



**Figure 1.** Rabies virus G gene phylogenetic tree showing specific clusters for different genera of bats in Brazil and dog-related samples in study of naturally acquired rabies in a white-eared opossum, Brazil (red text). The phylogeny was reconstructed by maximum-likelihood estimation from nucleotide sequences. Bootstrap values of >50% are depicted (1,000 bootstrap replicates). CVS corresponds to a fixed strain of the rabies virus. European bat lyssavirus-1 was used as an outgroup. The tree was visualized using iTOL version 6 (6). GenBank accession numbers are provided for reference sequences.



**Figure 2.** Kernel density map of concentration of *Artibeus* spp. bats in study of naturally acquired rabies in a white-eared opossum, Brazil. The kernel concentration layer of *Artibeus* is overlapped by layers of high concentration of *Eptesicus*, *Myotis*, and *Tadarida* spp. The opossum was found in a vegetated area with a high concentration of *Artibeus* spp. bats.

municipality of Campinas; 73.3% were found in areas of sparse vegetation and 26.7% in areas of remnants of vegetation (Appendix 1 Table). We identified bats in a regular spatial distribution throughout the city; we observed a small area of concentration in the north and a slight concentration of rabies-positive bats in the center of the city (Appendix 1 Figure 3). According to genus classification, *Artibeus* spp. bats were found in medium and high concentrations and overlapped spatially with a high concentration of insectivorous bats. Of note, opossums were found near areas of medium to high bat concentrations, and the rabies-positive opossum was captured in a vegetated area with a high concentration of *Artibeus* spp. bats (Figure 2). We also found a spatial diffusion of *Artibeus* spp. bats that overlapped with the rabies-positive opossum (Appendix 1 Figure 4), demonstrating a time overlap in August 2021.

### Conclusions

Experimental virus inoculation in the 1960s led to initial suggestions of resistance to infection by RABV

in *Didelphis* spp. opossums. (7). Reports of RABV in opossums are scarce; their low body temperature (34.4–36.1°C [94–97°F]) and the minimal possibility of surviving an attack by a rabid animal have been suggested as probable causes of the low prevalence of this disease in opossums in North America, where wild carnivorous mammals are natural reservoirs (2). Despite the low reports of rabies in opossums, a seroprevalence study conducted in São Paulo state observed a prevalence of RABV of 1.6% (5/312) in *Didelphis* spp., indicating contact between this animal population and RABV (8). Neurologic signs demonstrated by the rabies-positive opossum in this study are associated with paralytic form rabies, a common form transmitted by bats (9), and detection of viral particles in other organs indicates a phase of systemic spread. Interspecies interactions with bats in urban centers could be hypothesized as a route of RABV to the opossum, as has been observed in recent episodes of RABV in cats in Campinas (10,11). Unlike the scenario described in North America (2), opossums might survive interactions with

bats. Opossum deaths detected in this study occurred in anthropic areas of the city; they were more prevalent in homes and were caused by traumatic events, such as attacks by dogs, warning of the possible risk for infection with RABV in domestic animals.

Frugivorous and insectivorous bats are reservoirs of RABV in urban centers of South America; bat lineages are replacing RABV canid lineage after successful vaccination efforts were adopted in Brazil in the dog population (11,12). The spatial distribution of captured bats and opossums revealed an overlap in habitats between rabies-positive bats and opossums in urban areas. The rabies-positive opossum was found in a vegetated area within a very urbanized area densely occupied by *Artibeus* spp. bats; those areas of dense bat population might create conditions in which rabies transmission, and development of new hosts and strains, is more tied to ecologic factors than to the phylogenetic characteristics of the hosts (13). In São Paulo state, vaccination campaigns for dogs and cats were discontinued after dog RABV lineages had not been detected for >20 years. Spillover cases such as those described in this study indicate the importance of wildlife mammal surveillance to detect RABV, particularly in urban areas, where those animals can assume the role of host and act as a source of infection for humans. Spatial analysis can be a powerful tool to assist in rabies surveillance. Although some studies have conducted mapping of bat populations in cities in Brazil (11,14), such studies are scarce and need structured surveillance programs with trained teams. In addition, we noted seasonality in the RABV bat genus and in rabies-positive bats; insectivorous bats were commonly positive in summer and spring and the frugivorous genus *Artibeus* bats were more commonly positive in fall and winter, as described by Dias et al. (11).

This case shows that opossums are susceptible to rabies and can potentially acquire RABV from bats, as was suggested by the ecospatial analysis. Elucidating this possibility—through the detection of the dead opossum—occurred through integrated surveillance involving motivated field and laboratory teams. Our findings highlight the need for continuous surveillance of wildlife to clarify the dynamics of zoonotic diseases and to prevent their occurrence in humans and domestic animals, in agreement with a One Health approach.

#### Acknowledgments

We thank all members of the Surveillance Unit in Zoonoses, Bosque dos Jequitibás Zoo, Laboratory of Zoonoses and Vector-borne Diseases, Biological Sample Management Center, and Pathology Center of Adolfo Lutz Institute.

This study was supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) process number 404510/2021; Grupo de Apoio às Políticas de Prevenção e Proteção à Saúde/Fundo Especial de Saúde para Imunização em Massa e controle de Doenças (GAPS/FESIMA grant nos. 040/2019 and 28/2020); and The São Paulo Research Foundation—FAPESP (processes nos. 2022/02242-0 and 2023/04780-2). J.L.C.-D. is the recipient of a professorship from CNP (304106/2022-4).

#### About the Author

Mr. Ferreira-Machado is a PhD candidate in the school of Veterinary Medicine and Animal Sciences, University of São Paulo, São Paulo, Brazil. His primary research interests are wildlife comparative pathology and wildlife infectious diseases.

#### References

1. Scott TP, Nel LH. Lyssaviruses and the fatal encephalitic disease rabies. *Front Immunol.* 2021;12:786953. <https://doi.org/10.3389/fimmu.2021.786953>
2. Diana NK, Mitchell KC, Feldman KA. Shedding light on rabies in opossums. *J Am Vet Med Assoc.* 2015;247:1229.
3. Castilho JG, de Souza DN, Oliveira RN, Carnieli P Jr, Batista HBCR, Pereira PMC, et al. The epidemiological importance of bats in the transmission of rabies to dogs and cats in the state of São Paulo, Brazil, between 2005 and 2014. *Zoonoses Public Health.* 2017;64:423–30. <https://doi.org/10.1111/zph.12320>
4. Brazilian Institute of Geography and Statistics (IBGE). Cities and states: Campinas Panorama [cited 2023 Jul 2]. <https://cidades.ibge.gov.br/brasil/sp/campinas/panorama>
5. World Organization of Animal Health (WOAH). Chapter 3.1.18. Rabies (infection with rabies virus and other lyssaviruses). In: WOAH Terrestrial Manual 2023 [cited 2023 Mar 2]. [https://www.woah.org/fileadmin/Home/eng/Health\\_standards/tahm/3.01.18\\_RABIES.pdf](https://www.woah.org/fileadmin/Home/eng/Health_standards/tahm/3.01.18_RABIES.pdf)
6. Letunic I, Bork P. Interactive Tree of Life (iTOL): an online tool for phylogenetic tree display and annotation. *Nucleic Acids Res.* 2021;49:W293–6. PMID 33885785
7. Beamer PD, Mohr CO, Barr TR. Resistance of the opossum to rabies virus. *Am J Vet Res.* 1960;21:507–10.
8. Fornazari F, Scheffer KC, Dos Ramos Silva S, da Silva KR, Rodrigues AC, Teixeira CR, et al. Seroprevalence to rabies virus in wildlife in Brazil. *J Wildl Dis.* 2022;58:431–5. <https://doi.org/10.7589/JWD-D-21-00065>
9. Udow SJ, Marrie RA, Jackson AC. Clinical features of dog- and bat-acquired rabies in humans. *Clin Infect Dis.* 2013;57:689–96. <https://doi.org/10.1093/cid/cit372>
10. Gazarini J, Brito JE, Bernardi IP. Opportunistic predations of bats by *Didelphis albiventris* in southern Brazil [in Portuguese]. *Chiroptera Neotropical.* 2008;14:408–11.
11. Dias RA, Rocha F, Ulloa-Stanojlovic FM, Nitsche A, Castagna C, de Lucca T, et al. Spatiotemporal distribution of a non-haematophagous bat community and rabies virus circulation: a proposal for urban rabies surveillance in Brazil. *Epidemiol Infect.* 2019;147:e130. <https://doi.org/10.1017/S0950268818003229>
12. Queiroz LH, Favoretto SR, Cunha EM, Campos AC, Lopes MC, de Carvalho C, et al. Rabies in southeast

- Brazil: a change in the epidemiological pattern. *Arch Virol.* 2012;157:93–105. <https://doi.org/10.1007/s00705-011-1146-1>
13. Caraballo DA, Lema C, Novaro L, Gury-Dohmen F, Russo S, Beltrán FJ, et al. A novel terrestrial rabies virus lineage occurring in South America: origin, diversification, and evidence of contact between wild and domestic cycles. *Viruses.* 2021;13:2484. <https://doi.org/10.3390/v13122484>
14. Ribeiro J, Staudacher C, Martins CM, Ullmann LS, Ferreira F, Araujo JP Jr, et al. Bat rabies surveillance and


risk factors for rabies spillover in an urban area of Southern Brazil. *BMC Vet Res.* 2018;14:173. <https://doi.org/10.1186/s12917-018-1485-1>

Address for correspondence: Eduardo Ferreira-Machado, 87 Av. Prof. Dr. Orlando Marques de Paiva, Butantã, São Paulo, 05508-270, São Paulo, Brazil; email: eduardo.ferreira.machado@usp.br

February 2023

# Emerging Pathogens

- Infant Botulism, Israel, 2007–2021
- Sentinel Surveillance System Implementation and Evaluation for SARS-CoV-2 Genomic Data, Washington, USA, 2020–2021
- Crimean-Congo Hemorrhagic Fever, Spain, 2013–2021
- *Streptococcus dysgalactiae* Bloodstream Infections, Norway, 1999–2021
- Changing Disease Course of Crimean-Congo Hemorrhagic Fever in Children, Turkey
- Relationship between Telework Experience and Presenteeism during COVID-19 Pandemic, United States, March–November 2020
- Circovirus Hepatitis Infection in Heart-Lung Transplant Patient, France
- Incidence and Transmission Dynamics of *Bordetella pertussis* Infection in Rural and Urban Communities, South Africa, 2016–2018
- Influence of Landscape Patterns on Exposure to Lassa Fever Virus, Guinea
- Increased Multidrug-Resistant *Salmonella enterica* I Serotype 4,[5],12:i:- Infections Associated with Pork, United States, 2009–2018
- Novel Prion Strain as Cause of Chronic Wasting Disease in a Moose, Finland
- Novel Species of *Brucella* Causing Human Brucellosis, French Guiana
- Penicillin and Cefotaxime Resistance of Quinolone-Resistant *Neisseria meningitidis* Clonal Complex 4821, Shanghai, China, 1965–2020
- Combined Phylogeographic Analyses and Epidemiologic Contact Tracing to



Characterize Atypically Pathogenic Avian Influenza (H3N1) Epidemic, Belgium, 2019

- Age-Stratified Model to Assess Health Outcomes of COVID-19 Vaccination Strategies, Ghana
- Early Introduction and Community Transmission of SARS-CoV-2 Omicron Variant, New York, New York, USA
- Correlates of Protection, Thresholds of Protection, and Immunobridging among Persons with SARS-CoV-2 Infection
- Longitudinal Analysis of Electronic Health Information to Identify Possible COVID-19 Sequelae
- Nipah Virus Exposure in Domestic and Peridomestic Animals Living in Human Outbreak Sites, Bangladesh, 2013–2015
- (Mis)perception and Use of Unsterile Water in Home Medical Devices, PN View 360+ Survey, United States, August 2021

- Molecular Detection of *Candidatus Orientia chuto* in Wildlife, Saudi Arabia
- Neorickettsiosis in Symptomatic Immunocompetent Child, South Africa
- Successful Drug-Mediated Host Clearance of *Batrachochytrium salamandrivorans*
- Powassan Virus Lineage I in Field-Collected *Dermacentor variabilis* Ticks, New York, USA
- *Bartonella* spp. and Typhus Group Rickettsiae among Persons Experiencing Homelessness, São Paulo, Brazil
- *Candida auris* Discovery through Community Wastewater Surveillance during Healthcare Outbreak, Nevada, USA, 2022
- Estimated Cases Averted by COVID-19 Digital Exposure Notification, Pennsylvania, USA, November 8, 2020–January 2, 2021
- Next-Generation Sequencing for Identifying Unknown Pathogens in Sentinel Immunocompromised Hosts
- Orthopoxvirus Infections in Rodents, Nigeria, 2018–2019
- Occupational Monkeypox Virus Transmission to Healthcare Worker, California, USA, 2022
- Familial Monkeypox Virus Infection Involving 2 Young Children
- Relapsing Fever Caused by *Borrelia lonestari* after Tick Bite in Alabama, USA
- *Nocardia neocaledoniensis* as Rare Cause of Spondylodiscitis
- *Cryptococcus tetragattii* Meningitis Associated with Travel, Taiwan

**EMERGING  
INFECTIOUS DISEASES**

To revisit the February 2023 issue, go to:  
<https://wwwnc.cdc.gov/eid/articles/issue/29/2/table-of-contents>

*EID cannot ensure accessibility for supplementary materials supplied by authors. Readers who have difficulty accessing supplementary content should contact the authors for assistance.*

# Naturally Acquired Rabies in White-Eared Opossum, Brazil

## Appendix 1

### Material and Methods

**Material and Local of Analysis:** From January to December 2021, frozen brain tissue samples of 1,049 bats and 22 opossums were investigated for rabies in Campinas by Direct Fluorescent Antibody Test (dFAT) and confirmatory by Virus Isolation in Cell Culture (VICC). Data of the animal, e.g., taxonomic classification, sex, age class, and the address in which they were found, were recorded. Of the 1,049 bats, 119 individuals were discarded from analyses for inability to rabies test by autolysis or not determined by inconsistency in the diagnostic sheet. In addition, tissue fragments fixed in formalin buffered 10% of 15 opossums were collected and the necropsy exams were compiled by data from necropsy, causa mortis, gross lesions, and the place where the animals were found... The formalin fixed paraffin embedded samples were analyzed by histopathology. The positive opossum was tested additionally to the immunohistochemistry (IHC) test for rabies, RT-PCR and sequencing of the glycoprotein gene .

**dFAT and VICC:** dFAT was conducted with impression smears made from sections of the brain that were fixed with cold acetone ('30'), stained with the anti-rabies antibody conjugate (in-house Pasteur Institute) in a humid chamber ('30' at 37°C) and buffered with PBS ('10'). The slides were then examined under a fluorescence microscope (Nikon EFD-3) with a 40x objective in search of fluorescent antigen-antibody aggregates. The VICC was performed with frozen samples of a 20% suspension (mass/volume) of brain tissue, and PBS was obtained by grinding with mortar and pestle and centrifuged at 1500 x g (30' at 4°C). The clarified homogenate was

then inoculated in murine neuroblastoma cells (4x10<sup>5</sup> cells/mL) in triplicate and the microplate was incubated at 37°C in a 5% CO<sub>2</sub> humidified incubator for 96 hours. After the incubation period, the microplate was fixed and stained as described above and observed in a fluorescence microscope (Zeiss Axio Vert.A1) with a 20x objective in search of fluorescent infected cells.

**Histopathology:** Fixed samples were embedded in paraffin wax, sectioned at 4 µm-thick, and stained with hematoxylin and eosin for histologic analysis. For the CNS analysis, no degenerative changes, e.g., vacuolar degeneration and death neurons were considered due to the lack of knowledge between the time of death and the *postmortem* exams and collected performance.

**IHC:** Deparaffinized 3µm sections of tissues in silanized slides were submitted to enzymatic digestion (20' at 37°C) by proteinase K (125mg/ml). Endogenous peroxidase was blocked with 6% hydrogen peroxide (30') followed by overnight incubation with mouse hyperimmune antiserum polyclonal (in-house - Evandro Chagas Institute, Pará, Brazil) at a concentration of 1/2000. The signal was amplified by Polink-2AP Broad Kit (GBI Labs; WA; USA) for '60' conjugated to alkaline phosphatase and visualization was achieved by warp red(3') (Polink-2 AP Broad Detection System; GBI Labs, WA, USA, cat. D24-110). The samples were counterstained with Harris Hematoxylin (20''), followed by dehydration and slide mounting with synthetic resin. Brain mammal tissue fragments known to be positive and confirmed by IHC were used as positive controls. The same steps were followed for the negative control, except for the primary antibody incubation, replaced by non-immune serum from a mouse.

**RT-PCR and phylogeny:** Brain samples were submitted to RNA extraction with TRIzol (Thermo Fisher Scientific) following the manufacturer's instructions. Complementary DNA synthesis was made using the Superscript VILO cDNA synthesis kit (Thermo Fisher Scientific) following the manufacturer's protocol. PCR was performed using Platinum Taq DNA Polymerase (Thermo Fisher Scientific), and the primers Ga3222-40 (5' CGCTGCATTTTRTCARAGT 3') and Gb4119-39 (5' GGAGGGCACCATTGGMTC 3') targeting the glycoprotein gene as described elsewhere (1). Gel electrophoresis was used to confirm specific product amplification. The amplicon was purified by an enzymatic method (ExoSAP-IT – Thermo Fisher Scientific), and the BigDye Terminator v3.1 Cycle Sequencing kit (Thermo Fisher Scientific) was used for sequencing reactions. Sequences were obtained in ABI-

3500. PHRED algorithm was used for base quality analysis of the sequence generated. Values equal to or higher than 20 were considered real. Next, the software BioEdit Sequence Alignment Editor v. 7.0.3 (2) was used for generating the consensus sequence and multiple alignments. A database was built with sequences retrieved from GenBank representative of rabies virus lineages circulating in Brazil, and the phylogenetic reconstruction was inferred by Maximum Likelihood using PAUP 4.0 (3) with GTR+I+G as an evolutionary model with 1,000 bootstrap replicates. European Bat Lyssavirus was used as an outgroup.

**Geospatial Analysis:** Bats and opossums locations were georeferenced using an API in Google Sheets Geocoding service (4). The geographic coordinates were imported into ArcGIS Pro 2.9.5, of which 930 (88.6% of bats) and 18 (81.8% of opossums) locations of animal captures were successfully located. After validation, 94 bats (8.96%) and four opossums (18.18%) had to be manually geo-edited to ensure the correct locations. Each animal represented a point (shapefile) and was categorized according to its taxonomy and outcome diagnosis in the thematic maps. Spatial analysis was performed using Kernel estimator density (5). We defined a bandwidth of 1,000m to identify the density of bats by genus and overlap its information with the Opossum's location. In addition, areas of influence of 50m (buffers) were created around the location of the capture of positive animals to identify the presence of vegetation, classified as densely vegetated, remnants of vegetation, sparsely vegetated, and very sparsely vegetated, ranging from higher to lower arboreal vegetation cover, respectively (see Appendix 1 Table).

## References

1. Sato G, Itou T, Shoji Y, Miura Y, Mikami T, Ito M, et al. Genetic and phylogenetic analysis of glycoprotein of rabies virus isolated from several species in Brazil. *J Vet Med Sci.* 2004;66:747–53. [PubMed https://doi.org/10.1292/jvms.66.747](https://doi.org/10.1292/jvms.66.747)
2. Hall TA. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucl. Acids Symp. Ser.* 1999;41:95–8 [cited 2023 Mar 2]. [https://www.academia.edu/2034992/BioEdit\\_a\\_user\\_friendly\\_biological\\_sequence\\_alignment\\_editor\\_and\\_analysis\\_program\\_for\\_Windows\\_95\\_98\\_NT](https://www.academia.edu/2034992/BioEdit_a_user_friendly_biological_sequence_alignment_editor_and_analysis_program_for_Windows_95_98_NT)
3. Swofford D, Sullivan J. Phylogeny inference based on parsimony and other methods using PAUP. In: Lemey P, Salemi M, Vandamme A, eds. *The phylogenetic handbook: a practical approach to*







phylogenetic analysis and hypothesis testing. Cambridge: Cambridge University Press; 2009. p. 267–312.

4. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new developments. *Nucleic Acids Res.* 2019;47(W1):W256–9. [PubMed https://doi.org/10.1093/nar/gkz239](https://doi.org/10.1093/nar/gkz239)


5. Bailey TC, Gatrell AC. *Interactive spatial data analysis.* New York: Longman; 1995.

**Appendix 1 Table.** Classification of the land use and land cover where the rabies-positive animals were investigated


Interpretation keys	Buffer	Bats	Opossums
	Densely vegetated	–	1 positive
	Remnants of vegetation	8 (6 <i>Artibeus</i> , 2 <i>Myotis</i> )	–

Interpretation keys	Buffer	Bats	Opossums
	Sparsely vegetated	15 (10 <i>Artibeus</i> , 3 <i>Epitesicus</i> , 2 <i>Myotis</i> )	–
			
	Very sparsely vegetated	7 (2 <i>Tadarida</i> , 1 <i>Myotis</i> , 3 <i>Epitesicus</i> , 1 <i>Artibeus</i> )	–


Yellow buffer of 50m with 90% transparency around the animal capture location. ArcGIS Pro 2.9.5. Imagery basemaps (Source: Maxar 2021/07/24; 0.46m spatial resolution). Cartographic scale 1:1,000.




**Appendix 1 Figure 1.** CNS histology of rabies in opossum. HE, 200x, Cerebellum. Mononuclear perivascular cuffing.




**Appendix 1 Figure 2.** IHC of rabies in the positive opossum. IHQ, 400x, Brain. Neuronal cell with cytoplasmic antigen detected with AP for LR (arrow).




**Appendix 1 Figure 3.** Spatial distribution of bats classified by Family and positive diagnostic. Each gray dot represents one collection of bats: Molossidae are represented by circles, Phyllostomidae by squares, and Vespertilionidae by triangles. The positive ones are colored in red.



**Appendix 1 Figure 4.** Diffusion of *Artibeus* captured from January to December 2021. The dates were interpolated from February to December, varying from red to blue. The location where the positive opossum was found corresponds to August; however, considering the home range of Bats ( $\approx 19\text{ha}$ ), in addition to August, the diffusion shows also July and September 2021.



**Appendix 1 Figure 5.** Temporal distribution of total genus RABV detected in insectivorous and phytophagous bats.



**Appendix 1 Figure 6.** Temporal distribution of total bats RABV detected in insectivorous and phytophagous bats.