## About the Author

Dr. Guimarães is a research scientist and analyst at the Ezequiel Dias Foundation. Her research interests and work include the molecular diagnosis and sequencing of SARS-CoV-2, monkeypox virus, and arboviruses, including dengue, Zika, chikungunya, and yellow fever viruses, and other pathogens of importance to public health.

#### References

- McCollum AM, Damon IK. Human monkeypox. Clin Infect Dis. 2014;58:260–7. https://doi.org/10.1093/cid/ cit703
- Essbauer S, Pfeffer M, Meyer H. Zoonotic poxviruses. Vet Microbiol. 2010;140:229–36. https://doi.org/10.1016/ j.vetmic.2009.08.026
- Mauldin MR, McCollum AM, Nakazawa YJ, Mandra A, Whitehouse ER, Davidson W, et al. Exportation of monkeypox virus from the African continent. J Infect Dis. 2022;225:1367–76. https://doi.org/10.1093/infdis/jiaa559
- Minhaj FS, Ogale YP, Whitehill F, Schultz J, Foote M, Davidson W, et al.; Monkeypox Response Team 2022. Monkeypox outbreak – nine states, May 2022. MMWR Morb Mortal Wkly Rep. 2022;71:764–9. https://doi.org/10.15585/ mmwr.mm7123e1
- Pan American Health Organization (PAHO). Mpox [cited 2023 Mar 10]. https://www.paho.org/en/monkeypox
- World Health Organization (WHO). Global genomic surveillance strategy for pathogens with pandemic and epidemic potential, 2022–2032 [cited 2023 Mar 2]. https://www.who.int/publications/i/item/9789240046979
- Isidro J, Borges V, Pinto M, Sobral D, Santos JD, Nunes A, et al. Phylogenomic characterization and signs of microevolution in the 2022 multi-country outbreak of monkeypox virus. Nat Med. 2022;28:1569–72. https://doi.org/10.1038/s41591-022-01907-y
- Roychoudhury P, Sereewit J, Xie H, Nunley E, Bakhash SM, Lieberman NAP, et al. Genomic analysis of early monkeypox virus outbreak strains, Washington, USA. Emerg Infect Dis. 2023;29:644–6. https://doi.org/10.3201/eid2903.221446
- 9. Ministry of Health. Notifiable Diseases Information System–SINAN [in Portuguese] [cited 2023 Mar 10]. http://portalsinan.saude.gov.br
- Menezes YR, Miranda AB. Severe disseminated clinical presentation of monkeypox virus infection in an immunosuppressed patient: first death report in Brazil. Rev Soc Bras Med Trop. 2022;55:e0392-2022. https://doi.org/10.1590/0037-8682-0392-2022

Address for correspondence: Felipe Campos de Melo Iani, Ezequiel Dias Foundation, Rua Conde Pereira Carneiro, 80— Gameleira, Belo Horizonte 30510-010, Brazil; email: felipe.iani@funed.mg.gov.br

## Antimicrobial-Resistant Infections after Turkey/Syria Earthquakes, 2023

## Anthony Rizk, Antoine Abou Fayad, Louis-Patrick Haraoui

Author affiliations: Geneva Graduate Institute, Geneva, Switzerland (A. Rizk); American University of Beirut, Beirut, Lebanon (A. Abou Fayad); Université de Sherbrooke, Sherbrooke, Québec, Canada (L.-P. Haraoui)

DOI: https://doi.org/10.3201/eid2906.230316

Increased rates of multidrug-resistant microbes have been reported after earthquakes. After the 2023 earthquakes in Turkey and Syria, the number of associated highly drug-resistant pathogens and nosocomial transmission will probably surge in hospitals treating injured patients. It is not too late to act to prevent antimicrobialresistant infections from compounding these tragedies.

he 2023 earthquakes that affected Turkey and Syria, with Kahramanmaraş Province in Turkey at their epicenter, measured 7.8 and 7.5 on the Richter scale. The effects were devastating, making these the strongest earthquakes in Turkey since 1939. Combined with their multiple aftershocks, the earthquakes caused >50,000 deaths and severely damaged or collapsed >170,000 buildings (https:// www.aljazeera.com/news/2023/2/25/deathtoll-climbs-above-50000-after-turkey-syria-earthquakes). In their wake, the earthquakes left a growing humanitarian crisis. If previous experiences are any indication, we can also expect hospitals caring for the injured and wounded to struggle with highly antimicrobial-resistant infections, many of which will lead to excess illnesses and deaths.

Multidrug resistant microbes have often been reported after earthquakes and other natural disasters. Medical literature on earthquake-associated injuries, going as far back as the Marmara, Turkey, earthquake of 1999 (1), have consistently shown highly resistant microbial strains emerging in hospital settings and causing hospital-acquired infections in trauma patients. Antimicrobial-resistant *Acinetobacter baumannii* has been identified in disproportionately high rates from infections associated with large-scale earthquakes in Southeast Asia in 2004; northern Pakistan in 2005; Wenchuan, China, in 2008; central Italy in 2009; and Haiti in 2010 (Appendix, https://wwwnc.cdc.gov/EID/article/29/6/23-0316-App1.pdf).

Although earthquake-associated pathogens detected in hospital settings have been consistently multidrug resistant, their resistance profiles have varied. The etiology of antimicrobial resistance from earthquakes remains uncertain. After the Marmara earthquake of 1999, associated A. baumannii and Pseudomonas aeruginosa infections were mainly resistant to carbapenems, which had been preemptively administered in large numbers to patients before hospitalization (1). After the 2004 Southeast Asia earthquake, patients with multiple wounds admitted to the intensive care unit of the Cologne-Merheim Medical Center in Germany were reportedly contaminated with multidrug-resistant A. baumannii, extended-spectrum β-lactamase *Escherichia coli*, and methicillin-resistant Staphylococcus aureus (2). After the 2005 northern Pakistan earthquake, Kiani et al. reported emergence of multiply drug-resistant Acinetobacter spp., Enterobacter spp., and other gram-negative organisms, susceptible to amikacin only. Hospitals were overwhelmed with staff and antimicrobial drug shortages, and wounds were often treated empirically because of limited laboratory capacities (3). After the 2008 earthquake in Wenchuan, China, the predominant pathogen causing infections was A. baumannii; 1 hospital reported that >65% of Acinetobacter isolates were resistant to a wide range of antimicrobial drugs, except imipenem, and 24.6% of isolates were pan-drug resistant (4).

Since the 2023 Turkey/Syria earthquakes, multidrug resistance has not yet been reported. However, on the basis of the epidemiology of infections in earthquake-stricken regions, we expect hospitals in the region to struggle with containing nosocomial transmission of highly resistant pathogens that may increase deaths among hospitalized patients. Also of concern is the re-emergence of other infectious diseases, such as cholera, known to be precipitated by disaster settings (5). The Turkey/Syria earthquakes come on the heels of multiple crises in the region, many of which have been shown to influence the rising rates of multidrug resistance, as in the wake of the Iraq and Syrian conflicts (6). Rising rates of antimicrobial resistance are further compounded by the regionwide travel of wounded patients for treatment as healthcare infrastructure in Iraq and Syria collapsed (7). Medicine shortages, especially resulting from sanctions in Syria and Iraq and following the 2019 financial collapse in Lebanon, further compound selective pressures on microbes to develop specific resistance profiles.

Hospitals in and outside the region treating injuries from the Turkey/Syria earthquakes will probably observe a surge in highly resistant pathogens and nosocomial transmission. In addition to providing immediate care for the injured and displaced, relief efforts should therefore anticipate a probable increase in antimicrobial-resistant infections for which therapeutic options may be limited. Detecting and responding to this superimposed crisis urgently requires support and expansion of existing microbiology laboratory capacity. Recent initiatives, such as the deployment of telemicrobiology in northern Syria, are examples of how to rapidly assist local laboratory teams in settings of social disruption (8). Training healthcare workers on surgical techniques and wound care adapted to deal with traumatic injuries have also been shown to enhance wound healing and to limit complications such as chronic and recurrent skin and soft tissue infections, which are risk factors for antimicrobial resistance (9). Early-stage detection and aggressive infection-control practices (e.g., active surveillance, contact isolation, sampling of healthcare workers and hospital environments, and antimicrobial stewardship) during and after disasters play key roles in preventing resistant strains from becoming endemic to healthcare facilities (10). Healthcare facilities may need to consider patient decolonization through chlorhexidine bathing to forestall colonization by antimicrobial-resistant Acinetobacter strains (10). Communities affected by the recent earthquakes will probably experience their effects for months to come. It is not too late to act to prevent further complications from these natural disasters, such as antimicrobial-resistant infections, from compounding ongoing human tragedies.

This work was supported by the New Frontiers in Research Fund, Canada (NFRFE-2019-00444) (A.R. and L.-P.H.), the Fonds de Recherche du Quebec-Sante (282182) (L.-P.H.), and the Canadian Institute for Advanced Research (GS-0000000256) (L.-P.H.)

## About the Author

Mr. Rizk is a PhD candidate in anthropology at the Geneva Graduate Institute, Switzerland. His research interests include scientific laboratory capacities and antimicrobial resistance.

#### References

- 1. Oncül O, Keskin O, Acar HV, Küçükardali Y, Evrenkaya R, Atasoyu EM, et al. Hospital-acquired infections following the 1999 Marmara earthquake. J Hosp Infect. 2002;51:47–51. https://doi.org/10.1053/jhin.2002.1205
- Maegele M, Gregor S, Steinhausen E, Bouillon B, Heiss MM, Perbix W, et al. The long-distance tertiary air transfer and care of tsunami victims: injury pattern and microbiological and psychological aspects. Crit Care Med. 2005;33:1136–40. https://doi.org/10.1097/01.CCM.0000163269.42524.50
- 3. Kiani QH, Amir M, Ghazanfar MA, Iqbal M. Microbiology of wound infections among hospitalised patients following

the 2005 Pakistan earthquake. J Hosp Infect. 2009;73:71-8. https://doi.org/10.1016/j.jhin.2009.06.012

- Tao C, Kang M, Chen Z, Xie Y, Fan H, Qin L, et al. Microbiologic study of the pathogens isolated from wound culture among Wenchuan earthquake survivors. Diagn Microbiol Infect Dis. 2009;63:268–70. https://doi.org/ 10.1016/j.diagmicrobio.2008.11.009
- 5. Kelly-Hope LA. Conflict and emerging infectious diseases. Emerg Infect Dis. 2008;14:1004–5, author reply 1005.
- Haraoui LP, Valiquette L, Laupland KB. Antimicrobial resistance in conflicts. J Assoc Med Microbiol Infect Dis Can. 2018;3:119–22. https://doi.org/10.3138/jammi.2018.05.15
- Dewachi O, Skelton M, Nguyen VK, Fouad FM, Sitta GA, Maasri Z, et al. Changing therapeutic geographies of the Iraqi and Syrian wars. Lancet. 2014;383:449–57. https://doi.org/10.1016/S0140-6736(13)62299-0
- Karah N, Antypas K, Al-Toutanji A, Suveyd U, Rafei R, Haraoui LP, et al. Teleclinical microbiology: an innovative approach to providing web-enabled diagnostic laboratory services in Syria. Am J Clin Pathol. 2022;157:554–60. https://doi.org/10.1093/ajcp/aqab160
- Abu-Sittah GS, Hoballah JJ, Bakhach J, editors. Reconstructing the war injured patient. Cham, Switzerland: Springer International Publishing; 2017.
- Munoz-Price LS, Weinstein RA. Acinetobacter infection. N Engl J Med. 2008;358:1271–81. https://doi.org/10.1056/ NEJMra070741

Address for correspondence: Louis-Patrick Haraoui, Hôpital Charles-Le Moyne, Casier 251, 3120 Blvd Taschereau, Greenfield Park, QC J4V 2H1, Canada; email: louis.patrick.haraoui@usherbrooke.ca

## Detection of Severe Murine Typhus by Nanopore Targeted Sequencing, China

Panpan Qian, Xiaohua He, Mei Yang, Li Wei, Lihui Zhang, Xiqian Xing

Authors affiliation: The Affiliated Hospital of Yunnan University, Kunming, China

#### DOI: https://doi.org/10.3201/eid2906.221929

We report a case of murine typhus in China caused by *Rickettsia typhi* and diagnosed by nanopore targeted sequencing of a bronchoalveolar lavage fluid sample. This case highlights that nanopore targeted sequencing can effectively detect clinically unexplained infections and be especially useful for detecting infections in patients without typical signs and symptoms.

Murine typhus is caused by *Rickettsia typhi* bacteria transmitted by rat or cat flea vectors. Persons with murine typhus often have nonspecific or mild symptoms, such as fever, myalgia, and rash. In rare instances, murine typhus will cause atypical or multiple organ dysfunction syndrome (MODS) (1,2).

Murine typhus is an undifferentiated febrile illness, which makes it challenging to recognize and diagnose. We report a case of murine typhus and MODS in a patient without rash. We diagnosed murine typhus by using nanopore targeted sequencing (NTS) of a bronchoalveolar lavage fluid (BALF) sample, aiming to provide more reference for clinical practice.

A 60-year-old female farmer from Yunnan Province, China, had fatigue, anorexia, nausea, dizziness, and vomiting for 1 week. At admission, she was afebrile and hemodynamically stable and did not have headache, rash, or eschar. Chest computed tomography (CT) imaging showed pneumonia and a small plural effusion (Figure, panels A, B). By the next day, her condition had deteriorated. She experienced chills, fever (temperature 39°C), severe hypotension (70/53 mm Hg), dyspnea, and deterioration of the oxygenation index. Preliminary laboratory investigation demonstrated mild leukocytosis (13.86  $\times$  10<sup>9</sup> cells/L), moderately elevated transaminase levels (alanine aminotransferase 197 U/L, aspartate aminotransferase 128 U/L), severe thrombocytopenia ( $12 \times 10^9$  platelets/L), coagulation disorder (D-dimer 49.8 µg/mL), elevated C-reactive protein (207.4 mg/L) and procalcitonin (4.65 ng/mL) levels, and respiratory failure (partial pressure of oxygen 58.9 mm Hg).

The patient was given intravenous meropenem and norepinephrine and was admitted on noninvasive ventilation. We then conducted tests for malaria, *Legionella*, influenza virus, SARS-CoV-2, HIV, herpes simplex virus, cytomegalovirus, Epstein-Barr virus IgM, *Roxiella burnettii* IgM (phase II antigen), *R. typhi* IgM, *Mycoplasma pneumoniae* IgM, *Chlamydia* IgM, respiratory syncytial virus IgM, and adenovirus IgM; results were all negative. In addition, testing of blood, urine, stool, and sputum cultures and bone marrow biopsy all produced negative results.

On admission day 5, she remained normotensive. Her body temperature dropped, but she still had a low-grade fever, body temperature fluctuating from 37.5°C to 38°C. However, the cause of her severe infection remained unclear. The next day, she underwent bronchoscopy. BALF was sent to undergo NTS analysis to Wuhan Dgensee Clinical Laboratory Co., Ltd (https://www.dgensee.com). Two days later, NTS results revealed *R. typhi* DNA.

## Article DOI: https://doi.org/10.3201/eid29065.230316

EID cannot ensure accessibility for supplementary materials supplied by authors. Readers who have difficulty accessing supplementary content should contact the authors for assistance.

# Antimicrobial-Resistant Infections after Turkey/Syria Earthquakes, 2023

## Appendix

Table. Characteristics of earthquake-associated infections with antibiotic-resistant pathogens

Earthquake location,			
date	Severity	Casualties	Key contaminants of earthquake-associated wounds
Marmara, Turkey,	7.4 magnitude,	17,480 deaths,	Acinetobacter spp. and Pseudomonas aeruginosa resistant to
1999	affecting an area of	43,953 injured	carbapenems and sensitive to quinolones; methicillin-resistant
	$200 \text{ km} \times 40 \text{ km}$		Staphylococcus aureus (1)
Southeast Asia,	9.1 magnitude,	310,000 deaths,	MDR Acinetobacter spp.; ESBL Escherichia coli; methicillin-
2004	triggering a massive	millions destitute	resistant Staphylococcus aureus (MRSA); and Aeromonas
	tsunami		hydrophilia, Pseudomonas spp., and Candida albicans (2).
Northern Pakistan,	7.6 magnitude, with	>82,000 deaths, 3.3	MDR Pseudomonas spp., Enterobacter spp., and
2005	140 aftershocks	million injured	Acinetobacter spp. only susceptible to amikacin (3).
Wenchuan, China,	8.0 maginitude	>69,000 deaths,	MDR Acinetobacter baumannii, Escherichia coli,
2008		45.5 million destitute	Pseudomonas aeruginosa and Staphylococcus aureus (4,5).
Central Italy, 2009	5.9 magnitude	308 deaths, >1,000	Extensively drug-resistant Acinetobacter baumannii belonging
		injured	to A. baumannii sequence type 2 with bla <sub>OXA-23</sub> (6)
Haiti, 2010	7.0 magnitude	>100,000 deaths	Three A. baumannii isolates belonging to 2 distinct clones and
			were identified as ESBL producers and found to be <i>bla</i> <sub>CTX-M-15</sub> -
			positive. They were resistant to penicillins, broad-spectrum
			cephalosporins and aztreonam but susceptible to
			carbapenems (7).

## References

- Öncül O, Keskin O, Acar HV, Küçükardali Y, Evrenkaya R, Atasoyu EM, et al. Hospital-acquired infections following the 1999 Marmara earthquake. J Hosp Infect. 2002;51:47–51. <u>PubMed</u> https://doi.org/10.1053/jhin.2002.1205
- Maegele M, Gregor S, Steinhausen E, Bouillon B, Heiss MM, Perbix W, et al. The long-distance tertiary air transfer and care of tsunami victims: injury pattern and microbiological and psychological aspects. Crit Care Med. 2005;33:1136–40. <u>PubMed</u> <u>https://doi.org/10.1097/01.CCM.0000163269.42524.50</u>
- 3. Kiani QH, Amir M, Ghazanfar MA, Iqbal M. Microbiology of wound infections among hospitalised patients following the 2005 Pakistan earthquake. J Hosp Infect. 2009;73:71–8. <u>PubMed</u> <u>https://doi.org/10.1016/j.jhin.2009.06.012</u>

- Wang Y, Hao P, Lu B, Yu H, Huang W, Hou H, et al. Causes of infection after earthquake, China, 2008. Emerg Infect Dis. 2010;16:974–5. <u>PubMed https://doi.org/10.3201/eid1606.091523</u>
- Tao C, Kang M, Chen Z, Xie Y, Fan H, Qin L, et al. Microbiologic study of the pathogens isolated from wound culture among Wenchuan earthquake survivors. Diagn Microbiol Infect Dis. 2009;63:268–70. <u>PubMed https://doi.org/10.1016/j.diagmicrobio.2008.11.009</u>
- Perilli M, Sabatini A, Pontieri E, Celenza G, Segatore B, Bottoni C, et al. OXA-23 carbapenemase in multidrug-resistant *Acinetobacter baumannii* ST2 type: first identification in L'Aquila Hospital (Italy). Microb Drug Resist. 2015;21:97–101. <u>PubMed https://doi.org/10.1089/mdr.2014.0056</u>
- 7. Potron A, Munoz-Price LS, Nordmann P, Cleary T, Poirel L. Genetic features of CTX-M-15-producing Acinetobacter baumannii from Haiti. Antimicrob Agents Chemother. 2011;55:5946–8. <u>PubMed</u> <u>https://doi.org/10.1128/AAC.05124-11</u>