

8. Meyer Sauter PM, Beeton ML, Pereyre S, Bébéar C, Gardette M, Hénin N, et al.; ESGMAC the ESGMAC MAPS study group. *Mycoplasma pneumoniae*: gone forever? *Lancet Microbe*. 2023;4:e763. [https://doi.org/10.1016/S2666-5247\(23\)00182-9](https://doi.org/10.1016/S2666-5247(23)00182-9)
9. Meyer Sauter PM, Beeton ML, Pereyre S, Bébéar C, Gardette M, Hénin N, et al. European Society of Clinical Microbiology and Infectious Diseases (ESCMID) Study Group for Mycoplasma and Chlamydia Infections (ESGMAC), and the ESGMAC *Mycoplasma pneumoniae* Surveillance (MAPS) Study Group. *Mycoplasma pneumoniae*: delayed re-emergence after COVID-19 pandemic restrictions. *Lancet Microbe*. 2024;5:e100-1. [https://doi.org/10.1016/S2666-5247\(23\)00344-0](https://doi.org/10.1016/S2666-5247(23)00344-0)
10. Opota O, Brouillet R, Greub G, Jaton K. Methods for real-time PCR-based diagnosis of *Chlamydia pneumoniae*, *Chlamydia psittaci*, and *Chlamydia abortus* infections in an opened molecular diagnostic platform. *Methods Mol Biol*. 2017;1616:171-81. https://doi.org/10.1007/978-1-4939-7037-7_11

Address for correspondence: Gilbert Greub, Institute of Microbiology, Department of Laboratory Medicine and Pathology, Lausanne University Hospital and University of Lausanne, Bugnon 48, CH-1011 Lausanne, Switzerland; email: gilbert.greub@chuv.ch

Highly Pathogenic Avian Influenza A(H5N1) Viruses from Multispecies Outbreak, Argentina, August 2023

Agustina Rimondi, Ralph E.T. Vanstreels, Valeria Olivera, Agustina Donini, Martina Miqueo Lauriente, Marcela M. Uhart

Author affiliations: Robert Koch Institute, Berlin, Germany (A. Rimondi); Instituto Nacional de Tecnología Agropecuaria Instituto de Virología e Innovaciones Tecnológicas, Buenos Aires, Argentina (A. Rimondi, V. Olivera); University of California School of Veterinary Medicine, Davis, California, USA (R.E.T. Vanstreels, M.M. Uhart); Southern Right Whale Health Monitoring Program, Puerto Madryn, Argentina (A. Donini, M.M. Uhart); Secretaria de Ambiente y Cambio Climático de Provincia de Río Negro, Viedma, Argentina (M. Miqueo Lauriente)

DOI: <https://doi.org/10.3201/eid3004.231725>

We report full-genome characterization of highly pathogenic avian influenza A(H5N1) clade 2.3.4.4b virus from an outbreak among sea lions (August 2023) in Argentina and possible spillover to fur seals and terns. Mammalian adaptation mutations in virus isolated from marine mammals and a human in Chile were detected in mammalian and avian hosts.

In February 2023, the first case of highly pathogenic avian influenza (HPAI) A(H5N1) in Argentina was detected in a wild goose near the border with Bolivia and Chile (Appendix Figure 1, <https://wwwnc.cdc.gov/EID/article/30/4/23-1725-App1.pdf>) (1). In contrast with Peru and Chile, where extensive mortality of seabirds and marine mammals had been attributed to the virus in the preceding months (2,3), the initial spread of HPAI H5N1 in Argentina was largely limited to backyard and industrial poultry (94 outbreaks), causing the death or disposal of 2.2 million birds. Argentina declared itself free from the disease in poultry on August 8, 2023; before then, HPAI H5N1 detections in wildlife in Argentina had been scarce (7 events during February–April) and limited to aquatic birds (Anatidae, Laridae, and Rallidae) (1,4). However, soon thereafter, the national animal health services confirmed HPAI H5N1 in South American sea lions (*Otaria byronia*) from Río Grande, southernmost Argentina. Over subsequent weeks, the virus was detected in sea lions northward along the Argentina coast, and sporadic cases also occurred in South American fur seals (*Arctocephalus australis*). The most affected site was Punta Bermeja (Appendix Figure 1), the largest sea lion colony in Argentina, where an estimated 811 sea lions died over 2 months; minimal numbers (<5) of fur seals and terns were also affected (1,4).

In collaboration with provincial authorities and park rangers, we collected swab samples (oronasal, rectal, tracheal, lung, and brain) from 16 deceased sea lions, 1 fur seal, 1 great grebe (*Podiceps major*), and 1 South American tern (*Sterna hirundinacea*) discovered at Punta Bermeja on August 26, 2023. A sampled adult male sea lion was seen alive showing clinical signs consistent with HPAI infection (inability to stand or walk, muscular tremors and spasms, difficulty breathing, and abundant oral mucus). We tested the samples by real-time reverse transcription PCR targeting influenza A virus (5) and confirmed that all were positive. On the basis of viral RNA yields, we selected brain samples from 4 sea lions, 1 fur seal, and 1 tern for full-genome sequencing (Appendix Figure 2). We used maximum-likelihood tree phylogenetic analysis (6) and mutational analysis




Figure. Maximum-likelihood trees for hemagglutinin (A) and polymerase basic 2 (B) gene segments evaluated in study of highly pathogenic avian influenza A(H5N1) in Argentina compared with reference strains from other countries in South America. Tree areas have been enlarged at right to show detail. Red arrows indicate virus from marine mammals in Argentina; red asterisk indicates virus from a tern in Argentina. Black arrowhead along full tree in panel A indicates the hemagglutinin sequence from the first detection of HPAI H5N1 in a wild goose in Argentina. Node shape represents host group, and node color (and bars adjacent to trees) represents the region/country. Branch lengths are drawn proportionally to the extent of changes. Values adjacent to nodes represent bootstrap support ≥ 40 . Scale bars indicate nucleotide substitutions per site.

to compare the sequences (GenBank accession nos. OR987081–128) with representative HPAI H5N1 strains from South America.

Phylogenetic trees (Figure; Appendix Figure 2) showed that the viruses we identified belong to HPAI H5N1 clade 2.3.4.4b and are closely related to H5N1 viruses that circulated in South America during 2022–2023. Our finding supports the hypothesis that, after introduction from North America into Peru in November 2022, HPAI H5N1 viruses continued spreading across the continent and into Argentina. Of note, the viruses from Punta Bermeja did not cluster with the hemagglutinin and neuraminidase sequences available from HPAI H5N1 first detected in a wild goose in Argentina. Instead,

all gene segments from the viruses were closely related to virus sequences from sea lions in Chile and Peru (2; C. Pardo Roa, unpub. data, <https://www.biorxiv.org/content/10.1101/2023.06.30.547205v1>); 6 gene segments (all except polymerase basic protein 1 and nucleocapsid protein) also clustered with the virus isolated from a human in Chile (7). That finding suggests that viruses from Punta Bermeja may have been derived from a separate HPAI H5N1 introduction into Argentina. Because of the lack of genomic data for HPAI H5N1 viruses circulating in Argentina during February–July 2023, the finer scale pathways (local geographic routes and host species involved) of how these viruses arrived at Punta Bermeja remain unclear. Even so, the viruses

that we report did not cluster with those from birds in Uruguay, Brazil, or Bird Island (Antarctica), possibly suggesting separate pathways of virus spread.

On the basis of previous comparisons with HPAI H5N1 isolates from other countries in South America, we identified 9 mutations already present in viruses infecting sea lions in Peru and Chile but not in the goose/Guangdong reference strain or in viruses from birds and mammals from North America in 2022 (Table, <https://wwwnc.cdc.gov/eid/article/30/4/23-1725-t1>). Specifically, we found Q591K and D701N mutations in polymerase basic 2 associated with increased pathogenicity to mammals (8). The virus we detected in the South American tern also has those mutations, but they were absent from previously reported HPAI H5N1 viruses from avian hosts in South America (except for A/sanderling/Arica y Parinacota/240265/2023, which has the D701N mutation). That finding further supports the hypothesis that HPAI H5N1 viruses from sea lions from Peru and Chile acquired mammalian adaptation mutations that improved their ability to infect pinnipeds while possibly retaining the ability to infect avian hosts. Given the rapid and widespread dissemination of the viruses among pinnipeds in South America and the substantial associated mortalities (3,9), it seems likely that pinniped-to-pinniped transmission played a role in the spread of the mammal-adapted HPAI H5N1 viruses in the region. It is alarming that the HPAI H5N1 viruses infecting pinnipeds and seabirds in Argentina share the same mammalian adaptation mutations as the virus from the affected human in Chile, which highlights the potential threat posed by these viruses to public health.

Acknowledgments

We gratefully acknowledge all data contributors (i.e., the authors and their originating laboratories responsible for obtaining the specimens, and their submitting laboratories for generating the genetic sequence and metadata and sharing via the GISAID Initiative [<https://www.gisaid.org>], on which this research is based). We thank the Secretaría de Ambiente y Cambio Climático de la Provincia de Río Negro, especially Dina Migani and Federico Hollman, and Sebastián Ortega from Área Natural Protegida Bahía de San Antonio.

M.M.U. and A.R. designed the study and contributed funding; R.E.T.V., A.D., and M.M.L. collected samples; M.M.L. contributed surveillance data; V.O. performed the laboratory analyses; A.R. and R.E.T.V. performed the

bioinformatic analyses; A.R., R.E.T.V., and M.M.U. wrote the manuscript. All authors have read, revised, and approved the manuscript.

About the Author

Dr. Rimondi is a scientist at the National Institute of Agricultural Technology in Argentina and a postdoctoral fellow from Alexander von Humboldt Foundation from Germany working on HPAI H5N1 at the Robert Koch Institute. Her primary research interests focus on molecular epidemiology and host-pathogen interactions of avian influenza viruses.

References

1. Servicio Nacional de Sanidad y Calidad Agroalimentaria. State of the epidemiological situation in Argentina [in Spanish] [cited 2023 Dec 12]. <https://www.argentina.gob.ar/senasa/estado-de-la-situacion-epidemiologica-en-la-argentina>
2. Leguia M, Garcia-Glaessner A, Muñoz-Saavedra B, Juárez D, Barrera P, Calvo-Mac C, et al. Highly pathogenic avian influenza A (H5N1) in marine mammals and seabirds in Peru. *Nat Commun.* 2023;14:5489. <https://doi.org/10.1038/s41467-023-41182-0>
3. Ulloa M, Fernández A, Ariyama N, Colom-Rivero A, Rivera C, Nuñez P, et al. Mass mortality event in South American sea lions (*Otaria flavescens*) correlated to highly pathogenic avian influenza (HPAI) H5N1 outbreak in Chile. *Vet Q.* 2023;43:1–10. <https://doi.org/10.1080/01652176.2023.2265173>
4. World Organisation for Animal Health. WAHIS: World Animal Health Information System [cited 2023 Dec 12]. <https://wahis.woah.org>
5. Pan American Health Organization. CDC protocol for real-time RT-PCR for the new subtype of influenza A(H1N1) virus [in Spanish] [cited 2023 Dec 12]. <https://www.paho.org/es/node/36933>
6. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol.* 2015;32:268–74. <https://doi.org/10.1093/molbev/msu300>
7. Castillo A, Fasce R, Parra B, Andrade W, Covarrubias P, Hueche A, et al. The first case of human infection with H5N1 avian Influenza A virus in Chile. *J Travel Med.* 2023; 30(5):taad083. <https://doi.org/10.1093/jtm/taad083>
8. Lee CY, An SH, Choi JG, Lee YJ, Kim JH, Kwon HJ. Rank orders of mammalian pathogenicity-related PB2 mutations of avian influenza A viruses. *Sci Rep.* 2020;10:5359. <https://doi.org/10.1038/s41598-020-62036-5>
9. Campagna C, Uhart M, Falabella V, Campagna J, Zavattieri V, Vanstreels RET, et al. Catastrophic mortality of southern elephant seals caused by H5N1 avian influenza. *Mar Mamm Sci.* 2024;40:322–5. <https://doi.org/10.1111/mms.13101>

Address for correspondence: Agustina Rimondi, Robert Koch Institute, Holländerstrasse 102, 13407, Berlin, Germany; email: rimondia@rki.de or rimondi.agustina@inta.gob.ar

High-Pathogenicity Avian Influenza A(H5N1) Viruses in Multispecies Outbreak, Argentina, August 2023

Appendix

Details on the laboratory methods employed for the detection and full-genome characterization of HPAI H5N1 viruses in samples from marine mammals and seabirds at Punta Bermeja, Argentina.

Biosecurity

Beach surveys, documentation of clinical cases and sample collection from affected animals were performed by trained veterinarians in full personal protective equipment (hooded tyvek suits, double gloves, N95 respirators, goggles and rubber boots), and under government permits to perform these activities. All swabs collected were placed in cryotubes containing 1 mL of DNA/RNA Shield (Zymo Research, Irvine, CA, USA) for inactivation, then placed in individual ziploc bags (all samples per individual animal), sprayed with 10% bleach, then placed in a second ziploc bag that was stored in a cooler with icepacks. Personal protective equipment and disposable items were discarded as pathogenic waste, and all reusable items (boots, bucket, cooler, knives) were thoroughly washed and scrubbed with water, detergent and a handbrush, and then disinfected with Lysol and 10% bleach before leaving the site. Inactivated samples were transferred to an ultralow freezer (-80°C) within 24 hours after collection, with all samples from a single collection event in double ziploc bags sprayed with 10% bleach. At the laboratory, initial RNA extraction was performed under laminar flow and samples were added to lysis buffer from the QIAamp Viral RNA Mini Kit (Qiagen Inc., Valencia, CA, USA). After waiting the appropriate time for sample inactivation, further steps were performed without laminar flow.

Virus detection

Viral RNA was extracted from 140 µl of suspension from pooled swabs (Appendix Table 1) using a QIAamp Viral RNA Mini Kit Qiagen Inc., Valencia, CA, USA). RNA was eluted in a final volume of 60 µl and stored at –80°C until further use. Viral cDNA was prepared using 15 µl of viral RNA and random hexamers in a final volume of 30 µl using a High Capacity cDNA Archive kit (Applied Biosystems, Foster City, CA, USA). The cDNA was tested for IAV by real-time reverse transcription PCR (RRT-PCR) on an ABI Prism 7500 SDS (Applied Biosystems) using TaqMan Universal PCR Master Mix (Applied Biosystems) directed to the matrix (M) gene (*I*). Quantification cycles (Cq) values were used as a proxy to compare viral RNA load in different samples (Appendix Table 1). Based on average viral RNA loads (summarized in Appendix Table 2), brain samples were selected for further testing. Viral RNA was then extracted from individual brain swabs of four South American sea lions (RN-PB004, RN-PB007, RN-PB011, RN-PB013), one South American fur seal (RN-PB019) and one South American tern (RN-PB015) for further full genome sequencing. The four South American sea lions were randomly selected within age-sex cohorts (i.e., a sub-adult male, a sub-adult female, an adult male, and an adult female). No further analyses were performed on the samples from the great grebe due to low viral RNA yields.

Full genome sequencing

The viral genome was amplified from RNA using a multi-segment one-step RT-PCR with Superscript III high-fidelity RT-PCR kit (Invitrogen, Carlsbad CA) according to manufacturer's instructions using the Opti1 primer set (Opti1-F1, Opti1-F2 and Opti1-R1) as previously described by Zhou and colleagues (2). The RT-PCR amplification parameters were: 2 min at 55°C, 60 min at 42°C, and 2 min at 94°C, followed by 5 cycles of (94°C/30 s; 44°C/30 s; 68°C/3.5 min), 26 cycles of (94°C/30 s; 57°C/30 s; 68°C/3.5 min), and a final extension for 10 min at 68°C. Amplicons were visualized on a 1% agarose gel and purified with Agencourt AMPure XP beads (Beckman Coulter, Brea, CA). The concentration of purified amplicons was quantified using the Qubit High Sensitivity dsDNA kit and a Qubit Fluorometer (Invitrogen). The sequencing library preparation was done with the Oxford Nanopore Rapid Barcode library kit SQH-RBK110.96 and loaded on the Mk1c sequencer according to ONT instructions for the R.9 flow cells. Then, real time basecalling was performed with the MinIT to produce fast5 and fastQ files. The automatic real time division into passed and failed reads by the MinIT works as a

quality check, removing reads with quality scores lower than 7. The quality checked reads were demultiplexed and trimmed for adapters and primers, followed by mappings and a final consensus production using Qiagen CLC Genomics Workbench v23.0.2 (Qiagen, Hilden, Germany).

Phylogenetic analysis

Publicly available sequences (as of 9 December 2023) were obtained from GenBank (<https://www.ncbi.nlm.nih.gov/genbank/>) and the Global Initiative on Sharing All Influenza Data (<https://gisaid.org>) (3). All available sequences from influenza A viruses obtained from samples collected in South America since 1 January 2022 were included, as well as representative samples from other continents (accession codes provided in Appendix Table 3). Sequences were aligned with MAFFT v7.520 (<https://mafft.cbrc.jp/alignment/software/>) using the FFT-NS-2 algorithm (4). Maximum Likelihood trees were built using IQ-TREE v2.2.2.6 (<http://www.iqtree.org/>), using the integrated ModelFinder for model selection (5) and performing 5000 ultra-fast bootstrap repetitions.

References

- <eref>1. World Health Organization (WHO). Protocolo del CDC para el RT-PCR en tiempo real para el nuevo subtipo del virus de influenza A(H1N1): Revisión 1 (30 de abril de 2009) [cited 2023 Dec 12]. <https://www.paho.org/es/node/36933></eref>
- <jrn>2. Zhou B, Donnelly ME, Scholes DT, St George K, Hatta M, Kawaoka Y, et al. Single-reaction genomic amplification accelerates sequencing and vaccine production for classical and Swine origin human influenza a viruses. *J Virol.* 2009;83:10309–13. 10.1128/JVI.01109-09 [PubMed](#) <https://doi.org/10.1128/JVI.01109-09></jrn>
- <jrn>3. Shu Y, McCauley J. GISAID: Global initiative on sharing all influenza data - from vision to reality. *Euro Surveill.* 2017;22:30494. 10.2807/1560-7917.ES.2017.22.13.30494 [PubMed](#) <https://doi.org/10.2807/1560-7917.ES.2017.22.13.30494></jrn>
- <jrn>4. Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res.* 2002;30:3059–66. 10.1093/nar/gkf436 [PubMed](#) <https://doi.org/10.1093/nar/gkf436></jrn>

<jrn>5. Kalyanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermini LS. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods*. 2017;14:587–9. 10.1038/nmeth.4285 [PubMed https://doi.org/10.1038/nmeth.4285](https://doi.org/10.1038/nmeth.4285)</jrn>

Appendix Table 1. Details of Cq values obtained by RRT-PCR from pooled samples of wildlife collected at Punta Bermeja, Rio Negro, Argentina, in August 2023

| Pool | ID | Host species | Age | Sex | Swab | Cq |
|------|----------|-------------------------|-----------|--------|----------|-------|
| A | RN-PB002 | South American sea lion | Adult | Female | Oronasal | 27.24 |
| | RN-PB006 | South American sea lion | Sub-adult | Female | Oronasal | |
| | RN-PB007 | South American sea lion | Sub-adult | Female | Oronasal | |
| | RN-PB013 | South American sea lion | Adult | Female | Oronasal | |
| | RN-PB016 | South American sea lion | Adult | Female | Oronasal | |
| B | RN-PB005 | South American sea lion | Adult | Male | Oronasal | 26.14 |
| | RN-PB008 | South American sea lion | Sub-adult | Male | Oronasal | |
| | RN-PB011 | South American sea lion | Adult | Male | Oronasal | |
| | RN-PB012 | South American sea lion | Adult | Male | Oronasal | |
| | RN-PB017 | South American sea lion | Adult | Male | Oronasal | |
| | RN-PB018 | South American sea lion | Adult | Male | Oronasal | |
| C | RN-PB003 | South American sea lion | Juvenile | Female | Oronasal | 25.72 |
| | RN-PB004 | South American sea lion | Sub-adult | Male | Oronasal | |
| | RN-PB009 | South American sea lion | Adult | Male | Oronasal | |
| | RN-PB010 | South American sea lion | Juvenile | Female | Oronasal | |
| | RN-PB019 | South American fur seal | Juvenile | Male | Oronasal | |
| D | RN-PB002 | South American sea lion | Adult | Female | Rectal | 24.66 |
| | RN-PB006 | South American sea lion | Sub-adult | Female | Rectal | |
| | RN-PB007 | South American sea lion | Sub-adult | Female | Rectal | |
| | RN-PB013 | South American sea lion | Adult | Female | Rectal | |
| | RN-PB016 | South American sea lion | Adult | Female | Rectal | |
| E | RN-PB005 | South American sea lion | Adult | Male | Rectal | 30.71 |
| | RN-PB008 | South American sea lion | Sub-adult | Male | Rectal | |
| | RN-PB011 | South American sea lion | Adult | Male | Rectal | |
| | RN-PB012 | South American sea lion | Adult | Male | Rectal | |
| | RN-PB017 | South American sea lion | Adult | Male | Rectal | |
| | RN-PB018 | South American sea lion | Adult | Male | Rectal | |
| F | RN-PB003 | South American sea lion | Juvenile | Female | Rectal | 30.92 |
| | RN-PB004 | South American sea lion | Sub-adult | Male | Rectal | |
| | RN-PB009 | South American sea lion | Adult | Male | Rectal | |
| | RN-PB010 | South American sea lion | Juvenile | Female | Rectal | |
| | RN-PB019 | South American fur seal | Juvenile | Male | Rectal | |
| G | RN-PB002 | South American sea lion | Adult | Female | Tracheal | 22.97 |
| | RN-PB006 | South American sea lion | Sub-adult | Female | Tracheal | |
| | RN-PB007 | South American sea lion | Sub-adult | Female | Tracheal | |
| | RN-PB013 | South American sea lion | Adult | Female | Tracheal | |
| | RN-PB016 | South American sea lion | Adult | Female | Tracheal | |
| H | RN-PB005 | South American sea lion | Adult | Male | Tracheal | 19.13 |
| | RN-PB008 | South American sea lion | Sub-adult | Male | Tracheal | |
| | RN-PB011 | South American sea lion | Adult | Male | Tracheal | |
| | RN-PB012 | South American sea lion | Adult | Male | Tracheal | |
| | RN-PB017 | South American sea lion | Adult | Male | Tracheal | |
| | RN-PB018 | South American sea lion | Adult | Male | Tracheal | |
| I | RN-PB003 | South American sea lion | Juvenile | Female | Tracheal | 23.42 |
| | RN-PB004 | South American sea lion | Sub-adult | Male | Tracheal | |
| | RN-PB009 | South American sea lion | Adult | Male | Tracheal | |
| | RN-PB010 | South American sea lion | Juvenile | Female | Tracheal | |
| | RN-PB019 | South American fur seal | Juvenile | Male | Tracheal | |
| J | RN-PB002 | South American sea lion | Adult | Female | Lung | 21.58 |
| | RN-PB006 | South American sea lion | Sub-adult | Female | Lung | |
| | RN-PB007 | South American sea lion | Sub-adult | Female | Lung | |
| | RN-PB013 | South American sea lion | Adult | Female | Lung | |
| | RN-PB016 | South American sea lion | Adult | Female | Lung | |
| K | RN-PB005 | South American sea lion | Adult | Male | Lung | 21.87 |
| | RN-PB008 | South American sea lion | Sub-adult | Male | Lung | |
| | RN-PB011 | South American sea lion | Adult | Male | Lung | |
| | RN-PB012 | South American sea lion | Adult | Male | Lung | |
| | RN-PB017 | South American sea lion | Adult | Male | Lung | |

| Pool | ID | Host species | Age | Sex | Swab | Cq |
|------|----------|-------------------------|-----------|---------|-------|-------|
| L | RN-PB018 | South American sea lion | Adult | Male | Lung | 24.31 |
| | RN-PB003 | South American sea lion | Juvenile | Female | Lung | |
| | RN-PB004 | South American sea lion | Sub-adult | Male | Lung | |
| | RN-PB009 | South American sea lion | Adult | Male | Lung | |
| | RN-PB010 | South American sea lion | Juvenile | Female | Lung | |
| M | RN-PB019 | South American fur seal | Juvenile | Male | Lung | 19.30 |
| | RN-PB002 | South American sea lion | Adult | Female | Brain | |
| | RN-PB006 | South American sea lion | Sub-adult | Female | Brain | |
| | RN-PB007 | South American sea lion | Sub-adult | Female | Brain | |
| | RN-PB013 | South American sea lion | Adult | Female | Brain | |
| N | RN-PB016 | South American sea lion | Adult | Female | Brain | 21.84 |
| | RN-PB005 | South American sea lion | Adult | Male | Brain | |
| | RN-PB008 | South American sea lion | Sub-adult | Male | Brain | |
| | RN-PB011 | South American sea lion | Adult | Male | Brain | |
| | RN-PB012 | South American sea lion | Adult | Male | Brain | |
| O | RN-PB017 | South American sea lion | Adult | Male | Brain | 18.87 |
| | RN-PB018 | South American sea lion | Adult | Male | Brain | |
| | RN-PB003 | South American sea lion | Juvenile | Female | Brain | |
| | RN-PB004 | South American sea lion | Sub-adult | Male | Brain | |
| | RN-PB009 | South American sea lion | Adult | Male | Brain | |
| P | RN-PB010 | South American sea lion | Juvenile | Female | Brain | 17.56 |
| | RN-PB019 | South American fur seal | Juvenile | Male | Brain | |
| | RN-PB014 | South American sea lion | Fetus | Unknown | All | |
| Q | RN-PB015 | South American tern | Juvenile | Male | All | 27.74 |
| R | RN-PB020 | Great Grebe | Adult | Male | All | 35.88 |

Appendix Table 2. Average Cq values obtained by RRT-PCR from pooled samples collected from different tissues of sea lions at Punta Bermeja, Rio Negro, Argentina, in August 2023

| Statistic | Rectal | Oronasal | Lung | Tracheal | Brain |
|--------------------|--------|----------|------|----------|-------|
| Mean | 28.8 | 26.4 | 22.6 | 21.8 | 20.0 |
| Standard deviation | 1.6 | 1.5 | 0.8 | 3.6 | 2.4 |


Appendix Table 3. Accession codes for publicly-available gene sequences from GenBank and the Global Initiative on Sharing All Influenza Data used in the phylogenetic and mutation analyses.

| Isolate | Accession code |
|---|-------------------|
| A/Thalasseu acutiflavus/EspiritoSanto/1339 N2/2023 H5N1 Brazil 2023-05-15 | OR269884-OR269891 |
| A/Goose/Guangdong/1/96 H5N1 China 1996-01-01 | EPI_ISL_1254 |
| A/goose/Argentina/140223/2023 H5N1 Argentina 2023-02-11 | EPI_ISL_17527083 |
| A/Chile/25945/2023 H5N1 Chile 2023-03-24 | EPI_ISL_17468386 |
| A/sea lion/Tarapaca/240524-2/2023 H5N1 Chile 2023-03-07 | EPI_ISL_17885975 |
| A/sea lion/Arica y Parinacota/240270-1/2023 H5N1 Chile 2023-03-01 | EPI_ISL_17885976 |
| A/sanderling/Arica y Parinacota/240265/2023 H5N1 Chile 2023-03-03 | EPI_ISL_17885978 |
| A/sea lion/Peru/AQP-SER00R/2023 H5N1 Peru 2023-03-06 | EPI_ISL_18054510 |
| A/sea lion/Peru/TAC-INS-011/2023 H5N1 Peru 2023-03-07 | EPI_ISL_17777532 |
| A/sea lion/Peru/TAC-INS-010/2023 H5N1 Peru 2023-03-07 | EPI_ISL_17777531 |
| A/sea lion/Peru/AQP-SER00K/2023 H5N1 Peru 2023-02-07 | EPI_ISL_18054509 |
| A/Sea Lion/Peru/LIM-SER036/2023 H5N1 Peru 2023-01-23 | EPI_ISL_18054502 |
| A/dolphin/Peru/PIU-SER002/2022 H5N1 Peru 2022-11-22 | EPI_ISL_18265431 |
| A/lion/Peru/AIS0554/2023 H5N1 Peru 2023-02-08 | EPI_ISL_17805999 |
| A/pelican/Peru/PIU-SER019/2022 H5N1 Peru 2022-11-24 | EPI_ISL_18265435 |
| A/Brown skua/Bird Island/128287/2023 H5N1 Antarctica 2023-10-08 | EPI_ISL_18439562 |
| A/black-necked swan/Uruguay/UDELAR-078-M2/2023 H5N1 Uruguay 2023-03-15 | EPI_ISL_18310957 |
| A/Ecuador/6563/2023 H5N1 Ecuador 2023-01-05 | EPI_ISL_17021605 |
| A/wildbird-Fregata-magnificens/Ecuador/IC03-4587/2023 H5N1 Ecuador 2023-01-11 | EPI_ISL_17973443 |
| A/duck/Choco/ICA-3501/2022 H5N1 Colombia 2022-10-09 | EPI_ISL_17353839 |
| A/Pelican/Venezuela/Pel3/2022 H5N1 Venezuela 2022-11-25 | EPI_ISL_16013752 |
| A/Colorado/18/2022 H5N1 USA 2022-04-20 | EPI_ISL_12799972 |
| A/skunk/Washington/22-019274-001-original/2022 H5N1 USA 2022-06-07 | EPI_ISL_15078254 |
| A/harbor seal/Washington/23-025991-001-original/2023 H5N1 USA 2023-08-25 | EPI_ISL_18311025 |
| A/dolphin/Florida/22-025319-002-original/2022 H5N1 USA 2022-03-30 | EPI_ISL_15078255 |
| A/chicken/Wyoming/22-009599-002-original/2022 H5N1 USA 2022-03-26 | EPI_ISL_13009694 |
| A/Mink/Spain/3691-2 22VIR10586-8/2022 H5N1 Spain 2022-10-18 | EPI_ISL_15878541 |
| A/gray seal/Netherlands/30448/2023 H5N1 Netherlands 2023-01-01 | EPI_ISL_17672783 |
| A/common dolphin/Wales/040498/2023 H5N1 UK 2023-02-20 | EPI_ISL_17465833 |

| Isolate | Accession code |
|---|------------------|
| A/Brown skua/Bird Island/128288/2023 H5N1 Antarctica 2023-10-08 | EPI_ISL_18439563 |
| A/Brown skua/Bird Island/128289/2023 H5N1 Antarctica 2023-10-08 | EPI_ISL_18439564 |
| A/pelican/Peru/PIU-SER028/2022 H5N1 Peru 2022-11-24 | EPI_ISL_18265436 |
| A/pelican/Peru/PIU-SER016/2022 H5N1 Peru 2022-11-24 | EPI_ISL_18265437 |
| A/Guanay cormorant/Peru/PIU-SER024/2022 H5N1 Peru 2022-11-24 | EPI_ISL_18265434 |
| A/Sanderling/Peru/PIU-SER005/2022 H5N1 Peru 2022-11-22 | EPI_ISL_18265432 |
| A/pelican/Peru/PIU-SER013/2022 H5N1 Peru 2022-11-23 | EPI_ISL_18265433 |
| A/falcon/Peru/A273/2022 H5N1 Peru 2022-12-01 | EPI_ISL_17526123 |
| A/owl/Peru/A293/2022 H5N1 Peru 2022-12-01 | EPI_ISL_17526122 |
| A/sanderling/Peru/PIU-005/2022 H5N1 Peru 2022-11-22 | EPI_ISL_18054501 |
| A/guanay cormorant/Peru/PIU-024/2022 H5N1 Peru 2022-11-24 | EPI_ISL_18054500 |
| A/pelican/Peru/PIU-028/2022 H5N1 Peru 2022-11-24 | EPI_ISL_18054507 |
| A/pelican/Peru/PIU-019/2022 H5N1 Peru 2022-11-24 | EPI_ISL_18054506 |
| A/Pelecanus occidentalis/Venezuela/3S2/2022 H5N1 Venezuela 2022-11-25 | EPI_ISL_16854395 |
| A/pelican/Peru/PIU-013/2022 H5N1 Peru 2022-11-23 | EPI_ISL_18054504 |
| A/Pelecanus occidentalis/Venezuela/3S1/2022 H5N1 Venezuela 2022-11-25 | EPI_ISL_16854394 |
| A/Pelecanus occidentalis/Venezuela/3S4/2022 H5N1 Venezuela 2022-11-25 | EPI_ISL_16854397 |
| A/Pelecanus occidentalis/Venezuela/3S3/2022 H5N1 Venezuela 2022-11-25 | EPI_ISL_16854396 |
| A/Pelecanus occidentalis/Venezuela/3S6/2022 H5N1 Venezuela 2022-11-25 | EPI_ISL_16854399 |
| A/Pelecanus occidentalis/Venezuela/3S5/2022 H5N1 Venezuela 2022-11-25 | EPI_ISL_16854398 |
| A/duck/Peru/LAM-INS-014/2023 H5N1 Peru 2023-09-26 | EPI_ISL_18497946 |
| A/Pelecanus occidentalis/Venezuela/3S8/2022 H5N1 Venezuela 2022-11-25 | EPI_ISL_16854401 |
| A/Pelecanus occidentalis/Venezuela/3S7/2022 H5N1 Venezuela 2022-11-25 | EPI_ISL_16854400 |
| A/Pelecanus occidentalis/Venezuela/4S2/2022 H5N1 Venezuela 2022-11-25 | EPI_ISL_16854403 |
| A/Pelecanus occidentalis/Venezuela/4S1/2022 H5N1 Venezuela 2022-11-25 | EPI_ISL_16854402 |
| A/duck/Peru/LAM-INS-015/2023 H5N1 Peru 2023-09-20 | EPI_ISL_18497950 |
| A/Pelecanus occidentalis/Venezuela/4S4/2022 H5N1 Venezuela 2022-11-25 | EPI_ISL_16854405 |
| A/Pelecanus occidentalis/Venezuela/4S3/2022 H5N1 Venezuela 2022-11-25 | EPI_ISL_16854404 |
| A/Pelecanus occidentalis/Venezuela/4S6/2022 H5N1 Venezuela 2022-11-25 | EPI_ISL_16854407 |
| A/Pelecanus occidentalis/Venezuela/4S5/2022 H5N1 Venezuela 2022-11-25 | EPI_ISL_16854406 |
| A/Pelecanus occidentalis/Venezuela/4S8/2022 H5N1 Venezuela 2022-11-25 | EPI_ISL_16854409 |
| A/Pelecanus occidentalis/Venezuela/4S7/2022 H5N1 Venezuela 2022-11-25 | EPI_ISL_16854408 |
| A/american kestrel/Peru/UNMSM-A273/2022 H5N1 Peru 2022-12-01 | EPI_ISL_18238548 |
| A/Belcher gull/Peru/UNMSM-A267/2022 H5N1 Peru 2022-12-01 | EPI_ISL_18371664 |
| A/Guanay cormorant/Peru/UNMSM-A275/2022 H5N1 Peru 2022-12-01 | EPI_ISL_18371665 |
| A/Peruvian booby/Peru/UNMSM-A296/2022 H5N1 Peru 2022-12-01 | EPI_ISL_18371666 |
| A/Pelican/Venezuela/Pel4/2022 H5N1 Venezuela 2022-11-25 | EPI_ISL_16013753 |
| A/pelican/Peru/PIU-INS-001/2022 H5N1 Peru 2022-11-12 | EPI_ISL_16249274 |
| A/chicken/Magdalena/ICA-3503/2022 H5N1 Colombia 2022-11-18 | EPI_ISL_17353838 |
| A/belchers gull/Peru/UNMSM-A102/2022 H5N1 Peru 2022-11-01 | EPI_ISL_18238610 |
| A/duck/Peru/CAL-INS-013/2023 H5N1 Peru 2023-07-24 | EPI_ISL_18238540 |
| A/peruvian pelican/Peru/UNMSM-A074/2022 H5N1 Peru 2022-11-01 | EPI_ISL_18238608 |
| A/belchers gull/Peru/UNMSM-A267/2022 H5N1 Peru 2022-12-01 | EPI_ISL_18238609 |
| A/western barn owl/Peru/UNMSM-A293/2022 H5N1 Peru 2022-12-01 | EPI_ISL_18238622 |
| A/peruvian pelican/Peru/UNMSM-A106/2022 H5N1 Peru 2022-11-01 | EPI_ISL_18238607 |
| A/owl/Peru/23-005629-001-original/2022 H5N1 Peru 2022-11-11 | EPI_ISL_17660073 |
| A/chicken/Peru/23-005607-003-original/2022 H5N1 Peru 2022-11-11 | EPI_ISL_17660072 |
| A/brown booby/Peru/23-005629-003-original/2022 H5N1 Peru 2022-11-11 | EPI_ISL_17660074 |
| A/chicken/Peru/23-005607-002-original/2022 H5N1 Peru 2022-11-11 | EPI_ISL_17660071 |
| A/chicken/Peru/23-005607-001-original/2022 H5N1 Peru 2022-11-11 | EPI_ISL_17660070 |
| A/Pelecanus/Peru/VFAR-140/2022 H5N1 Peru 2022-12-01 | EPI_ISL_17099964 |
| A/wildbird-Fregata-magnificens/Ecuador/IC06-4590/2023 H5N1 Ecuador 2023-01-11 | EPI_ISL_17973458 |
| A/Gull/Chile/7023-3/2022 H5N1 Chile 2022-12-07 | EPI_ISL_17011958 |
| A/Gull/Chile/7023-2/2022 H5N1 Chile 2022-12-07 | EPI_ISL_17011964 |
| A/turkey vulture/Valparaiso/230187-1/2022 H5N1 Chile 2022-12-31 | EPI_ISL_17885968 |
| A/domestic duck/Maule/240466-1/2023 H5N1 Chile 2023-03-07 | EPI_ISL_18005779 |
| A/heron/Antofagasta/228705-3/2022 H5N1 Chile 2022-12-20 | EPI_ISL_17885969 |
| A/turkey vulture/Antofagasta/228252-1/2022 H5N1 Chile 2022-12-17 | EPI_ISL_17885970 |
| A/heron/Antofagasta/228705-2/2022 H5N1 Chile 2022-12-20 | EPI_ISL_17885971 |
| A/kelp gull/Maule/239349/2023 H5N1 Chile 2023-03-01 | EPI_ISL_17885972 |
| A/domestic duck/Araucania/239189-3/2023 H5N1 Chile 2023-02-28 | EPI_ISL_18005783 |
| A/tern/Maule/238507/2023 H5N1 Chile 2023-02-23 | EPI_ISL_17885973 |
| A/domestic duck/Araucania/241914-2/2023 H5N1 Chile 2023-03-14 | EPI_ISL_18005780 |
| A/domestic duck/Araucania/240481-2/2023 H5N1 Chile 2023-03-07 | EPI_ISL_18005781 |
| A/black crowned night heron/Antofagasta/228705-2/2022 H5N1 Chile 2022-12-20 | EPI_ISL_18005786 |
| A/backyard chicken/Uruguay/UDELAR-040-M5/2023 H5N1 Uruguay 2023-03-03 | EPI_ISL_18310942 |
| A/black crowned night heron/Antofagasta/228705-3/2022 H5N1 Chile 2022-12-20 | EPI_ISL_18005784 |
| A/sanderling/Arica y Parinacota/230758-1/2022 H5N1 Chile 2022-12-30 | EPI_ISL_17885980 |
| A/pelican/Valparaiso/234040/2023 H5N1 Chile 2023-01-25 | EPI_ISL_17885982 |

| Isolate | Accession code |
|--|------------------|
| A/pelican/Valparaiso/233450-1/2023 H5N1 Chile 2023-01-23 | EPI_ISL_17885983 |
| A/chicken/Araucania/239569-2/2023 H5N1 Chile 2023-03-02 | EPI_ISL_17885952 |
| A/elegant tern/Tarapaca/229133-1/2022 H5N1 Chile 2022-12-23 | EPI_ISL_17885953 |
| A/chicken/Araucania/239569-1/2023 H5N1 Chile 2023-03-02 | EPI_ISL_17885954 |
| A/elegant tern/Arica y Parinacota/229476-1/2022 H5N1 Chile 2022-12-24 | EPI_ISL_17885955 |
| A/pelican/Valparaiso/233091-2/2023 H5N1 Chile 2023-01-20 | EPI_ISL_17885956 |
| A/duck/Maule/240466-1/2023 H5N1 Chile 2023-03-07 | EPI_ISL_17885957 |
| A/chicken/Araucania/239189-2/2023 H5N1 Chile 2023-02-28 | EPI_ISL_17885958 |
| A/duck/Araucania/241914-2/2023 H5N1 Chile 2023-03-14 | EPI_ISL_17885959 |
| A/whimbrel/Coquimbo/239964/2023 H5N1 Chile 2023-03-05 | EPI_ISL_17885960 |
| A/duck/Araucania/240481-2/2023 H5N1 Chile 2023-03-07 | EPI_ISL_17885961 |
| A/turkey/Nuble/241568-1/2023 H5N1 Chile 2023-03-10 | EPI_ISL_17885962 |
| A/duck/Araucania/239189-3/2023 H5N1 Chile 2023-02-28 | EPI_ISL_17885963 |
| A/turkey/Nuble/240489-1/2023 H5N1 Chile 2023-03-07 | EPI_ISL_17885964 |
| A/chiloe wigeon/OHiggins/240893-2/2023 H5N1 Chile 2023-03-09 | EPI_ISL_17885965 |
| A/turkey/Araucania/241892-3/2023 H5N1 Chile 2023-03-14 | EPI_ISL_17885966 |
| A/blackish oystercatcher/OHiggins/240628/2023 H5N1 Chile 2023-03-06 | EPI_ISL_17885967 |
| A/backyard chicken/Uruguay/UDELAR-040-M7/2023 H5N1 Uruguay 2023-03-03 | EPI_ISL_18310967 |
| A/backyard chicken/Uruguay/UDELAR-047-M1/2023 H5N1 Uruguay 2023-03-06 | EPI_ISL_18310966 |
| A/backyard chicken/Uruguay/UDELAR-047-M3/2023 H5N1 Uruguay 2023-03-06 | EPI_ISL_18310965 |
| A/backyard chicken/Uruguay/UDELAR-124-M1/2023 H5N1 Uruguay 2023-04-16 | EPI_ISL_18310964 |
| A/backyard chicken/Uruguay/UDELAR-127-M1/2023 H5N1 Uruguay 2023-04-17 | EPI_ISL_18310963 |
| A/backyard chicken/Uruguay/UDELAR-127-M4/2023 H5N1 Uruguay 2023-04-17 | EPI_ISL_18310962 |
| A/backyard chicken/Uruguay/UDELAR-144-M3/2023 H5N1 Uruguay 2023-05-03 | EPI_ISL_18310961 |
| A/backyard duck/Uruguay/UDELAR-124-M3/2023 H5N1 Uruguay 2023-04-16 | EPI_ISL_18310960 |
| A/chicken/Bolivar/ICA-3500/2022 H5N1 Colombia 2022-11-08 | EPI_ISL_17353507 |
| A/chicken/Choco/ICA-3502/2022 H5N1 Colombia 2022-11-13 | EPI_ISL_17353508 |
| A/chicken/Choco/ICA-3504/2022 H5N1 Colombia 2022-11-01 | EPI_ISL_17353509 |
| A/chicken/Cordoba/ICA-3499/2022 H5N1 Colombia 2022-11-26 | EPI_ISL_17353510 |
| A/pelican/Valparaiso/233447-2/2023 H5N1 Chile 2023-01-21 | EPI_ISL_17885985 |
| A/whimbrel/Valparaiso/239964/2023 H5N1 Chile 2023-03-05 | EPI_ISL_17885987 |
| A/backyard turkey/Uruguay/UDELAR-124-M6/2023 H5N1 Uruguay 2023-04-16 | EPI_ISL_18310959 |
| A/black-necked swan/Uruguay/UDELAR-014-M3/2023 H5N1 Uruguay 2023-02-18 | EPI_ISL_18310958 |
| A/peruvian booby/Peru/LIM-INS-012/2023 H5N1 Peru 2023-04-12 | EPI_ISL_17777533 |
| A/guanay cormorant/Peru/CAL-INS-009/2023 H5N1 Peru 2023-03-17 | EPI_ISL_17777530 |
| A/peruvian booby/Peru/CAL-INS-008/2023 H5N1 Peru 2023-03-17 | EPI_ISL_17777529 |
| A/peruvian booby/Peru/CAL-INS-007/2023 H5N1 Peru 2023-03-17 | EPI_ISL_17777528 |
| A/gull/Peru/LIM-INS-006/2023 H5N1 Peru 2023-03-16 | EPI_ISL_17777527 |
| A/peruvian booby/Peru/LIM-INS-005/2023 H5N1 Peru 2023-03-20 | EPI_ISL_17777526 |
| A/peruvian booby/Peru/LIM-INS-004/2023 H5N1 Peru 2023-04-03 | EPI_ISL_17777525 |
| A/Peruvian pelican/Chile/C61740/2022 H5N1 Chile 2022-12-02 | EPI_ISL_16891401 |
| A/black skimmer/Chile/C61962/2022 H5N1 Chile 2022-12-02 | EPI_ISL_16891402 |
| A/wildbird-Sula-nebouxii/Ecuador/7607/2023 H5N1 Ecuador 2023-05-18 | EPI_ISL_18137626 |
| A/wildbird-Sula-nebouxii/Ecuador/7611/2023 H5N1 Ecuador 2023-05-18 | EPI_ISL_18137671 |
| A/pelican/Peru/A074/2022 H5N1 Peru 2022-11-01 | EPI_ISL_17477219 |
| A/gull/Peru/A267/2022 H5N1 Peru 2022-12-01 | EPI_ISL_17477222 |
| A/gull/Peru/A102/2022 H5N1 Peru 2022-11-01 | EPI_ISL_17477220 |
| A/pelican/Peru/A106/2022 H5N1 Peru 2022-11-01 | EPI_ISL_17477221 |
| A/chicken/Peru/LIM-INS-003/2022 H5N1 Peru 2022-11-29 | EPI_ISL_16249730 |
| A/chicken/Ecuador/04/2022 H5N1 Ecuador 2022-11-25 | EPI_ISL_16161675 |
| A/chicken/Ecuador/03/2022 H5N1 Ecuador 2022-11-25 | EPI_ISL_16161673 |
| A/black skimmer/Maule/240379/2023 H5N1 Chile 2023-03-06 | EPI_ISL_17885849 |
| A/chicken/Ecuador/02/2022 H5N1 Ecuador 2022-11-25 | EPI_ISL_16157545 |
| A/chicken/OHiggins/241252-1/2023 H5N1 Chile 2023-03-12 | EPI_ISL_17885872 |
| A/chicken/Nuble/241681-1/2023 H5N1 Chile 2023-03-10 | EPI_ISL_17885873 |
| A/avian/Peru/AISA0451/2022 H5N1 Peru 2022-11-22 | EPI_ISL_17805998 |
| A/chicken/Nuble/241557-1/2023 H5N1 Chile 2023-03-10 | EPI_ISL_17885874 |
| A/avian/Peru/AISA0446/2022 H5N1 Peru 2022-11-22 | EPI_ISL_17805997 |
| A/chicken/Peru/AIS0549/2022 H5N1 Peru 2022-12-22 | EPI_ISL_17805996 |
| A/chicken/Peru/AIS0550/2022 H5N1 Peru 2022-12-12 | EPI_ISL_17805995 |
| A/chicken/Nuble/240684/2023 H5N1 Chile 2023-03-08 | EPI_ISL_17885876 |
| A/chicken/Peru/AIS0548/2022 H5N1 Peru 2022-12-22 | EPI_ISL_17805994 |
| A/chicken/Nuble/240155/2023 H5N1 Chile 2023-03-06 | EPI_ISL_17885877 |
| A/chicken/Nuble/239136/2023 H5N1 Chile 2023-02-27 | EPI_ISL_17885878 |
| A/chicken/Peru/AIS0547/2022 H5N1 Peru 2022-12-28 | EPI_ISL_17805993 |
| A/chicken/Peru/LAM-INS-002/2022 H5N1 Peru 2022-11-28 | EPI_ISL_16249681 |
| A/chicken/Peru/AIS0545/2022 H5N1 Peru 2022-12-03 | EPI_ISL_17805992 |
| A/chicken/Peru/AIS0544/2022 H5N1 Peru 2022-12-03 | EPI_ISL_17805991 |
| A/chicken/Peru/AIS0543/2022 H5N1 Peru 2022-12-01 | EPI_ISL_17805990 |

| Isolate | Accession code |
|---|------------------|
| A/chicken/Peru/AIS0542/2022 H5N1 Peru 2022-12-01 | EPI_ISL_17805989 |
| A/chicken/Peru/AIS0540/2022 H5N1 Peru 2022-11-28 | EPI_ISL_17805988 |
| A/chicken/Peru/AIS0539/2022 H5N1 Peru 2022-11-18 | EPI_ISL_17805987 |
| A/chicken/Peru/AIS0546/2022 H5N1 Peru 2022-12-18 | EPI_ISL_17805986 |
| A/pelican/Peru/AISA0464/2022 H5N1 Peru 2022-11-22 | EPI_ISL_17806003 |
| A/goose/Araucania/239189-1/2023 H5N1 Chile 2023-02-28 | EPI_ISL_17885869 |
| A/pelican/Peru/AIS0541/2022 H5N1 Peru 2022-11-16 | EPI_ISL_17806002 |
| A/chicken/OHiggins/241252-6/2023 H5N1 Chile 2023-03-12 | EPI_ISL_17885870 |
| A/chicken/Peru/AIS0551/2022 H5N1 Peru 2022-12-12 | EPI_ISL_17806001 |
| A/pelican/Peru/AIS0538/2022 H5N1 Peru 2022-11-10 | EPI_ISL_17806000 |
| A/chicken/OHiggins/241252-3/2023 H5N1 Chile 2023-03-12 | EPI_ISL_17885871 |
| A/pelican/OHiggins/233721-1/2023 H5N1 Chile 2023-01-24 | EPI_ISL_17885912 |
| A/pelican/OHiggins/233663-1/2023 H5N1 Chile 2023-01-24 | EPI_ISL_17885915 |
| A/pelican/Nuble/233947-2/2023 H5N1 Chile 2023-01-26 | EPI_ISL_17885918 |
| A/pelican/Tarapaca/227436-2/2022 H5N1 Chile 2022-12-09 | EPI_ISL_17885942 |
| A/pelican/Antofagasta/228244-2/2022 H5N1 Chile 2022-12-16 | EPI_ISL_17885944 |
| A/chicken/Araucania/241914-1/2023 H5N1 Chile 2023-03-14 | EPI_ISL_17885945 |
| A/chicken/Araucania/241892-2/2023 H5N1 Chile 2023-03-14 | EPI_ISL_17885946 |
| A/humboldt penguin/Tarapaca/238744-2/2023 H5N1 Chile 2023-02-23 | EPI_ISL_17885947 |
| A/humboldt penguin/Coquimbo/239590/2023 H5N1 Chile 2023-02-28 | EPI_ISL_17885948 |
| A/chicken/Araucania/240481-1/2023 H5N1 Chile 2023-03-07 | EPI_ISL_17885949 |
| A/great egret/Araucania/240518/2023 H5N1 Chile 2023-03-07 | EPI_ISL_17885950 |
| A/gray gull/Tarapaca/232825-1/2023 H5N1 Chile 2023-01-18 | EPI_ISL_17885951 |
| A/pelican/Maule/231155-2/2023 H5N1 Chile 2023-01-09 | EPI_ISL_17885921 |
| A/pelican/Coquimbo/231946-1/2023 H5N1 Chile 2023-01-14 | EPI_ISL_17885922 |
| A/pelican/Coquimbo/230310-1/2023 H5N1 Chile 2023-01-02 | EPI_ISL_17885923 |
| A/pelican/Atacama/230158-2/2022 H5N1 Chile 2022-12-30 | EPI_ISL_17885924 |
| A/pelican/Atacama/230158-1/2022 H5N1 Chile 2022-12-30 | EPI_ISL_17885925 |
| A/pelican/Atacama/229450-2/2022 H5N1 Chile 2022-12-26 | EPI_ISL_17885926 |
| A/pelican/Atacama/229424-2/2022 H5N1 Chile 2022-12-22 | EPI_ISL_17885927 |
| A/pelican/Antofagasta/228318-1/2022 H5N1 Chile 2022-12-16 | EPI_ISL_17885928 |
| A/pelican/Antofagasta/228272-1/2022 H5N1 Chile 2022-12-18 | EPI_ISL_17885929 |
| A/pelican/Valparaiso/233091-1/2023 H5N1 Chile 2023-01-20 | EPI_ISL_17885930 |
| A/pelican/Antofagasta/228246-3/2022 H5N1 Chile 2022-12-15 | EPI_ISL_17885931 |
| A/pelican/Valparaiso/233418-1/2023 H5N1 Chile 2023-01-20 | EPI_ISL_17885932 |
| A/pelican/Antofagasta/228246-2/2022 H5N1 Chile 2022-12-15 | EPI_ISL_17885934 |
| A/south american sea lion/Peru/AQP-SER00R/2023 H5N1 Peru 2023-03-06 | EPI_ISL_18265422 |
| A/sea lion/Peru/AQP-SER00B/2023 H5N1 Peru 2023-01-25 | EPI_ISL_18054508 |



Appendix Figure 1. Location of Punta Bermeja and other HPAI H5N1 detections in pinnipeds since August 2023 (red dots) and HPAI H5N1 detections in poultry (blue dots) and wild birds (purple dots) in Argentina.

A


0.035
Note: Branches shorter than 0.0025 are shown as having length 0.0025

Country (Node color)

- Antarctica
- Argentina
- Brazil
- Chile
- Colombia
- Ecuador
- Other
- Peru
- Uruguay
- Venezuela

Host group (Node shape)

- Human
- Mammal
- Poultry
- Wild bird



B


0.020
Note: Branches shorter than 0.0013 are shown as having length 0.0013

Country (Node color)

- Antarctica
- Argentina
- Brazil
- Chile
- Colombia
- Ecuador
- Other
- Peru
- Uruguay
- Venezuela

Host group (Node shape)

- Human
- Mammal
- Poultry
- Wild bird



C


Note: Branches shorter than 0.0016 are shown as having length 0.0016

Country (Node color)

- Antarctica
- Argentina
- Brazil
- Chile
- Colombia
- Ecuador
- Other
- Peru
- Uruguay
- Venezuela

Host group (Node shape)

- Human
- Mammal
- Poultry
- Wild bird



D


Note: Branches shorter than C 0.009 are shown as having length 0.0009

Country (Node color)

- Antarctica
- Argentina
- Brazil
- Chile
- Colombia
- Ecuador
- Other
- Peru
- Uruguay
- Venezuela

Host group (Node shape)

- Human
- Mammal
- Poultry
- Wild bird



E

0.030


Note: Branches shorter than 0.0021 are shown as having length 0.0021

Country (Node color)

- Antarctica
- Argentina
- Brazil
- Chile
- Colombia
- Ecuador
- Other
- Peru
- Uruguay
- Venezuela

Host group (Node shape)

- Human
- Mammal
- Poultry
- Wild bird



F


0.009
Note: Branches shorter than 0.0005 are shown as having length 0.0005

Country (Node color)

- Antarctica
- Argentina
- Brazil
- Chile
- Colombia
- Ecuador
- Other
- Peru
- Uruguay
- Venezuela

Host group (Node shape)

- Human
- Mammal
- Poultry
- Wild bird



G


0.007
Note: Branches shorter than 0.0006 are shown as having length 0.0006

Country (Node color)

- Antarctica
- Argentina
- Brazil
- Chile
- Colombia
- Ecuador
- Other
- Peru
- Uruguay
- Venezuela



Host group (Node shape)

- Human
- Mammal
- Poultry
- Wild bird



H

Note: Branches shorter than 0.0558 are shown as having length 0.0058



Appendix Figure 2. Maximum likelihood tree for PB2, PB1, PA, HA, NP, NA, MP, and NS gene segments of HPAI H5N1 strains from Punta Bermeja, Argentina, in relation to all HPAI H5N1 strains from South America and a few strains from North America and Europe to show viral evolution in the southern hemisphere. Node shape represents host group, and node color (and bars adjacent to trees) represents the region/country. Branch lengths are drawn proportionally to the extent of changes (scale-bars are shown). Values adjacent to nodes represent bootstrap support.