

Data table for Figure 2. Age-adjusted death rates for selected causes of death for all ages, by sex: United States, 2006–2016

Excel and PowerPoint: https://www.cdc.gov/nchs/hus/contents2017.htm#Figure_002

Sex and year	All causes	Heart disease	Cancer	Cause of death ¹				
				Unintentional injuries ²	CLRD ³	Stroke ⁴	Diabetes ⁵	Alzheimer's disease
Male				Age-adjusted deaths per 100,000 population ⁶				
2006	943.5	254.9	221.7	56.0	48.4	45.2	27.7	19.4
2007	922.9	243.7	218.8	55.9	48.8	43.7	26.6	19.5
2008	918.8	238.5	214.9	54.3	52.3	42.2	25.9	21.3
2009	890.9	229.4	210.9	51.4	49.5	39.9	25.0	20.2
2010	887.1	225.1	209.9	51.5	48.7	39.3	24.9	21.0
2011	875.3	218.1	204.0	52.8	48.6	37.9	26.0	20.4
2012	865.1	214.7	200.3	52.6	47.2	37.1	25.5	19.8
2013	863.6	214.5	196.0	53.1	47.5	36.7	25.6	19.3
2014	855.1	210.9	192.9	54.6	45.4	36.9	25.6	20.6
2015	863.2	211.8	189.2	58.7	46.0	37.8	26.2	23.7
2016	861.0	209.1	185.4	65.0	45.1	37.5	26.0	24.3
Female				Age-adjusted deaths per 100,000 population ⁶				
2006	672.2	167.2	154.7	25.8	36.4	43.9	20.4	25.9
2007	658.1	159.0	152.3	26.1	36.6	42.7	19.8	26.2
2008	659.9	155.9	149.6	25.4	39.8	41.4	19.1	28.2
2009	636.8	146.6	147.4	24.8	38.3	38.8	17.9	26.3
2010	634.9	143.3	146.7	25.6	38.0	38.3	17.6	27.3
2011	632.4	138.7	144.0	26.5	38.5	37.2	18.2	27.1
2012	624.7	135.5	142.1	26.4	37.8	36.1	17.7	26.1
2013	623.5	134.3	139.5	26.6	38.5	35.2	17.6	25.9
2014	616.7	131.8	138.1	27.3	37.1	35.6	17.2	28.3
2015	624.2	133.6	135.9	28.7	38.6	36.9	17.3	32.8
2016	617.5	130.4	134.0	30.8	37.4	36.5	16.9	33.9
Male				Standard error				
2006	0.88	0.46	0.42	0.20	0.20	0.20	0.15	0.13
2007	0.86	0.45	0.41	0.20	0.20	0.19	0.14	0.13
2008	0.84	0.43	0.40	0.20	0.20	0.18	0.14	0.14
2009	0.82	0.42	0.39	0.19	0.20	0.18	0.14	0.13
2010	0.81	0.41	0.39	0.19	0.19	0.17	0.13	0.13
2011	0.79	0.40	0.38	0.19	0.19	0.17	0.14	0.13
2012	0.78	0.39	0.37	0.19	0.18	0.16	0.13	0.12
2013	0.77	0.38	0.36	0.19	0.18	0.16	0.13	0.12
2014	0.75	0.38	0.35	0.19	0.18	0.16	0.13	0.12
2015	0.75	0.37	0.34	0.20	0.17	0.16	0.13	0.13
2016	0.74	0.37	0.34	0.21	0.17	0.16	0.13	0.13
Female				Standard error				
2006	0.61	0.30	0.30	0.13	0.14	0.15	0.11	0.11
2007	0.60	0.29	0.30	0.13	0.14	0.15	0.11	0.11
2008	0.60	0.29	0.29	0.12	0.15	0.15	0.10	0.12
2009	0.59	0.27	0.29	0.12	0.14	0.14	0.10	0.11
2010	0.58	0.27	0.28	0.12	0.14	0.14	0.10	0.11
2011	0.57	0.26	0.28	0.12	0.14	0.14	0.10	0.11
2012	0.57	0.26	0.27	0.12	0.14	0.13	0.10	0.11
2013	0.56	0.25	0.27	0.12	0.14	0.13	0.09	0.11
2014	0.55	0.25	0.27	0.13	0.14	0.13	0.09	0.11
2015	0.55	0.25	0.26	0.13	0.14	0.13	0.09	0.12
2016	0.55	0.24	0.26	0.13	0.13	0.13	0.09	0.12

¹ Underlying causes of death are based on the *International Classification of Diseases, 10th Revision* (ICD-10).

² Unintentional injuries is accidents.

³ CLRD is chronic lower respiratory disease. Starting with 1999 data, the rules for selecting CLRD and Pneumonia as the underlying cause of death changed, resulting in an increase in the number of deaths for CLRD and a decrease in the number of deaths for pneumonia. Therefore, trend data for these two causes of death should be interpreted with caution. For more information, see Comparability of cause of death between ICD-9 and ICD-10 in [Appendix II](#), Table V.

⁴ Stroke is cerebrovascular disease.

⁵ Starting with 2011 data, the rules for selecting Renal failure as the underlying cause of death were changed, resulting in an increase in the number of deaths for Diabetes mellitus. Therefore, trend data for diabetes should be interpreted with caution. For more information, see Technical Notes in Deaths: Final data for 2011, available from: https://www.cdc.gov/nchs/data/nvsr/nvsr63/nvsr63_03.pdf.

⁶ Estimates are age-adjusted to the year 2000 standard population with unrounded population numbers. See [Appendix II](#), Age adjustment.

NOTES: See [Appendix II](#), Cause of death; Cause-of-death ranking; Table IV.

SOURCE: NCHS, National Vital Statistics System, Mortality. See [Appendix I](#), National Vital Statistics System (NVSS).