## Data table for Figure 5. Cigarette smoking among adults aged 18 and over, by age, and tobacco use among adolescents in grades 9–12, by age and type of product: United States, 2006–2016

Excel and PowerPoint: https://www.cdc.gov/nchs/hus/contents2017.htm#Figure\_005

Current cigarette smoking among adults aged 18 and over, by age: 2006-2016

Year	18–44 years		45–64 years		65 years and over	
	Percent	SE	Percent	SE	Percent	SE
2006	23.6	0.5	21.8	0.6	10.2	0.5
2007	22.6	0.6	21.0	0.6	8.3	0.5
2008	23.1	0.6	22.6	0.6	9.3	0.5
2009	23.4	0.5	21.9	0.6	9.5	0.5
2010	21.5	0.5	21.1	0.5	9.5	0.5
2011	21.2	0.5	21.4	0.5	7.9	0.4
2012	20.4	0.5	19.5	0.5	8.9	0.4
2013	19.7	0.5	19.9	0.5	8.8	0.4
2014	19.1	0.5	18.0	0.5	8.5	0.4
2015	16.5	0.5	17.0	0.5	8.4	0.4
2016	16.4	0.5	18.0	0.5	8.8	0.4

## Tobacco use in past 30 days among students in grades 9-12, by type of product: 2011-2016

Year	Any tobacco products	Cigarettes	Cigars	Electronic cigarettes	Hookah	Smokeless tobacco	Pipe tobacco	Bidis
				Perc	ent			
2011	24.2	15.8	11.6	1.5	4.1	7.9	4.0	2.0
2012	23.3	14.0	12.6	2.8	5.4	7.3	4.5	0.9
2013	22.9	12.7	11.9	4.5	5.2	6.2	4.1	0.6
2014	24.6	9.2	8.2	13.4	9.4	6.3	1.5	0.9
2015	25.3	9.3	8.6	16.0	7.2	6.0	1.0	0.6
2016	20.2	8.0	7.7	11.3	4.8	5.8	1.4	0.5
	Standard error							
2011	1.2	1.1	0.6	0.2	0.4	0.8	0.3	0.2
2012	0.9	8.0	0.6	0.3	0.4	0.6	0.3	0.1
013	0.9	0.7	0.6	0.4	0.4	0.7	0.3	0.1
014	1.0	0.6	0.5	1.2	0.6	0.6	0.2	0.1
015	1.1	0.8	0.5	1.0	0.5	0.7	0.2	0.1
016	1.0	0.7	0.6	0.8	0.4	0.6	0.1	0.1

SE is standard error.

NOTES: Current cigarette smoking by adults is defined as having smoked 100 or more cigarettes in their lifetime and smoking now, every day or some days. Tobacco use by students in grades 9–12 is defined as having used the product one or more times in the past 30 days. Electronic cigarettes are cigarette shaped devices containing nicotine and other chemicals, which is vaporized and inhaled to simulate smoking. Bidis are small, thin, hand-rolled cigarettes imported to the United States. See Appendix II, Cigarette smoking. Data are for the civilian noninstitutionalized population.

SOURCE: NCHS, National Health Interview Survey, and CDC, National Youth Tobacco Survey. See Appendix I, National Health Interview Survey (NHIS) and National Youth Tobacco Survey (NYTS).