

NATIONAL OCCUPATIONAL RESEARCH AGENDA (NORA)

NATIONAL OCCUPATIONAL RESEARCH AGENDA FOR HEALTHY WORK DESIGN AND WELL-BEING

January 2020

Developed by the NORA Healthy Work Design and Well-Being (HWD) Cross-Sector Council

For more information about the National Occupational Research Agenda (NORA), visit the web site: https://www.cdc.gov/nora/

For monthly updates on NORA, subscribe to NIOSH eNews at <u>www.cdc.gov/niosh/eNews</u>

Disclaimer

This is a product of the National Occupational Research Agenda (NORA) Healthy Work Design and Well-Being Cross-Sector Council. It does not necessarily represent the official position of the National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention, or U.S. Department of Health and Human Services.

INTRODUCTION

What is the National Occupational Research Agenda?

The National Occupational Research Agenda (NORA) is a partnership program to stimulate innovative research and workplace interventions. In combination with other initiatives, the products of this program are expected to reduce the occurrence of injuries and illnesses at work. Unveiled in 1996, NORA has become a research framework for the Nation and National Institute for Occupational Safety and Health (NIOSH). Diverse parties collaborate to identify the most critical issues in workplace safety and health and develop research objectives for addressing those needs.

NORA entered its third decade in 2016 with an enhanced structure. The ten sectors formed for the second decade continue to prioritize occupational safety and health research by major areas of the U.S. economy. In addition, there are seven cross-sectors organized according the major health and safety issues affecting the U.S. working population. While NIOSH is serving as the steward to move this effort forward, it is truly a national effort. NORA is carried out through multi-stakeholder councils, which are developing and implementing research agendas for the occupational safety and health community over the decade (2016-2026). Councils address objectives through information exchange, partnership building, and enhanced dissemination and implementation of evidence-based solutions.

NORA groups health and safety issues into seven cross-sectors. The Healthy Work Design and Well-Being (HWD) Cross-Sector focuses on protecting and advancing worker safety, health, and well-being by improving the design of work, management practices, and the physical and psychosocial work environment. HWD's holistic perspective focuses on how work affects overall safety, health, and well-being, including physical, psychological, social, and economic aspects.

What are NORA Councils?

Participation in NORA councils is broad, including stakeholders from universities, large and small businesses, professional societies, government agencies, and worker organizations. Councils are co-chaired by one NIOSH representative and another member from outside NIOSH.

Statement of Purpose

NORA councils are a national venue for individuals and organizations with common interests in occupational safety and health topics to come together. Councils will start the third decade by identifying broad occupational safety and health research objectives for the nation. These research objectives will build from advances in knowledge in the last decade, address emerging issues, and be based on council member and public input. Councils will spend the remainder of the decade working together to address the agenda through information exchange, collaboration, and enhanced dissemination and implementation of solutions that work.

Although NIOSH is the steward of NORA, it is just one of many partners that make NORA possible. Councils are not an opportunity to give consensus advice to NIOSH, but instead a way to maximize resources towards improved occupational safety and health nationwide. Councils are platforms that help build close partnerships among members and broader collaborations between councils and other organizations. The resulting information sharing and leveraging efforts promote widespread adoption of improved workplace practices based on research results.

Councils are diverse and dynamic and are open to anyone with an interest in occupational safety and health. Members benefit by hearing about cutting-edge research findings, learning about evidence-based ways to improve safety and health efforts in their organization, and forming new partnerships. In turn, members share their knowledge and experiences with others and reciprocate partnerships.

NORA Healthy Work Design and Well-Being (HWD) Council

The NORA HWD Cross-Sector Council was formed in early 2017. The Council aims to provide national leadership in the area of HWD through gathering and synthesizing scientific information, creating knowledge, providing recommendations, and delivering products to those who are developing and implementing workplace prevention efforts. The Council intends for the National Occupational Research Agenda for HWD to succinctly summarize research objectives that will address important gaps and needs, and advance knowledge and practice. The members of the NORA HWD Cross-Sector Council are:

David DeJoy, PhD, University of Georgia, Athens,	Russell Matthews, PhD, University of Alabama
Co-Chair	Patricia McGovern, PhD, MPH, University of
Rene Pana-Cryan, PhD, NIOSH, Co-Chair	Minnesota
L. Casey Chosewood, MD, MPH, NIOSH, Co-Chair	Lee Newman, MD, MA, University of Colorado Denver
Naomi Swanson, PhD, NIOSH, Co-Chair	
David Ballard, PsyD, MBA, American Psychological Association	Jeannie Nigam, MS, NIOSH
	Brian Quay, PhD, NIOSH
Les Boden, PhD, Boston University	Tapas Ray, PhD, NIOSH
Carol Brown, PhD, University of Colorado	Bonnie Rogers, DrPH, COHN-S, FAAN, University of North Carolina at Chapel Hill
Tim Bushnell, PhD, NIOSH	
Claire C. Caruso, PhD, RN, FAAN, NIOSH	Steve Sauter, PhD, ConsultantLaura Stock, MPH, University of California, Berkeley
BongKyoo Choi, ScD, MPH, University of California, Irvine	Ron Stout, MD, Proctor & Gamble
Lorraine M. Conroy, ScD, University of Illinois at Chicago	Jessica Streit, PhD, NIOSH
	Jennifer Swanberg, PhD, MMHS, OTR, University of
Michael Foley, MA, Washington State Department of Labor and Industries	Maryland, Baltimore
	Sara Tamers, PhD, MPH, NIOSH
Leslie Hammer, PhD, Portland State University and Oregon Health & Science University	Ibraheem (Abe) Tarawneh, PhD, Ohio Center for Occupational Safety and Health
Karen Heaton, PhD, CRNP, FNP-BC, FAAN, University of Alabama at Birmingham	Carolyn Winslow, PhD, University of California, Berkeley
Heidi Hudson, MPH, NIOSH	Past Participants:
Pamela Hymel, MD, Walt Disney World Parks and Resorts	Lisa Brosseau, ScD, University of Illinois at Chicago
Dave LeGrande, MA, RN, Communication Workers of America	
David Lombardi, PhD, Harvard University	

What does the National Occupational Research Agenda for HWD represent?

The National Occupational Research Agenda for HWD is intended to identify the knowledge and actions most urgently needed to identify occupational risk factors to prevent avoidable adverse health outcomes among workers. This Agenda provides a vehicle for stakeholders to describe the most relevant issues, research gaps, and safety and health needs for the cross-sector. It is meant to be broader than any one agency or organization. It identifies the priorities for the entire country and all of its research and development entities, whether government, higher education, or industry. Because the Agenda is intended to guide national occupational health and safety efforts for HWD, it cannot at the same time be an *inventory* of all issues worthy of attention. The omission of a topic does not mean that topic was viewed as unimportant. Those who developed this Agenda did, however, believe that the number of topics should be small enough so that resources could be focused on a manageable set of objectives, thereby increasing the likelihood of real impact in the workplace.

NIOSH used the draft Agendas created by the sector and cross-sector NORA councils as an input into the development of a <u>NIOSH Strategic Plan</u>. Programs used the <u>burden</u>, <u>need and impact method</u> to write research goals that articulate and operationalize the components of the NORA sector and cross-sector Agendas that NIOSH will take up. NORA Agendas and the NIOSH Strategic Plan are separate but linked.

Who are the target audiences?

The National Occupational Research Agenda for HWD was developed to provide information about the most relevant HWD research gaps and needs and is targeted to researchers, health and safety professionals, practitioners, and human resources professionals in industry, labor, academia, and government. The objectives are meant to assist in setting priorities around occupational health and safety research and produce information that can help inform workplace practices.

How was the research agenda developed?

This Agenda was developed through a series of conference calls and email communications. A draft list of relevant topics was drawn up, then refined and grouped into seven broad objectives by the HWD Council. Working subgroups were formed for each of the seven objectives. A variety of information sources were used, including articles from the scientific literature, and HWD Council member expertise and experience. HWD Council members went through three iterations of review and discussion and redrafting the objectives in the development of the Agenda. Although there is some overlap in the risk factors and outcomes across the objectives, each has a unique primary focus.

THE OBJECTIVES

The Healthy Work Design and Well-Being (HWD Cross-Sector focuses on protecting and advancing worker safety, health, and well-being by improving the design of work, management practices, and the physical and psychosocial work environment. HWD's holistic perspective focuses on how work affects overall safety, health and well-being, including physical, psychological, social, and economic aspects. The following seven research objectives address the most critical HWD issues identified by the HWD Council. Interactions among research strategies aiming to address individual objectives are encouraged even when not explicitly noted, in order to identify and address solutions at the intersections of the objectives.

Objective 1: Identify and examine the impact of worker demographics on employer or organizational practices and worker safety, health, and well-being

Background

Demographic characteristics, including age, race, ethnicity, country of origin, language, and disability may adversely and disproportionately affect worker safety, health, and well-being. Workers in certain demographic groups often face elevated safety and health risks due to disparities in employment opportunities; management practices and treatment; wages; and exposure to physically demanding, labor-intensive jobs; and unsafe workplace environments. For example, workers with certain demographic characteristics are often more concentrated in work arrangements that are temporary in nature (Ray, Kenigsberg, & Pana-Cryan, 2017) and where exposure to physical and psychosocial risks is more pronounced (Howard, 2017; Benavides et al., 2006). (See Objective 2 for more information on work arrangements.) In addition, many of these jobs do not provide employee benefits such as health insurance, paid leave, and retirement, thereby affecting overall worker wellbeing. In March of 2017, while 93% of managers had paid sick leave benefits, only 46% of service workers and 47% of construction, extraction, farming, fishing, and forestry workers had paid sick leave benefits. Less than half (48%) of private industry establishments offered retirement benefits, and a little over a half (58%) of these establishments offered healthcare benefits to their workers. (BLS, 2018a; BLS, 2018b).

Workers in certain demographic groups are more likely to earn low wages and there are wage disparities even among low wage workers. For example, a recent study (Vincent, Fusaro, & Shafer, 2016) that categorized workers in three low wage categories (lowest if paid up to \$9.25 per hour, highest if paid \$10.75 to \$13.50 per hour) found that in 2013, the majority of workers in the middle and highest categories were White non-Hispanic. Both Hispanic and Black workers were less prevalent in the highest wage category than in the lowest category.

In addition, there is an ever-widening recognition of work-related hazards, including low wages, which in turn seem to result in poor health or health behaviors (Leigh & De Vogli, 2016).

Age

Older workers

The Bureau of Labor Statistics (BLS) projects that by 2024, approximately 25% of the American workforce (about 41 million workers) will be aged 55 or older, with workers aged 65 or older expected to be the fastest growing segment (Toossi & Torpey, 2017). While older workers are often psychologically healthier, report more positive psychosocial work environments (Sauter, Keel, & Hanseman, 2009), and have fewer occupational injuries overall than younger workers, the rate of fatalities due to occupational injuries suffered by workers aged 65 and older is nearly three times that of the overall workforce (BLS, 2017). In addition, older workers suffer from more chronic health conditions that can affect their ability to stay safe and healthy at work. Maintaining productivity in an aging workforce has important economic implications. The Congressional Budget Office has projected that by 2030, aggregate expenditures on federal programs that include social security, Medicare, and Medicaid will likely be

16% of the total output of the economy (CBO, 2001). These challenges demonstrate a need to develop and implement age-friendly work design, management practices, and physical and psychosocial work environments to create workplaces that are safe for all workers as they age (Silverstein, 2008).

Younger workers

In 2016, there were about 19 million workers aged 16 to 24, representing about 13% of the total workforce. In 2016, over 380 workers under the age of 24 died from occupational injuries (BLS, 2017). In 2014, the rate of occupational injuries treated in emergency departments for workers aged 15–19 was 2.18 times greater than the rate for workers aged 25 and older. The rate for workers aged 20–24 was 1.76 times greater than for workers aged 25 and older (NIOSH, 2017). Injuries to young workers are often the result of working in high-risk jobs, lack of work experience and safety training, and fear of speaking up about the workplace hazards that they and newly hired workers face.

Race and ethnicity

The American workforce continues to become more diverse. In 1996, White non-Hispanic workers comprised about 75% of the workforce. In 2016, White non-Hispanic workers comprised 63% of the workforce, and are predicted to be about 58% of the workforce by 2026 (BLS, 2017). By 2026, Hispanics, non-Hispanic Blacks, and non-Hispanic Asians are expected to constitute about 21%, 13%, and 7% of the workforce, respectively (BLS, 2017).

A recent study used national survey data to test for differences between members of minority groups and non-Hispanic white workers in the risk of workplace injuries and the prevalence of work-related disabilities. The researcher found that, after adjusting for covariates, disparities in economic opportunities expose members of minority groups to increased risk of workplace injury and disability (Seabury, Terp, & Boden, 2017).

There also is evidence of adverse health effects attributed to racial and ethnic discrimination. Shannon et al. (2009) found that experiences of racial discrimination at work were associated with risk for work-related illness, injury, or assault across two time points. The odds of reporting racial discrimination at work were between 3 and 8 times greater among minorities compared with Whites; those who experienced discrimination at work were two times as likely as those without discrimination experiences to report illness/injury/assault; and, the odds of illness/injury/assault were double among Hispanics compared with White respondents (Blacks and those from 'other races' did not differ significantly from Whites).

Country of origin and language

In 2016, there were 27 million foreign-born workers in the U.S., comprising about 17% of the total labor force (BLS, 2017). Foreign-born workers may face workplace discrimination that results in an increased exposure to adverse working conditions. Non-English-speaking workers may lack access to safety and health training in their own language and may lack familiarity with programs that protect workers (Byler, 2013). In addition, concerns about immigration status may result in fear of speaking up and of exercising their right to a safe and healthy workplace. These disparities have resulted in high rates of injuries, illnesses, and fatalities among foreign-born workers (Byler, 2013).

- Enable better sharing of data among state and national agencies to improve the understanding about the nature, source, and consequences of injuries and illnesses among older and younger workers.
- Gather and disseminate best practices for age- and disability-friendly workplace policies, including job redesign interventions, flexible scheduling models, and anti-discrimination policies.
- Conduct intervention research on strategies including youth leadership training programs, community awareness campaigns, and school based and job training programs to promote awareness among young

workers about occupational safety and health and development of necessary skills to advocate for improved working conditions, and active participation in efforts to address occupational hazards.

- Conduct research on the impact of racial and ethnic discrimination and disparities in the workplace and its effect on worker safety, health, and well-being.
- Conduct research on the impact of wages and hours on workers' health status.
- Conduct research into the extent of underreporting of injuries and illnesses by foreign-born and other vulnerable workers and by their employers, and identify best practices in data collection strategies and systems.
- Collect and disseminate best practices for addressing the occupational safety and health of foreign-born workers, including policies that prevent workplace discrimination, promote worker involvement in identifying and addressing hazards without fear of retaliation, and education and training efforts that reflect the specific language and literacy needs of diverse workers.
- Conduct intersectional research that examines the effects of all aspects of worker demographics on worker safety, health, and well-being.

Objective 2: Improve the Safety, Health, and Well-Being of Workers with Non-Standard Work Arrangements

Background

There has been an upward trend in the share of the U.S. workforce in non-standard work arrangements during the 2000s (Katz & Krueger, 2019). According to definitions and data used in recent studies by Katz and Krueger (2016, 2019), there was a 1-2 percentage point increase in the share of workers in alternative work from 2005 to 2015. The authors acknowledged it was difficult to compare trends in work arrangements across surveys with different sampling frames and at different points of the business cycle, as well as that work status seems to not be accurately reported, especially during a period of changing work relationships.

In addition, details on work arrangements are commonly collected for a worker's main job. This leads to a poor understanding of the complexities associated with holding and managing multiple jobs, simultaneously. One commonality shared by many workers in non-standard work arrangements is that there is no expectation of permanence (Howard, 2017). Many non-standard work arrangements are characterized by temporariness, instability, unpredictability of work hours, limited schedule control, and lack of legal protections and employer sponsored benefits for workers (Bushnell et al., 2017). All of these factors can influence worker safety, health, and well-being.

For example, unpredictable work schedules can be associated with reduced worker well-being (Henly & Lambert, 2014). Non-standard work arrangements may also be associated with increased chronic opioid use. Pensa et al. (2018) found that being paid hourly was a predictor for increased chronic opioid prescribing. Although there is research (Vives et al., 2011) and systematic reviews (e.g., Quinlan, 2015) on the consequences of non-standard work arrangements, and despite the rise in the prevalence of non-standard work arrangements, these types of arrangements continue to be understudied, and their determinants and safety, health, and well-being consequences continue to be poorly understood.

This is partially due to the current occupational safety and health surveillance systems that do not adequately capture information about both non-standard work arrangements and associated adverse health outcomes. In addition, traditional workplace-based interventions to improve safety, health, and well-being are likely to be

inadequate for addressing any harmful characteristics of non-standard work. Likewise, traditional intervention dissemination models may be inadequate for improving the safety, health, and well-being of workers in certain non-standard work arrangements.

Categories of non-standard work arrangements

Although there is currently no consistent definition for non-standard work arrangements among surveys that aim to collect these data, workers in these arrangements can be grouped into several categories: (1) contingent workers, hired for a specific project or other short-term need; (2) subcontracted workers, where working conditions are controlled by an entity other than the workers' employer (for example, temporary agency workers); (3) independent contractors, who are self-employed and work at a location controlled by a business with which they have a contractual relationship; (4) "gig" workers, who perform work intermediated by a digital on-line platform; (5) part-time workers, many of whom are in low-wage hourly jobs, have unpredictable work hours and schedules, and would prefer to work full-time; and (6) workers assigned to schedules other than the traditional 8:00 am to 5:00 pm, Monday through Friday routine (Howard, 2017; Swanberg, Watson, & Eastman, 2014), including unpredictable schedules (Henly & Lambert, 2014).

Temporary and precarious employment

Compared with standard work arrangements, several non-standard arrangements involve temporary jobs and insecurity of employment and income, which have been linked to low levels of health and well-being (Virtanen, et al., 2005). Insufficient household income and non-standard or irregular hours are associated with poor mental health (De Moortel, Vandenheede, & Vanroelen, 2014). Similarly to newly hired workers, temporary workers frequently lack safety and other job training, and experience higher injury rates than permanently employed workers (Foley, 2017). (See Objective 1, younger workers.) A recent study that used data from years between 2002 and 2014 demonstrated that work arrangement was an important predictor of job stress. Compared with non-stressed workers, stressed workers across all work arrangements reported more days of poor physical and mental health and more days with activity limitations (Ray et al., 2017). (See Objective 4 and 6 for more information on the adverse effects of job stress.)

- Improve occupational health surveillance systems' data collection on work arrangements, including information on arrangements in multiple jobs held simultaneously, and associated adverse health outcomes.
- Develop surveillance methods, including those used for community-level surveillance systems that measure the prevalence and characteristics of non-standard work and identify workers and workplaces with non-standard arrangements.
- Describe the characteristics of non-standard work that are detrimental to worker health and the mechanisms of these effects in different economic sectors and for various demographic groups.
- Understand specific socioeconomic and other risk factors, such as job insecurity, that affect the prevalence of precarious work and jobs with non-standard work arrangements; this, considered together with information on the gap identified immediately above, can increase awareness and point to prevention guidance for employers, organizations, and policy makers.
- Identify and evaluate interventions that improve the safety, health, and well-being of workers in nonstandard work arrangements.
- Disseminate effective interventions that improve the safety, health, and well-being of workers in nonstandard work arrangements.

 Collect and disseminate best practices for addressing the occupational safety and health of workers in non-standard work arrangements, such as community based participatory research and policies that prevent workplace discrimination, promote worker involvement in identifying and addressing hazards without fear of retaliation, and education and training efforts that reflect the specific needs of these workers.

Objective 3: Address the Safety and Health Implications of Advancing Technology

Background

Computers have become a central component of many jobs, and there are rapidly expanding applications of robotics and sophisticated information and communication technologies affecting most industries and occupations. Continuing developments in artificial intelligence and machine learning are driving much of this expansion. Recently, an expert panel concluded that "advances in information technology are far from over, and some of the biggest improvements in areas like artificial intelligence are likely still to come" (National Academies of Sciences, Engineering, and Medicine, 2017). The full impact of changes in technology follows from both the development of new technologies and the diffusion and maturation of existing technologies. For most of the topics discussed below, there has been only scattered direct research. There is a clear need to develop surveillance methods, prioritize research, and create and disseminate strategies for safeguarding workers during a time of rapid technological dynamism.

New technological demands on workers

New technologies have the potential to make jobs safer and easier to perform but can also produce the opposite effect: jobs can become more complex and workers can be confronted with new and unanticipated hazards. Technology can also result in the creation of entirely new jobs and the elimination of other jobs. A recent study indicated that as many as 45% of the activities people are paid to perform could be automated by taking advantage of currently available technologies (Chui, Manyika, & Miremadi, 2016). The introduction of new technologies may require changes in business processes, job definitions, and worker adaptation. Concerns involving data security and related issues are also likely to become more complex. Workers must learn how to use the new technology and how best to incorporate it into their workflow. The term "technostress" has been coined to refer to the demands placed on workers by the incorporation of new technologies into the workplace (Tarafdar et al., 2015).

New technologies may force workers to use multiple streams of information from different devices and sources, leading to increased mental workload. Existing systems and components may be updated or otherwise modified with little warning and inadequate training or technical support. In some cases, technologies can make work tasks monotonous or less engaging through oversimplification. Tasks that once involved specific actions and direct involvement in production processes may now mostly involve passive monitoring of computer screens or gauges. Technology can also be used to monitor employee performance, which may be stressful for workers. Thus far, technology has been viewed as primarily impacting low-skill jobs and less educated workers. Future impacts, however, are likely to be much broader as increasingly advanced technologies are introduced into skilled professions such as medicine, banking, and law. For highly skilled workers, technology is more likely to complement their jobs than eliminate them. Still, the incorporation of these changes is likely to be disruptive and stressful at least in the short term.

Human-machine interaction

Workers have long interacted with technology of one kind or another. Currently, however, these interactions are changing in fundamental ways. Early robotic systems performed basic tasks with high mechanical precision and

initial safety concerns associated with industrial robots focused mostly on preventing workers from being struck by the moving robotic arm or being trapped within its range of movement. Many of today's robots are programmable and utilize various types of sensors. Some can learn a new task from observation of humans. Progress is also being made in computer perception of speech, vision, and sound (National Academies of Sciences, Engineering, and Medicine, 2017). In the future, workers will interact with and communicate with robots in a variety of different ways. Workers will supervise and support robots at the same time that they are interacting with other people (Hancock, 2009; Rus, 2015) and robots will also communicate and interact with each other. These future types of interactions will raise new and more complex issues in protecting workers who interact with robotic devices.

Virtual work, re-defining the "workplace"

Technology can clearly impact how people work; it can also impact where they work, when they work, and even how much they work. Today, workers in many jobs can telecommute or work remotely, and interact with colleagues in multiple locations around the world. The wide availability of information and communication technologies have blurred the separation between work-life and home-life. A 2015 Pew Research Center report (Purcell & Rainie, 2015) indicated that 35% of employed adults who use the internet at work reported that the internet, email, and cell phones have increased the amount of time they spend working. In essence, workers can be continuously connected to their work. Technological change can also alter the fundamental contract between employer and employee. Software algorithms are increasingly being used to assign, optimize, and evaluate employee performance in a variety of traditional as well as new or emerging employment relationships (e.g., gig employment) (Lee et al., 2015). Non-standard work arrangements such as independent contracting and working for temporary employment agencies have been expanding for some time (Katz & Krueger, 2016). Information and communication technologies have helped fuel this trend by making it easier for employers and potential workers to connect, communicate, and complete a variety of tasks, remotely. This has led to the creation of new companies that make use of technology-mediated on-demand or "gig" employment (U.S. Census Bureau, 2015). While these non-standard work arrangements have led to increased employment insecurity for some, they have provided others with increased flexibility and expanded employment opportunities. (See Objective 2 for more information on non-standard work arrangements and Objective 7 for more information on workplace flexibility policies and practices.)

- Develop a taxonomy of types of work-related technological changes and the types and extent of worker interaction with them.
- Conduct surveillance on the use of different types of technology by industry and occupation over time and the relationship with worker safety, health, and productivity outcomes.
- Conduct systematic, multidisciplinary research on the impact on workers of different types of technology, including through changes in job design, work schedules, training, demands, and safety.
- Conduct research on how to best design technology, including research on how humans interact with technology, to enhance worker safety, health, and productivity.
- Develop best practices for supporting worker safety and health in the introduction of new work-related technologies.

Objective 4: Reduce Work Organization-Related Chronic Health Conditions among Workers

Background

Although more than a century's worth of research documents the link between work organization, job design, and acute and chronic worker health outcomes, research on these linkages has intensified in recent years (Sauter & Hurrell, 2017; Vinchur & Koppes, 2011). Management and human resources policies, leadership and management practices, organizational climate and culture, and economic conditions can have profound implications for the health of workers. Many chronic health conditions, both physical and psychological, have been shown to be associated with job design, work organization and associated job stress. For example, occupations in the public sector, such as education and social services, often face increasing demands for services with limited support, decision-making authority, and job security contributing to job stress and the risk of chronic conditions such as cardiovascular disease and violence-related injuries (Landsbergis et al, 2017). Lean production practices can create a climate of intensified work pace and high demands—also often associated with low decision latitude—thus contributing to job strain and subsequent health effects (Landsbergis et al, 1999).

Working conditions and other elements of work organization and employment play a critical role in the Nation's opioid overdose epidemic among workers. Preventing opioid and other substance use disorders by improving surveillance and expanding research, which explores the connections between work, work injury, working conditions, and employment factors and antecedents or risk factors for drug misuse, is vital to stemming the tide of deaths from opioids and other substances. While work-related factors can be antecedents to opioid use disorder, workplaces can be helpful platforms for awareness building; decreasing stigma; coverage of, screening and referral for treatment; and reduction of secondary harms. Workplace policies can expand access to high-quality treatment programs for substance use disorder and recovery-friendly workplaces can support return to and sustained employment, improving opportunities for individuals and communities.

Given the enormous health and economic burden of chronic health conditions for both individuals and society as a whole, there is a need to understand the links between job design, working conditions, work organization, and these chronic health conditions.¹ (Objective 6 in this document addresses additional opportunities to improve the safety, health and well-being of workers through improved job design and organizational practices.)

Obesity

Obesity is a complex and costly condition which affected approximately 36% of adults in the U.S. during 2011–2014. (Ogden et al., 2015). Being overweight or obese is linked to an increased risk of developing a variety of serious health problems, including hypertension, type 2 diabetes, metabolic syndrome, stroke, coronary heart disease, sleep apnea, osteoarthritis, and some cancers (National Center for Chronic Disease Prevention and Health Promotion, 2016). Obesity and its associated health problems have had a significant economic impact on both health care costs and work-related productivity (Cawley & Meyerhoefer, 2012). Using 2006 and 2008 data, annual costs attributable to obesity among full-time employees in the U.S. have been estimated to exceed 70 billion dollars (Finkelstein et al., 2010). Overweight and obesity also have important implications in the work environment and lead to a number of significant negative impacts on workers as well as employers. Overweight and obese workers, on average, earn lower salaries, are viewed more negatively than others of normal weight, both by supervisors and coworkers, and can be victims of discrimination in the workplace (Judge & Cable, 2011).

¹ Musculoskeletal health conditions related to the design of work and working conditions were not included in Objective 4 as they are covered in the <u>National Occupational Research Agenda for Musculoskeletal Health</u>.

In one study among manufacturing firms, psychosocial factors associated with higher BMI were found to vary by age with higher overall stress reported among those ages 45-55 years of age (Garza et al, 2015). When obesity is present, companies incur costs linked to absenteeism and both workers and employers have higher medical expenditures (Finkelstein, Fiebelkorn & Wang, 2005). Additionally, workers with obesity experience lower productivity at work due to health problems and musculoskeletal injuries (Gu, et al, 2016) along with "presenteeism" (Ricci & Chee, 2005) and higher rates of obesity-related disability (Sturm, Ringel, & Andreyeva, 2004).

Cardiovascular disease

Cardiovascular disease (CVD), including heart and cerebrovascular disease, is the leading cause of death in the U.S. (Heron, 2016). CVD is prevalent among the working population; 53% of those with CVD are less than 60 years of age, according to pre-2000 data (Cooper et al., 2000; Leigh & Miller, 1998). CVD is a leading cause of death and permanent disability among workers, resulting in an average loss of seven years of life expectancy (American Heart Association, 2000). While many workplace prevention efforts for CVD have primarily focused upon altering individual health behaviors (e.g., smoking, physical activity, and nutritional practices), a growing body of research has identified the need to focus on the association between CVD and work organization factors, including psychosocial risks common in many workplaces (Theorell et al., 2016). The Sixth International Conference on Work Environment and Cardiovascular Disease adopted a statement in 2013 estimating that "10%-20% of all causes of CVD deaths among working age populations…are work-related" (International Commission on Occupational Health, 2013, page 4).

Depression and anxiety

In 2016, there were an estimated 44.7 million adults aged 18 or older in the U.S. (about 18% of the adult population) with any mental illness (AMI) within the past year (National Institute of Mental Health, 2016). AMI can vary in impact, ranging from no impairment to mild, moderate, or severe impairment. Moreover, nearly one in four of those with AMI (approximately 10.4 million Americans) have a serious mental illness, which results in significant functional impairment. Depression, anxiety, and other mental health conditions are costly in both human and economic terms. In 2015, more than 16 million Americans (or nearly 7% of all adult Americans) had a major depressive episode within the past year (National Institute of Mental Health, 2015). In 2003, depression was estimated to be responsible for \$36.6 billion to \$51.5 billion in lost productivity in American workplaces (Stewart, Ricci, Chee, et al., 2003). It is also widely believed that mental health conditions interact with many other chronic health conditions, such as CVD, obesity, and respiratory disease, increasing the severity, costs, and risk for disability from these chronic health conditions (Huffman et al., 2013; Wells et al., 1989). Fast-paced work, high job demands, job insecurity, and low wages in many work settings may increase risks for poor mental health outcomes. (See Objective 6.) Depression symptoms are also strongly related to a number of productivity outcomes such as impaired work performance and work absences (National Institute for Occupational Safety and Health, 1999). A 2012 analysis found that depression topped the list of risk factors linked to increased employee and employer health-care spending (Goetzel et al., 2012).

Cancer

In 2018, the International Agency for Research on Cancer estimated that there were 18.1 million new cancer cases and 9.5 million cancer deaths around the globe (International Agency for Research on Cancer, 2018. In the U.S., cancer remains a leading cause of death. Many factors play a role in the development of cancer including personal characteristics, family history, diet and personal habits, exposures to cancer-causing agents in the environment, and a variety of exposures related to work. The link between cancer and work exposures is well-documented. Up to 14% of cancer deaths in men may be related to work (Purdue et al., 2015). Up to 20% of lung cancer deaths and 7% of bladder cancer deaths were attributed to occupational exposures in Great Britain in 2004-2005 (Rushton et al., 2012). Research on work organization and job design factors and cancer is expanding. Examples include findings related to circadian disruption and shift work, job stress, long hours of work, missed health screenings and lack of or limitations in healthcare benefits related to work arrangement or employment status (Costa, House, & Stevens, 2010; Straif et al., 2007). (See Objectives 2, 5, and 6.)

Substance misuse and the opioid overdose crisis

Substance misuse among the working population is a serious and growing problem. The recent acuity and severity of a crisis related to opioid misuse and overdose deaths from opioids have brought renewed attention to this issue. Overdose deaths from prescription and non-medical opioid use doubled from 2010 to 2016, with more than 43,000 deaths in 2016 (CDC 2016). Of the 70,237 drug overdose deaths in 2017, nearly 68% (47,600) involved an opioid (Scholl et al., 2017). The rate of overdose deaths continues to increase. In response, the President declared the opioid epidemic to be a public health emergency in October 2017 (The White House, 2017). The Bureau of Labor Statistics (BLS) reported that overdose deaths at work from non-medical use of drugs or alcohol increased by at least 38% annually between 2013 and 2016. The 217 workplace overdose deaths reported in 2016 accounted for 4.2% of occupational injury deaths that year, compared to 1.8% in 2013 (BLS 2017). This large increase in overdose deaths in the workplace (from all drugs) parallels a surge in overall overdose deaths from opioids reported by CDC (2017).

Opioids are often initially prescribed to manage pain arising from a work injury. Risky workplace conditions that lead to injury, such as slip, trip, and fall hazards or heavy physical workloads, can be associated with prescription opioid use (Kowalski-McGraw et al., 2017). Other factors, such as job insecurity, job loss, and high demand/low control jobs may also be associated with prescription opioid use (Kowalski-McGraw et al., 2017). Some people who are prescribed opioids may misuse them and/or develop dependence. A recent study of Massachusetts workers showed rates of opioid overdose deaths were significantly higher among workers employed in industries and occupations known to have high rates of work-related injuries and illnesses. Additionally, rates were significantly higher among workers in occupations with lower availability of paid sick leave and lower job security, suggesting that the need to return to work soon after an injury may contribute to high rates of opioid-related overdose death (MDPH, 2018). Additional insights into the connection between opioid misuse and employment factors such as employment benefits, working conditions, work design, and work organization are needed to more effectively respond to this challenge.

- Expand research to understand the work organization-related etiology of chronic health conditions, including obesity and metabolic disorders, CVD, mental health disorders, substance use disorders, and cancer, to add to the current knowledge base on possible mechanisms of action and to resolve incongruent findings.
- Expand research to examine the contribution of new forms of work arrangements and design of work (as further detailed in Objectives 2 and 6) to chronic disease risk in the workforce.
- Improve the understanding and awareness of how to design work to decrease risk factors related to chronic health conditions to include comparative effectiveness studies and economic evaluations to compare interventions, policies and practices.
- Conduct research and analyses to better evaluate the monetized and non-monetized costs and other burdens of unhealthy work design related to disease incidence and prevalence, healthcare costs, and organizational factors such as absenteeism, presenteeism, productivity loss, turnover, and reputational risk, to include ways of assessing their importance for inclusion in a business case to improve work design.

- Incentivize employers to improve the design of work and implement healthier work organization principles shown to decrease risks for chronic disease and improve the workplace management of them.
- Identify and establish new networks and collaborations with professionals in chronic disease prevention and management, leadership and management, human resources and employee benefits, occupational health, and other areas to expand the visibility and application of HWD research findings.
- Conduct implementation and dissemination research that examines factors that reduce chronic disease
 outcomes among workers and influences the adoption of work design policies and practices among
 organizations and industries.
- Develop and evaluate workplace and related clinical interventions that integrate assessment and remediation of work-related contributors to chronic health conditions.
- Conduct research on policies, programs and practices that integrate protection from work-related safety and health hazards with promotion of injury and illness prevention efforts to advance worker well-being and to communicate knowledge and effective best practices in collaboration with existing and new partners, including employers and the community.

Objective 5: Decrease the Burden of Shift Work, Long Hours of Work, and Sleep Deficiency

Background

Sleep deficiency is a broad term that includes inadequate sleep duration, poor sleep quality, untreated sleep disorders, and mistimed sleep that is not synchronized with circadian rhythms. Sleep deficiency can be caused by both work and personal factors (e.g., race and ethnicity, culture, socioeconomic status, long commutes) and contributes to a variety of safety and health risks both on- and off-the-job. In 2004, almost 15% of full-time workers were employed in evening, night, rotating, split, or employer-arranged irregular shifts (McMenamin, 2007). According to American Time Use Survey analyses during 2003–2011 and National Health Interview Survey analyses during 1997-2011, 10% of Americans worked multiple jobs. Working multiple jobs is reported to impact sleep duration and quality and increase the risk of and incidence of injury (Marucci-Wellman, Lombardi, & Willetts, 2016; Marucci-Wellman, Willetts, Lin, Brennan, & Verma, 2014). In some instances, long work hours and night work cannot be eliminated because society needs critical services around-the-clock for occupations such as police and fire protection, health care, transportation, and public utilities. A recent study estimated that insufficient sleep cost the overall U.S. economy upwards of \$411 billion (2.28% of its GDP) in 2015, due to a variety of negative impacts (Hafner et al., 2016).

Research is needed to develop effective administrative controls for managers and individual strategies for workers to use in their personal life. Research is also needed on many types of interventions to reduce risks, including education programs, testing various work scheduling patterns, remote work, manipulating light exposure, pharmacological agents, diet regimens, work organization strategies and efforts to change workplace culture, workplace interventions including policies, fatigue risk management systems, mathematical models to predict risks, and studies of the impact of broader public policy measures such as the impact of State laws that ban mandatory overtime in nurses.

Work factors

Work factors associated with poor sleep quality and duration include shift work, long work hours, and job stress.

Shift work and long work hours also negatively affect personal relationships and responsibilities due to tired workers' poor moods and reduced quality time to spend with their families. Several studies show shift work and long work hours increase the risk for work-life conflict (Albertsen et al., 2008; Carlson & Perrewe 1999). (See Objective 7 on work-life conflict.) Additional research demonstrates the adverse impacts of workplace stressors, such as work-life conflict and unsupportive supervisors, on sleep quality, duration, and fatigue (Crain et al., 2014). Certain employee groups, such as nurses, experience excessive physical and psychological stress, and increased work pace. Because physiologic recovery from stress is delayed past the cessation of stressful events, sleep may become more difficult to initiate and maintain among these workers. Over time, this may lead to maladaptive fatigue, characterized by problems with supervisors, peers, increased emotional demand and mental load (Winwood & Lushington, 2006).

Personal factors

Personal factors that lead to fatigue on the job include sleep disorders and not allowing enough time for sleep for a variety of personal and other reasons. For example, care-giving responsibilities, long commutes, and working at more than one job can cut down on the time available for sleep. An estimated 18 million American adults had obstructive sleep apnea in 2006 (Institute of Medicine, 2006) and an estimated 30 million had insomnia in 2017 (National Sleep Foundation, 2017). A recent literature review, representing more than 12,000 participants in 10 studies published from 2000 to 2015, indicated that workers with obstructive sleep apnea experienced just over twice as many occupational injuries and other adverse events as workers without obstructive sleep apnea (Garbarino et al., 2016). In truck drivers, sleep restriction, sleep apnea, and daytime sleepiness were all significantly associated with motor vehicle crashes and near misses.

Sleep deficiency endangers safety and health

Workers with sleep deficiency can show reduced ability to perform work and other tasks. Studies report that performance when awake 17 or more hours is similar to performance when intoxicated with alcohol (Dawson & Reid, 1997; Williamson & Feyer, 2000). Workers with sleep deficiency can show a general slowing of cognitive functioning and degradation in attention, vigilance, processing speed, and alertness (Kilgore, 2010). They can also show impairment in memory consolidation (Havekes & Abel, 2017). In addition, a person who experiences sleep deprivation may not see or recognize a hazard quickly enough to avoid it, or may recognize the hazard, but respond too slowly to counteract it (Rakitin et al., 2012). Cognitive responses due to sleep deficiency degrade in a dose-response manner (Goel et al., 2014), but do not recover in a dose-response manner. That is, multiple nights of recovery sleep may be required to recover from sleep-related cognitive impairments (Banks et al., 2010).

The safety of the public is a concern when workers do not get enough good quality sleep. Rosekind (2015) reported that up to 20% of the fatal vehicle crashes investigated by the National Transportation Safety Board were due to drowsy driving. Investigators of several well-known industrial disasters reported worker fatigue to be one of the causal factors. These disasters include the 2005 BP Texas City explosion, the Buffalo jet crash, and the Exxon Valdez oil spill (Baker Panel, 2007; National Transportation Safety Board, 2004; National Transportation Safety Board, 2009).

In addition to the safety risks, sleep deficiency, shift work, and long work hours are associated with a broad range of health risks. These include premature death, obesity, adverse reproductive outcomes, infections, and a wide range of chronic illnesses including cardiovascular, gastrointestinal, and musculoskeletal disorders, diabetes mellitus, cancer, Alzheimer's disease, and disturbances to mood and cognition (Bubu et al., 2017; Durmer & Dinges, 2005: Irwin, 2015; Luyster et al., 2012; NIOSH, 2015).

Research gaps

• Develop surveillance methods that measure the prevalence and characteristics of shift work, long work

hour schedules, and other factors that lead to fatigue in the workplace.

- Identify and test interventions that reduce the adverse safety and health consequences of shift work, long work hour schedules, and other factors that lead to fatigue in the workplace.
- Develop and test interventions for workers on shift work and long work hour schedules in occupations involving public safety (e.g., health care, law enforcement, transportation, utilities) where the consequences of errors due to fatigue can be especially great to the public.
- Develop implementation and dissemination strategies for effective interventions that reduce the adverse health and safety consequences of shift work, long work hour schedules, and other factors that lead to fatigue in the workplace.

Objective 6: Improve the Safety, Health, and Well-being of Workers through Healthier Work Design and Better Organizational Practices

Background

Work design has implications for the safety, health, well-being, and functioning of individuals, families, groups, organizations, and communities. Workplaces are settings to target not only work-related risks, such as unsafe working conditions, high job demands, and low job control, but also to promote workplace conditions and programs that provide support for workers' health and well-being, such as smoking cessation or promotion of healthy physical activity (Sorensen et al., 2010). Worker well-being characterizes quality of life with respect to an individual's health and work-related environmental, organizational, and psychosocial factors. Worker well-being includes the concept of work that allows individuals to reach their full potential along with the recognition that positive work design has an impact on worker functioning and satisfaction both within the organization and in the community and society (Schulte & Vainio, 2010). Healthy work design includes attention to job tasks, the physical and psychosocial work environment, workplace policies and practices, and organizational climate and culture. Efforts that focus on prevention of safety and health hazards and promotion of well-being typically involve multilevel approaches that include commitment and involvement from management as well as worker input on identification of strategies most effective in reducing unreasonable work demands (e.g., workload, work pace), workplace physical hazards, precarious work, and hazards associated with non-standard work schedules. (See Objective 2 on non-standard work arrangements and Objective 5 on work schedules). Organizational factors that can differentiate successful from unsuccessful employer efforts to reduce occupational injury and illness and promote worker well-being include aligning programs with the organization's mission and values; coordinating multi-level or cross-organization efforts; tailoring initiatives to meet the unique needs of the organization and its workers; garnering support from senior leaders; designing effective two-way communications; and promoting ongoing evaluation and a philosophy of continuous improvement (Fitz-enz, 1993; Goetzel et al., 2007; Grawitch et al., 2009).

Hazard control and job design

Safety and health initiatives that focus on eliminating workplace safety and health hazards first and then changing workplace systems to improve the physical and mental health of workers are the preferred approach. A wide variety of workplace practices can help workers improve their physical and mental health, reduce health risks, minimize chronic illness and disability, and reduce stress. These include efforts to identify and eliminate workplace safety and health hazards, ensure that workspaces and tools are ergonomically designed to avoid injury, ensure that workload and work pace are reasonable, and provide sufficient autonomy, control, and task variety. These efforts also include primary-level interventions that change the design of both physical workspaces and work

processes. Although sustained efforts have been made to control workplace safety and health hazards, more than half of Americans still report exposure to potentially hazardous working conditions (RAND, 2017).

Psychosocial workplace conditions

There is a need for additional strategies and interventions to improve psychosocial workplace conditions. For example, in 2017, most American workers reported having to work at high speeds or under tight deadlines, or having too little time to do their job, and almost one in five American workers reported experiencing a hostile or threatening work environment (RAND, 2017). Job insecurity is another critical psychosocial stressor within many sectors of the workforce that is deserving of more attention (Sinclair et al., 2010). While job stress is related to adverse safety, health, and well-being outcomes and some interventions have been identified that address contributors to job stress (Hurrell & Sauter, 2013; Sauter & Murphy, 2006), more research is needed.

Organizational culture and psychosocial climate

Kelloway's (2015) commentary on Grawitch, Ballard, & Erb (2015) points out that "a psychologically healthy workplace is far more than a mere list of programs and initiatives. Instead, it is a description of an organizational culture—a culture that provides the foundation for the psychological health and wellbeing of employees" (p. 263). The culture should facilitate the development and utilization of workplace practices intended to improve worker well-being. Psychological factors such as respect, trust, feeling valued, and management and organizational support also play key roles in worker well-being. In the American Psychological Association's (APA) 2016 survey, workers who felt valued were more likely to say they regularly participated in their employer's health and wellbeing programs, training activities, and involvement efforts and were less likely to report chronic job stress (APA, 2016). Access to a health and well-being program at work can have health and productivity benefits. NIOSH analyses of 2014 Quality of WorkLife survey data indicate that workers who had a stress management program at their workplace had 1.5 fewer days of poor mental health per month than workers who did not have stress management programs at their workplace (NIOSH, 2017).

Employee involvement and autonomy

Increasing employee involvement in decision-making and providing sufficient autonomy and control can increase job satisfaction, employee morale, and commitment to the organization as well as increase productivity, reduce turnover and absenteeism, and enhance the quality of products and services (Freeman & Rogers, 1999; Lawler, 1991; Vandenberg, Richardson, & Eastman, 1999). Increased worker control, particularly in high demand work conditions, has been associated with reduced risk of cardiovascular disease, improved mental health (Mikkelsen & Saksvik, 1999), and with a significantly reduced risk of mortality (Mule & Cockburn, 2017). Bourbonnais et al. (2006) have demonstrated the positive effects of worker participation in reducing adverse workplace psychosocial factors and improving worker mental health. Despite strong evidence of the value of worker involvement and autonomy, according to a 2017 survey of the U.S. workforce, only 58% of working Americans reported that their employer provided sufficient opportunities to be involved in decision-making, and even fewer (53%) said they regularly participated in activities involving decision-making, solving problems, or setting goals (APA, 2017). Another national study by RAND (2017) found that one in three Americans reported no control over their work schedules. Thus, increasing control over how and when workers do their work, and increasing opportunities for workers to be involved in workers are important to address.

Career development

Opportunities for advancement and career development, promotion and compensation policies and practices that are fair and equitable, and reward and recognition mechanisms that are commensurate with employee effort, are important contributors to employee well-being. The opportunity to gain new skills and experiences can increase employee motivation and job satisfaction and help workers more effectively manage job stress (Pfeffer, 1994;

Browne, 2000). These practices include self-set goals and feedback, as well as the ways performance is measured and how related data are used. In APA's 2017 survey, lack of opportunity for growth or advancement was second only to low salaries as a source of job stress; only 60% of the U.S. workforce reported being satisfied with the development opportunities offered by their employer, just 54% said that employee recognition was based on a fair and useful evaluation system, and less than half (49%) said their employer provided sufficient opportunity for internal advancement (APA, 2017). As job satisfaction has been shown to influence worker mental and physical health, career development and advancement opportunity influences on job satisfaction are important from a worker health standpoint as well (Ferragher et al., 2005).

- Explore effective strategies for comprehensive systems-based interventions (i.e., collective sets of organization and work practices and how they relate to and interact with each other in real-world settings) and evaluation of those interventions.
- Develop interventions to improve the impact of management systems, quality improvement methods, leadership styles, and supervisor and co-worker support approaches on employee and organizational outcomes.
- Identify best-practice methods for assessing worker well-being in individual workplaces and identifying
 deficits that need to be addressed. Special attention should focus on understanding how these best
 practice methods may vary by worker or organization characteristics such as age, ethnicity, occupational
 group, and industry sector.
- Develop effective evidence-based design, implementation and dissemination strategies for best practice assessment, interventions, and program evaluation, critical to the translation of research to practice. Emphasize evidence-based methods that are cost effective and realistic for small- and medium-size organizations without dedicated staff or access to internal experts.
- Evaluate labor-management relationships and involvement in establishing overall work policies and practices and the related implications for worker safety, health, and well-being.
- Collect and analyze data on differences in worker and organizational needs, preferences, practices, and outcomes by industry, region, size, structure, and job type.
- Identify characteristics of the built environment (e.g., ergonomics, workspace design) that, in combination with psychosocial work environment characteristics, promote worker safety, health, and well-being.
- Develop and evaluate organizational and managerial strategies to respond to misuse of opioids and illicit drugs by employees. These strategies could include:
 - Identifying and correcting demanding work organization factors that increase the risk for substance misuse by workers;
 - o Identifying ways to assist workers with substance misuse while protecting worker privacy;
 - Identifying ways of supporting workers with substance misuse through return to work assistance and continued employment while protecting worker and public safety;
 - Facilitating effective treatment through anti-stigma interventions and improved treatment access.

Objective 7: Promote a Sustainable Work-Nonwork Interface

Background

The boundaries between people's work and non-work lives (e.g., family and leisure) continue to blur, with 42% of U.S. workers in 2014 reporting that they take care of personal or family needs during work and about a quarter indicating that they regularly bring work home, work during vacations, and allow work to interrupt time with family and friends (APA, 2015). A U.S. nationally representative study found that, in 2015, about one-half of workers indicated doing some work in their free time to meet the demands of their jobs and 31% of workers reported that it was somewhat or very difficult to adjust their work schedule to accommodate a personal matter (RAND, 2017). For many workers, such blurring of the boundaries between work and non-work demands leads to work-life conflict, which is now viewed as an occupational hazard (Hammer & Sauter, 2013).

Work-life conflict

Research has demonstrated that work-life conflict, or the conflicting demands of work-nonwork responsibilities, is associated with adverse outcomes for workers, their families, and their employers (Amstad, Meier, Fasel, Elfering, & Semmer, 2011; Hammer & Sauter, 2013). Work-life conflict is related to adverse workplace safety outcomes (Cullen & Hammer, 2007; Smith & DeJoy, 2012; Turner et al., 2014). Workers who experience work-life conflict also report low career, job, family, and life satisfaction, poor sleep quality, poor health behaviors, and physical and mental health problems. In addition, work-family conflict affects employers through high levels of employee absenteeism, burnout, and turnover, as well as poor job performance, low commitment, and poor organizational citizenship behaviors.

Managing the work-nonwork interface

Despite the research demonstrating the adverse worker health and safety effects of work-life conflict, few workers have access to benefits and services that can help them better manage work-nonwork responsibilities. Such access does have health and safety benefits for both the workers and their families. For example, parental access to paid sick leave is associated with significantly higher odds of children receiving annual medical checkups and vaccinations, and significantly lower odds of children receiving delayed medical care or being taken to the emergency room (Asfaw et al., 2017). Low-income women, in particular, cannot take time off to care for their children or themselves (Clemens-Cope et al., 2008; Shepherd-Banigan & Bell, 2014).

Other employer-provided services and benefits are similarly limited for workers. In 2016, only 13% of private sector organizations provided access to paid family leave (BLS, 2016). In 2017, only 4% of employers subsidized childcare and 17% of employers offered referral to childcare services (SHRM, 2017). Also in 2017, 13% of employers offered referral to elder care services, 62% offered some type of teleworking, and 51% offered some type of flexible schedules (SHRM, 2017). Interest in workplace flexibility and managing the work and non-work interface is not based on family obligations, exclusively. The non-work domain includes many activities, including leisure, educational, spiritual, and community (Keeney et al., 2013), and workers without families also desire access to work-life initiatives (Ryan & Kossek, 2008).

While reducing work-life conflict and the related negative consequences is critical, successful management of the work-nonwork interface requires more than simply mitigating negative experiences. Organizational strategies that can be implemented to improve the work-nonwork interface include increasing supervisor support for managing work and family demands; minimizing telepressure; increasing worker control over where, when, and how they work; and decreasing the excessive demands of work, especially among lower-income workers including workers in low hourly wage jobs (Kelly et al., 2008). Training supervisors on family supportive behaviors has been shown to have positive impacts on worker job satisfaction and physical health (Hammer et al., 2011), as well as workplace

productivity (Odle-Dusseau et al., 2016). (See objective 6 on organizational practices.) Work can also be a source of personal enrichment and enjoyment, with opportunities for meaningful work and personal development.

- Define well-being as it relates to successful management of the work-nonwork interface. Raise awareness of how workplace experiences across the occupational spectrum can spill over to affect the well-being of families and communities.
- Conduct research on contextual factors and their implications for the work-nonwork interface. Examples of organizational contextual factors include job type, industry, organizational culture, leadership, management skills, and available technology. Examples of individual contextual factors include decision-making and demographic considerations, such as age, race and ethnicity, language ability, education, income, marital status, and family make-up.
- Include measures that characterize quality of work-life, work-non-work balance, and worker well-being in ongoing public health surveillance efforts.
- Investigate the safety and health effects of work-life conflict of workers in jobs without much stability or security and workers in low-paying hourly jobs with limited flexibility to deal with family demands or emergencies and continual financial stress.
- Examine safety and health implications associated with commuting, work travel, and work-life conflict.
- Design and implement work-non-work interventions, including policies, programs, and practices, that reduce conflict and support worker and family safety, health, and well-being. These interventions should be sustainable over time, inclusive of all employees with differing non-work needs and demands, promoted by employers, and willingly adopted by organizational members.
- Translate science on known work-life risks to increase awareness of those risks and use it to inform practice-based research.

REFERENCES

Albertsen, K., Rafnsdóttir, L., Grimsmo, A., Tómasson, K., & Kauppinen, K. (2008). Workhours and worklife balance. Scandinavian Journal of Work, Environment and Health, Suppl 5:14-21.

American Heart Association (2000). "Eliminating Cardiovascular Disease Would Increase U.S. Life Expectancy By Seven Years." Science Daily, 4 January 2000. Retrieved from http://www.sciencedaily.com/releases/2000/01/000104065154.htm

American Psychological Association. (2015). Work-Life Survey. Retrieved from http://www.apaexcellence.org/assets/general/2015-work-life-survey-results.pdf

American Psychological Association. (2016). 2016 Work and Well-Being Survey. Retrieved from http://www.apaexcellence.org/assets/general/2016-work-and-wellbeing-survey-results.pdf

American Psychological Association. (2017). 2017 Work and Well-Being Survey. Retrieved from http://www.apaexcellence.org/assets/general/2017-work-and-wellbeing-survey-results.pdf

Amstad, F. T., Meier, L. L., Fasel, U., Elfering, A., & Semmer, N. K. (2011). A meta-analysis of work–family conflict and various outcomes with a special emphasis on cross-domain versus matching-domain relations. Journal of Occupational Health Psychology, 16, 151-169. doi:10.1037/a0022170

Asfaw, A. & Colopy, M. (2017). Association between parental access to paid sick leave and children's access to and use of healthcare services. American Journal of Industrial Medicine, 60(3), 276-284.

Baker Panel (2007). The report of the BP U.S. refineries independent safety review panel. Retrieved from <u>http://www.csb.gov/assets/1/19/csbfinalreportbp.pdf</u>

Banks, S., Van Dongen, H. P. A., Maislin, G., & Dinges, D. F. (2010). Neurobehavioral dynamics following chronic sleep restriction: Dose-response effects of one night for recovery. Sleep, 33, 1013 – 1026.

Benavides, F. G., Benach, J., Muntaner, C., Delclos, G. L., Catot, N., & Amable, M. (2006). Associations between temporary employment and occupational injury: What are the mechanisms? Occupational and Environmental Medicine, 63, 416–421.

BLS. (2015). Employment Projections - 2014-2024.BLS News release USDL-15-2327. Accessed at <u>https://www.bls.gov/news.release/pdf/ecopro.pdf</u>.

BLS (2016). <u>https://www.bls.gov/opub/ted/2016/13-percent-of-private-industry-workers-had-access-to-paid-family-leave-in-march-2016.htm</u>

BLS (2017). <u>https://www.bls.gov/opub/ted/2017/93-percent-of-managers-and-46-percent-of-service-workers-had-paid-sick-leave-benefits-in-march-2017.htm</u>

BLS (2017). https://www.bls.gov/ncs/ebs/benefits/2017/ownership/private/table01a.pdf

BLS (2017). https://www.bls.gov/iif/oshwc/cfoi/cfch0015.pdf

<u>BLS (2017). Economic News Release: Census of Fatal Occuational Injuries Summary, 2016. Washington, D.C.:</u> <u>Bureau of Labor Statistics, December 19. https://www.bls.gov/news.release/cfoi.nr0.htm</u>

BLS (2017). Employment Projections 2016 to 26. USDL – 17-1429.

Bourbonnais, R., Brisson, C., Vinet, A., Vezina, M., Abdous, B., & Gaudet, M. (2006). Effectiveness of a participative intervention on psychosocial work factors to prevent mental health problems in a hospital setting. Occupational and Environmental Medicine, 63, 335-342.

Browne, J. H. (2000). Benchmarking HRM practices in healthy work organizations. The American Business Review, 18, 54-61.

Bubu, O. M., Brannick, M., Mortimer, J., Umasabor-Bubu, O., Sebastiao, Y. V., Wen, Y., et al. (2017). Sleep, Cognitive impairment, and Alzheimer's disease: A Systematic Review and Meta-Analysis. Sleep, 40, 10.

Bushnell, T., Scharf, T., Alterman, T., Cummings, K. J., Luckhaupt, S. E., Ray, T. K., Rosa, R. R., & Su, C. P. (2017). Developing a Taxonomy of Work Arrangements to Examine Relationships with Worker Safety, Health, and Well-Being. Poster presented at Work, Stress and Health 2017, Minneapolis, MN, June 8, <u>http://www.apa.org/wsh/preliminary-program.pdf</u>

Byler, C. G. (2013). Hispanic/Latino fatal occupational injury rates. Monthly Labor Review. February, 2013, 14-23.

Carlson, D. S. & Perrewe, P. L. (1999). The role of social support in the stressor-strain relationship: an examination of work-family conflict. Journal of Management, 25, 513-540.

Cawley, J. & Meyerhoefer, C. (2012). The medical care costs of obesity: an instrumental variables approach. Journal of Health Economics. 31, 219-230.

CBO. 2001. Social Security: The Challenges of an Aging Population. CBO testimony on aging before the special committee of aging. Accessed at

https://www.cbo.gov/sites/default/files/cbofiles/ftpdocs/32xx/doc3214/ssptestimony.pdf

CDC (2016). CDC guideline for prescribing opioids for chronic pain – United States, 2016. MMWR, 65(1), 1-49.

CDC (2017). Provisional counts of drug overdose deaths as of 8/16/2017. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Health Statistics, National Vital Statistics System. <u>https://www.cdc.gov/nchs/data/health_policy/monthly-drug-overdose-deathestimates.pdf</u>

Hui, M., Manyika, J., & Miremadi, M. (2016). Where machines could replace humans – and where they can't (yet). McKinsey Quarterly, 2016 Issue 3, 49-57.

Clemens-Cope, L., Perry, C. D., Kenney, G. M., Pelletier, J. E., & Pantelli, M. S. (2008). Access to and use of paid sick leave among low-income families with children. Pediatrics, 122, e480-e486.

Cooper, R., Cutler, J., Desvigne-Nickens, P., et al. (2000). Trends and disparities in coronary heart disease, stroke, and other cardiovascular diseases in the United States: Findings from the National Conference on Cardiovascular Disease Prevention. Circulation 102:3137–3147.

Costa, G., Haus, E., & Stevens, R. (2010). Shift work and caner – Considerations on rationale, mechanisms, and epidemiology. Scandinavian Journal Work, Environment, and Health, 36, 163-179.

Crain, T. L., Hammer, L. B., Bodner, T., Kossek, E. E., Moen, P., Lilienthal, R., & Buxton, O. M. (2014). Work-family conflict, family-supportive supervisor behaviors (FSSB), and sleep outcomes. Journal of Occupational Health Psychology, 19, 155–167.

Cullen, J. C. & Hammer, L. B. (2007). Developing and testing a theoretical model linking work-family conflict to employee safety. Journal of Occupational Health Psychology, 12, 266-278.

Dawson, D. & Reid, K. (1997). Fatigue, alcohol and performance impairment. Nature, 388, 235.

De Moortel, D., Vandenheede, H., & Vanroelen, C. (2014). Contemporary employment arrangements and mental well-being in men and women across Europe: A cross-sectional study. International Journal for Equity in Health, 13(1), 90.

Durmer, J. S. & Dinges, D. F. (2005). Neurocognitive consequences of sleep deprivation. Seminars in Neurology, 25, 117-129.

EEOC (2018). : <u>https://www.eeoc.gov/eeoc/publications/brochure-gender_stereotyping.cfm.</u>

Faragher, E. B., Cass, M., & Cooper, C. L. (2005). The relationship between job satisfaction and health: a metaanalysis. Occupational and Environmental Medicine, 62, 102-112.

Finkelstein, E., Fiebelkorn, I. C., & Wang, G. (2005). The Costs of Obesity among Full-Time Employees. American Journal of Health Promotion, 20, 45 - 51

Finkelstein E. A., DiBonaventura M. D., Burgess S. M. & Hale B. C. (2010). The costs of obesity in the workplace. Journal of Occupational and Environmental Medicine, 52, 971-97

Fitz-enz, J. (1993). The truth about "best practice". Human Resource Planning, 16, 19-26.

Foley, M. (2017). Factors underlying observed injury rate differences between temporary workers and permanent peers. American Journal of Industrial Medicine, 60(10):841-851. doi: 10.1002/ajim.22763

Freeman, R. B. & Rogers, J. (1999). What workers want. Ithaca, NY: Russell Sage Foundation.

Fusaro, V. A. & Shaefer, H. L. (2016). How should we define "low-wage" work? An analysis using the Current Population Survey. Monthly Labor Review, U.S. Bureau of Labor Statistics, October. (https://doi.org/10.21916/mir.2016.44.)

Gallup (2017). In U.S. more adults identifying as LGBT. <u>http://news.gallup.com/poll/201731/lgbt-identification-rises.aspx</u>

Garbarino, S., Guglielmi, O., Sanna, A., Mancardi, G. L., & Magnavita, N. (2016). Risk of Occupational Accidents in Workers with Obstructive Sleep Apnea: Systematic Review and Meta-analysis. Sleep, 39, 1211-1218.

Garza, J. L., Dugan, A.G., Faghri, P.D., Gorin, A.A., Huedo-Medina, T. B., Kenny, A.M., Cherniack, M.G., & Cavallari, J.M. (2015) Demographic, health-related, and work-related factors associated with body mass index and body fat percentage among workers at six Connecticut manufacturing companies across different age groups: a cohort study. BMC Obesity. 2:43:1-18.

Gates, G. J. (2014). LGBT demographics: Comparisons among population-based surveys. <u>https://williamsinstitute.law.ucla.edu/wp-content/uploads/lgbt-demogs-sep-2014.pdf</u>

Goel, N., Abe, T., Braun, M., & Dinges, D. (2014). Cognitive workload and sleep restriction interact to influence sleep homeostatic responses. Sleep, 37, 1745 – 1756.

Goetzel, R. Z., Shechter, D., Ozminkowski, R. J., Marmet, P. F., Tabrizi, M. J., & Roemer, E. C. (2007). Promising practices in employer health and productivity management efforts: Findings from a benchmarking study. Journal of Occupational and Environmental Medicine, 49(2):111-130.

Goetzel, R. Z., Pei, X., Tabrizi, M. J., Henke, R. M., Kowlessar, N., Nelson, C. F., & Metz, R. D. (2012). Ten modifiable health risk factors are linked to more than one-fifth of employer-employee health care spending. Health Affairs, 31, 2474-2484.

Gonzales-Mule, E. & Cockburn, B. (2017). Worked to death: The relationship of job demands and job control with mortality. Personnel Psychology, 70, 73-112.

Grawitch, M. J., Ballard, D. W., & Erb, K. R. (2015). To be or not to be (stressed): The critical role of a psychologically healthy workplace in effective stress management. Stress and Health, 31, 264-273.

Grawitch, M. J., Ledford, G. E., Ballard, D. W., & Barber, L. K. (2009). Leading the healthy workforce: The integral role of employee involvement. Consulting Psychology Journal: Practice and Research, 61, 122-135.

Gu, J. K., Charles, L. E., Andrew, M. E., Ma, C.C., Hartley, T. A., Violanti, J. M., & Burchfiel, C. M.(2016) Prevalence of work-site injuries and relationship between obesity and injury among U.S. workers: NHIS 2004-2012. Journal of Safety Research. 58:21-30.

Hafner, M., Stepanek, M., Taylor, J., Troxel, W. M., & Van Stolk, C. (2016). *Why sleep matters - the economic costs of insufficient sleep: A cross-country comparative analysis.* Santa Monica, CA: RAND Corporation.

Hammer, L. B. & Sauter, S. L. (2013). Total worker health and work-life stress. Journal of Environmental and Occupational Medicine, 55, S25-S29.

Hancock, P. A. (2009). Mind, machine and morality: Toward a philosophy of human-technology symbiosis. Farnham: Ashgate.

Havekes, R. & Abel, T. (2017). The tired hippocampus: The molecular impact of sleep deprivation on hippocampal function. Current Opinions in Neurobiology, 44, 13-19.

Henly, J. R. & Lambert, S. (2014). Unpredictable work timing in retail jobs: Implications for employee work-life outcomes. Industrial and Labor Relations Review, 67(3), 986-1016.

Heron, M. (2016). Deaths: Leading causes for 2014. National vital statistics reports; 65(5). Hyattsville, MD: National Center for Health Statistics. <u>http://www.cdc.gov/nchs/data/nvsr/nvsr65/nvsr65_05.pdf</u>.

Howard, J. (2017). Nonstandard work arrangements and worker health and safety. American Journal of Industrial Medicine, 60(1):1-10.

Huffman, J. C., Celano, C. M., Beach, S. R., Motiwala, S. R., & Januzzi, J. L. (2013). Depression and cardiac disease: Epidemiology, mechanisms, and diagnosis. Cardiovascular Psychiatry and Neurology, Volume 2013, Article ID 695925: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3638710/pdf/CPN2013-695925.pdf.

Hurrell, J. J., Jr. & Sauter, S. L. (2013). Job stress prevention: An overview of approaches. In A.M. Rossi, J.A. Meurs & P.L. Perrewé (Eds.), *Stress & quality of working life: Improving employee health & well-being* (pp. 187-206). Greenwich, CT: Information Age Publishing.

Institute of Medicine (IOM) Committee on Sleep Medicine and Research, Colten, H.R. & Altevogt, B.M. (ed.). (2006). *Sleep Disorders and Sleep Deprivation: An Unmet Public Health Problem*, Washington, DC: National Academy of Sciences.

International Agency for Research on Cancer (2018). https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3638710/pdf/CPN2013-695925.pdf

International Commission on Occupational Health (2013). The Tokyo declaration on prevention and management of work-related cardiovascular disorders. International Commission on Occupational Health (ICOH) Newsletter 11;(2,3):4.

Irwin, M. R. (2015). Why sleep is important for health: A psychoneuroimmunology perspective. Annual Review of Psychology, 66, 143-172.

Judge, T. A. & Cable, D. M. (2011). When it comes to pay, do the thin win? The effect of weight on pay for men and women. Journal of Applied Psychology, 96, 95-112. doi: 10.1037/a0020860.

Katz, L. F. & Krueger, A. B. (2016). The rise and nature of alternative work arrangements in the United States 1995-2015 (Working Paper 22667). Cambridge, MA: National Bureau of Economic Research.

Katz, L. F. & Krueger, A. B. (2019). Understanding trends in alternative work arrangements in the United States. Working Paper 25425, <u>http://www.nber.org/papers/w25425</u>. National Bureau of Economic Research, Cambridge, MA.

Keeney, J., Boyd, E. M., Sinha, R., Westring, A. F., & Ryan, A. M. (2013). From "work-family" to "work-life": Broadening our conceptualization and measurement. Journal of Vocational Behavior, 82(3), 221-237.

Kelloway, E. K. (2015). The psychologically healthy workplace. Stress and Health, 31, 263.

Kelly, E. L., Kossek, E. E., Hammer, L. B., Durham, M., Bray, J., Chermack, K., Murphy, L. A., & Kaskubar, D. (2008). Getting there from here: Research on the effects of work-family initiatives on work-family conflict and business outcomes. The Academy of Management Annals, 2, 305-349.

Killgore, W. D. S. (2010). Effects of sleep deprivation on cognition. Progress in Brain Research, 185, 105 – 129.

Kowalski-McGraw, M., Green-McKenzie, J., Pandalai, S., & Schulte, P.A. (2017). Characterizing the interrelationships of prescription opioid and benzodiazepine drugs with worker health and workplace hazards. Journal of Occupational and Environmental Medicine,

Landsbergis, P., Cahill, J., & Schnall, P. (1999) The Impact of Lean Production and Related New Systems of Work Organization on Worker Health. Journal of Occupational Health Psychology.492):108-30.

Landsbergis, P., Zoeckler, J., Kashem, Z., Rivera, B., Alexander, D., & Baruth, A. (2017) Organizational Policies and Programs to Reduce Job Stress and Risk of Workplace Violence Among K-12 Education Staff. New Solutions. 27(4):559-580.

Lawler, E. E., III. (1991). Participative management strategies. In J.W. Jones, B.D. Steffy, & D.W. Bray (Eds.), *Applying psychology in business: The handbook for managers and human resource professionals* (pp. 578-586). Lexington, KY: Lexington Books.

Lee, M. K., Kusbit, D., Metsky, E., & Dabbish, L. (2015). Working with machines: The impact of Algorithmic and data-driven management on human workers. Presented at CHI 2015, April 18-23, Seoul, Korea.

Leigh, P. J. & Miller, T. R. (1998). Job-related diseases and occupations within a large workers' compensation data set. American Journal of Industrial Medicine. 33:197–211.

Leigh, P. J. & De Vogli, R. (2016). Low wages as occupational health hazards. Journal of Occupational and Environmental Medicine, 58(5), 444-447.

Luyster, F. S., Strollo, P. J., Zee, P. C., & Walsh, J. K. (2012). Sleep: A health imperative. Sleep, 35(6):727-734. DOI: 10.5665/sleep.1846

Marucci-Wellman, H. R., Lombardi, D. A., & Willetts, J. L. (2016). Working multiple jobs over a day or a week: Short-term effects on sleep duration. Chronobiology International, 33, 630-649.

Marucci-Wellman, H. R., Willetts, J. L., Lin, T. C., Brennan, M. J., & Verma, S. K. (2014). Work in multiple jobs and the risk of injury in the US working population. American Journal of Public Health, 104, 134-142.

McMenamin, T. M. (2007). A time to work: recent trends in shift work and flexible schedules. Monthly Labor Review, 130, 3-15.

MDPH (Massachusetts Department of Public Health) (2018). Opioid-Related Overdose Deaths in Massachusetts by Industry and Occupation, 2011-2015. Boston, MA: MDPH Occupational Health Surveillance Program.

Mikkelsen, A. & Saksvik, P.O. (1999). Impact of a participatory organizational intervention on job characteristics and job stress. International Journal of Health Services, 29(4), 871-893.

National Academies of Sciences, Engineering, and Medicine (2017). Information Technology and the U.S. Workforce: Where Are We and Where Do We Go from Here? Washington, DC: The National Academies Press. doi:10.17226/24649.

National Center for Chronic Disease Prevention and Health Promotion (2016). Adult Obesity Causes and Consequences. Accessed 8/18/2017, <u>https://www.cdc.gov/obesity/adult/causes.html</u>

National Institute of Mental Health (2015). Major Depression among Adults. Accessed September, 2017. <u>https://www.nimh.nih.gov/health/statistics/prevalence/major-depression-among-adults.shtml</u>

National Institute of Mental Health (2016). From Any Mental Illness (AMI) Among U.S. Adults <u>https://www.nimh.nih.gov/health/statistics/prevalence/any-mental-illness-ami-among-us-adults.shtml</u>

National Institute for Occupational Safety and Health (1999). Stress at Work. DHHS (NIOSH) Publication Number 99-101. Accessed September, 2017. <u>https://www.cdc.gov/niosh/docs/99-101/</u>

National Institute for Occupational Safety and Health (2015). NIOSH Training For Nurses On Shift Work and Long Work Hours. By Caruso, C.C., Geiger-Brown, J., Takahashi, M., Trinkoff, A., Nakata, A. Cincinnati, OH: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 2015-115, Retrieved February 1, 2017, from www.cdc.gov/niosh/docs/2015-115/

National Institute for Occupational Safety and Health (2017). Healthy Work Design and Well-Being Program. U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 2017-198.

National Sleep Foundation. (2017). Sleep apnea. Retrieved from <u>https://sleepfoundation.org/sleep-disorders-problems/sleep-apnea</u>

National Transportation Safety Board [2004]. Grounding of U.S. Tank ship Exxon Valdez on Bligh Reef, Prince William Sound near Valdez, Alaska, March 04 1989. NTSB/ MAR-90/04, Washington D.C.: NTSB.

National Transportation Safety Board [2009]. Aircraft Accident Report: loss of control on approach, Colgan Air, Inc., operating as Continental connection flight 3407, Bombardier DHC-8-400, N200WQ, Clarence Center, New York, February 12, 2009. Washington, DC: National Transportation Safety Board. <u>http://www.ntsb.gov/investigations/AccidentReports/Reports/AAR1001.pdf</u>.

NIDA (2018). Opioids: Brief Description. Washington, DC: National Institutes of Health, National Institute on Drug Abuse. <u>https://www.drugabuse.gov/drugs-abuse/opioids</u>

Odle-Dusseau, H. N., Hammer, L. B., Crain, T. L., & Bodner, T. (2016). The influence of family-supportive supervisors on job performance: An organizational work-family initiative. Journal of Occupational Health Psychology, 21, 296-308.doi: 10.1037/a0039961.

Ogden, C. L., Carroll, M. D., Fryar, C. D., & Flegal, K. M. Prevalence of obesity among adults and youth: United States, 2011–2014. NCHS data brief, no 219. Hyattsville, MD: National Center for Health Statistics. 2015.

Pensa, M.A.; Galusha, D.H.; Cantley, L.F. (2018). Patterns of opioid prescribing and predictors of chronic opioid use in an industrial cohort, 2003 to 2013. Journal of Occupational and Environmental Medicine, 60(5), 457-461.

Pfeffer, J. (1994). Competitive advantage through people. California Management Review, 36, 9-28.

Purcell, K. & Rainie, L. (2015). Technology's impact on workers. Available at: http://www.pewInternet.org/2014/12/30/technologys-impact-on-workers/

Purdue, M. P., et al (2015). The proportion of cancer attributable to occupational exposures. Annals of Epidemiology, 25: 188e192.

Quinlan, M. (2015). The effects of non-standard forms of employment on worker health and safety; International Labour Office, Inclusive Labour Markets, Labour Relations and Working Conditions Branch. Geneva: ILO. (Conditions of work and employment series No. 67.)

Rakitin, B. C., Tucker, A. M., Basner, R. C., & Stern, Y. (2012). The effects of stimulus degradation after 48 hours of total sleep deprivation. Sleep, 35, 113 – 121.

RAND (2017). Working Conditions in the United States: Results of the 2015 American Working Conditions Survey. <u>https://www.rand.org/pubs/research_reports/RR2014.html</u>

Ray, T. K., Kenigsberg, T. A., & Pana-Cryan, R. (2017). Employment arrangement, job stress, and health-related quality of life. Safety Science, May 2017, Epub: <u>https://doi.org/10.1016/j.ssci.2017.05.003</u>.

Rosekind, M. R. (2015). Awakening a nation: a call to action. Sleep Health, 1, 9-10.

Ricci, J. A. & Chee, E. Lost Productive Time Associated with Excess Weight in the U.S. Workforce. Journal of Occupational and Environmental Medicine. 2005; 47 (12):1227–34.

Rus, D. (2015). The robots are coming: How technological breakthroughs will transform everyday life. Foreign Affairs, 94, 2-6.

Rushton, L., Hutchings, S. J., Fortunato, L., et al. (2012). Occupational cancer burden in Great Britain. British Journal of Cancer, 107(Suppl 1):S3-7.

Ryan, A. M & Kossek, E. E. (2008). Work–life policy implementation: Breaking down or creating barriers to inclusiveness? Human Resource Management, 47, 295–310.

Sauter, S. L. & Hurrell, J. J., Jr. (2017). Occupational Health Contributions to the Development and Promise of Occupational Health Psychology. Journal of Occupational Health Psychology, 22, 251–258.

Sauter, S. L. & Murphy, L. R. (2006). Approaches to the management of job stress in the US. In P. Perrewe, A. M. Rossi, & S. L. Sauter (Eds.), *Stress & quality of working life: Current perspectives in occupational health* (pp. 168-183), Greenwich, CT: Information Age Publishing, Inc.

Sauter, S. L., Keel, J., & Hanseman, D. M. (2009). Work organization and health in an aging workforce: Observations from the NIOSH Quality of Work Life Survey. In S. Czaja & J. Sharit (Eds.), *The future of work for an aging population* (pp. 359-393). Baltimore: The Johns Hopkins Press.

Scholl, L., Seth, P., Kariisa, M., Wilson, N., & Baldwin, G. Drug and Opioid-Involved Overdose Deaths — United States, 2013–2017. MMWR Morb Mortal Wkly Rep 2019;67:1419–1427. DOI: http://dx.doi.org/10.15585/mmwr.mm675152e1

Schulte, P. & Vainio, H. (2010). Well-being at work: Overview and perspective. Scandinavian Journal of Work, Environment and Health, 36, 422-429.

Seabury, S. A., Terp, S., & Boden, L. L. (2017). Racial and ethnic differences in the frequency of workplace injuries and prevalence of work-related disability. Health Affairs, 36(2). (<u>https://dol.org/10.1377/hlthaff.2016.1185</u>.

Shannon, C. A., Rospenda, K. M., Richman, J. A., & Minich, L. M. (2009). Race, racial discrimination, and the risk of work-related illness, injury, or assault: Findings from a national study. Journal of Occupational and Environmental Medicine, 51(4), 441-448.

Shepherd-Banigan, M. & Bell, J. F. (2014). Paid leave benefits among a national sample of working mothers with infants in the United States. Maternal and Child Health Journal, 18, 286-295.

SHRM (2017). 2017 employee benefits: Remaining competitive in a challenging talent marketplace. Retrieved from: <u>https://www.shrm.org/hr-today/trends-and-forecasting/research-and-surveys/Documents/2017-Employee-Benefits.pdf</u>.

Sinclair, R., Sears, L. E., Probst, T., & Zajack, M. (2010). A multilevel model of economic stress and employee wellbeing. Contemporary occupational health psychology: Global perspectives on research and practice, 1, 1-20.

Silverstein, M. (2008) Meeting the Challenges of an Aging Workforce. American Journal of Industrial Medicine, 51, 269-280. Dol: 10.1002/ajim.20569.

Smith, T. D. & DeJoy, D. M. (2012). Occupational injury in America: An analysis of risk factors using data from the General Social Survey (GSS). Journal of Safety Research, 43, 67-74.

Sorensen, G., Stoddard, A., Quintiliani, L., Ebbeling, C., Nagler, E., Yang, M., ... & Wallace, L. (2010). Tobacco use cessation and weight management among motor freight workers: Results of the gear up for health study. Cancer Causes & Control, 21, 2113-2122.

Stewart, W. F., Ricci, J. A., Chee, E., et al. (2003). Cost of lost productive work time among US workers with depression. Journal of the American Medical Association, 289: 3135-3144.

Straif, K., Baan, R., Grosse, Y., Secretan, B., Ghissassi, F. R., Bouvard, V., Altieri, A., Benbrahim-Tallaa, L., & Cogliano, V. (2007). WHO International Agency for Research on Cancer Monograph Working Group . Carcinogenicity of shift-work, painting, and fire-fighting. Lancet Oncology, 8, 1065-1066.

Sturm, R., Ringel, J. S., & Andreyeva, T. (2004). Increasing obesity rates and disability trends. Health Affairs, 23, Beyond Managed Care. Published March/April 2004 <u>https://doi.org/10.1377/hlthaff.23.2.199</u>

Swanberg, J., Watson, E., & Eastman, M. (2014). Scheduling Challenges among Workers in Low-wage Hourly Jobs: Similarities and Differences among Workers in Standard and Non-Standard Hour Jobs. Community, Work & Family. <u>http://dx.doi.org/10.1080/13668803.2014.931837</u>.

Tarafdar, M., D'Arcy, J., Turel, O., & Gupta, A. (2015). The dark side of information technology. MIT Sloan Management Review, 56, 61-70.

Theorell, T., Jood, K., Jarvholm, L. S., Vingard, E., Perk, J., Ostergren, P. O., & Hall, C. (2016). A systematic review of studies in the contributions of the work environment to ischemic heart disease development. European Journal of Public Health 26: 470-477.

The White House (2017). "The Opioid Crisis". <u>https://www.whitehouse.gov/opioids/</u>. Accessed August 1, 2018.

Toossi, M. (2016, September). A look at the future of the U.S. labor force to 2060. Spotlight on Statistics, Bureau of Labor Statistics.

Toossi, M. & Torpey, E. (2017, May). Older workers: Labor force trends and career options. Career Outlook, U.S. Bureau of Labor Statistics.

Toossi, M. & Morisi, T.L. (2017) <u>https://www.bls.gov/spotlight/2017/women-in-the-workforce-before-during-and-after-the-great-recession/pdf/women-in-the-workforce-before-during-and-after-the-great-recession.pdf</u>

Turner, N., Hershcovis, M. S., Reich, T. C., & Totterdell, P. (2014). Work–family interference, psychological distress, and workplace injuries. Journal of Occupational and Organizational Psychology, 87, 715-732.

U.S. Census Bureau. (2015). Nation Gains more than 4 Million Non-employer Businesses Over the Last Decade, Census Bureau Reports, U.S. Census Bureau Newsroom (blog), May 27, http://www.census.gov/newsroom/press-releases/2015/cb15-96.html.

Vandenberg, R. J., Richardson, H. A., & Eastman, L. J. (1999). The impact of high involvement work processes on organizational effectiveness: A second-order latent variable approach. Group and Organization Management, 24, 300-339.

Vinchur, A. J. & Koppes, L. L. (2011). A historical survey of research and practice in industrial and organizational psychology. In S. Zedeck (Ed.), *APA handbook of industrial and organizational psychology* (Vol. 1, pp. 3–36). Washington, DC: American Psychological Association.

Virtanen, M., Kivimaki, M., Joensuu, M., Virtanen, P., Elovainio, M., & Vahtera, J. (2005). Temporary employment and health: a review. International Journal of Epidemiology, 34 (3): 610-622.

Wells, K. B., Stewart, A., Hays, R. D., Burnam, M.A., Rogers, W., Daniels, M., Berry, S., Greenfield, S., & Ware, J. (1989). The functioning and well-being of depressed patients: Results from the Medical Outcomes Study. Journal of the American Medical Association, 262(7), 914-919.

Williamson, A. M. & Feyer, A. M. (2000). Moderate sleep deprivation produces impairments in cognitive and motor performance equivalent to legally prescribed levels of alcohol intoxication. Occupational & Environmental Medicine, 57, 649-655.

Winwood, P. C. & Lushington, K. (2006). Disentangling the effects of psychological and physical work demands on sleep, recovery and maladaptive chronic stress outcomes within a large sample of Australian nurses. Journal of Advanced Nursing, 56, 679-689.