

Appendix F: Selected References of Descriptive NHANES Papers on Diet- and-Nutrition Biochemical Indicators

Overarching

Haynes MBH, Pfeiffer CM, Sternberg MR, Schleicher RL. Selected pre-analytical and physiological factors are weakly to moderately associated with twenty-nine biomarkers of diet and nutrition, NHANES 2003–2006. *J Nutr.* 2013;143:1001S–1010S.

Pfeiffer CM, Schleicher RL, Johnson CL, Coates PM. Assessing vitamin status in large population surveys by measuring biomarkers and dietary intake – two case studies: folate and vitamin D. *Food Nutr Res.* 2012;56:5944.

Pfeiffer CM, Sternberg MR, Schleicher RL, Haynes MBH, Rybak ME, Pirkle JL. The CDC’s Second National Report on Biochemical Indicators of Diet and Nutrition in the US Population is a valuable tool for researchers and policy makers to assess nutritional status. *J Nutr.* 2013;143:938S–947S.

Pfeiffer CM, Lacher DA, Schleicher RL, Johnson CL, Yetley EA. Challenges and lessons learned in generating and interpreting NHANES nutritional biomarker data. *Adv Nutr.* 2017;8:290–307.

Pfeiffer CM, Sternberg MR, Powers CD, Ogden CL, Jefferds MED, Crider KS, Juan W, Gahche JJ, Rybak ME, Fazili Z, et al. Centers for Disease Control and Prevention’s 2026 Nutrition Report presents data for 131 biomarkers from the cross-sectional National Health and Nutrition Examination Survey (NHANES 1999–August 2023). *Curr Dev Nutr.* 2026a;10:107688.

Pfeiffer CM, Sternberg MR, Powers CD, Ogden CL, Jefferds MED, Crider KS, Juan W, Gahche JJ, Rybak ME, Fazili Z, et al. Population-level changes in folate, iodine, and vitamin D status over more than 20 years: biomarker data from the cross-sectional National Health and Nutrition Examination Survey (NHANES 1999–2000 through August 2021–August 2023). *Curr Dev Nutr.* 2026b;10:107687.

Pfeiffer CM, Sternberg MR, Powers CD, Fothergill A, Juan W, Jefferds MED, Wambogo E, Leachman J, Mineva EM, Rybak ME, et al. The prevalence of nutritional imbalances in the US population over time: Biomarker data from the cross-sectional National Health and Nutrition Examination Survey (NHANES 1999–2000 through August 2021–August 2023). *Curr Dev Nutr.* 2026c;10:107689.

Sternberg MR, Schleicher RL, Pfeiffer CM. Regression modeling plan for twenty-nine biochemical indicators of diet and nutrition measured in NHANES 2003–2006. *J Nutr.* 2013; 143:948S–956S.

Sternberg MR, Pfeiffer CM, Powers CD, Ahluwalia N, Hamner H, Wang A, Wambogo E, Gahche JJ, Juan W. Dietary supplement use explains more variance in nutritional biomarker concentrations than 24-hour dietary intake from food only: evidence from a cross-sectional survey of the United States population (NHANES 2003–2006). *Curr Dev Nutr.* 2026;10:107692.

Water-Soluble Vitamins

Bailey RL, Mills JL, Yetley EA, Gahche JJ, Pfeiffer CM, Dwyer JT, et al. Unmetabolized serum folic acid and its relation to folic acid intake from diet and supplements in a nationally representative sample of adults aged ≥ 60 y in the United States. *Am J Clin Nutr.* 2010;92:383–389.

Bailey RL, Carmel R, Green R, Pfeiffer CM, Cogswell ME, Osterloh JD, Sempos CT, Yetley EA. Monitoring of vitamin B12 nutritional status in the United States using plasma methylmalonic acid and serum vitamin B12. *Am J Clin Nutr.* 2011;94:552–561.

Bailey RL, Mills JL, Yetley EA, Gahche JJ, Pfeiffer CM, Dwyer JT, Dodd KW, Sempos CT, Betz JM, Picciano MF. Serum unmetabolized folic acid in a nationally representative sample of adults aged ≥ 60 y in the United States, 2001–2002. *Food Nutr Res.* 2012;56:5616.

Bailey RL, Durazo-Arvizu RA, Carmel R, Green R, Pfeiffer CM, Sempos CT, Carriquiry A, Yetley EA. Modeling a methylmalonic acid-derived change point for serum vitamin B-12 for adults in the National Health and Nutrition Examination Survey S. *Am J Clin Nutr.* 2013;98:460–467.

Bailey RL, Fulgoni VL, Taylor CL, Pfeiffer CM, Thuppal SV, McCabe GP, Yetley EA. Correspondence of folate dietary intake and biomarker data. *Am J Clin Nutr.* 2017;105:1336–1343.

Boulet SL, Yang Q, Mai C, Mulinare J, Pfeiffer CM. Folate status in women of childbearing age by race/ethnicity—United States 1999–2000, 2001–2002, and 2003–2004. *Morb Mortal Wkly Rep.* 2007;55(51):1377–1380.

Crider KS, Adisa O, Pfeiffer CM, Wang A, Zhou Y, Yeung LF, Bullard KM, Qi YP, Rose C, Fazili Z, et al. Prediabetes, diabetes, and folate status among United States women of reproductive age: NHANES 2011–March 2020. *Am J Clin Nutr.* 2026;123:101193.

Fazili Z, Sternberg MR, Potischman N, Wang C-Y, Storandt RJ, Yeung L, Yamini S, Gahche JJ, Juan WY, Qi YP, Paladugula N, Gabey G, Pfeiffer CM. Demographic, physiologic, and lifestyle characteristics observed with serum total folate differ among folate forms: cross-sectional data from fasting samples in the NHANES 2011–2016. *J Nutr.* 2020;150:851–860.

Hamner HC, Mulinare J, Cogswell ME, Flores AL, Boyle CA, Prue CE, et al. Predicted contribution of folic acid fortification of corn masa flour to the usual folic acid intake for the US population: National Health and Nutrition Examination Survey, 2001–2004. *Am J Clin Nutr.* 2009;89:305–315.

Mineva EM, Sternberg MR, Zhang M, Aoki Y, Storandt R, Bailey RL, Pfeiffer CM. Age-specific reference ranges are needed to interpret serum methylmalonic acid concentrations in the US population. *Am J Clin Nutr.* 2019;110:158–168.

Mineva E, Sternberg M, Bailey R, Storandt R, Pfeiffer CM. Fewer US adults had low or transitional vitamin B12 status based on the novel combined indicator of vitamin B12 status compared with single, conventional biomarkers, NHANES 1999–2004. *Am J Clin Nutr.* 2021;114:1070–1079.

Mineva EM, Sternberg MR, Pfeiffer CM. The diagnostic accuracy of methylmalonic acid to detect inadequate vitamin B12 status relative to 3cB-12 was higher compared with that of total cobalamin and total homocysteine in a cross-sectional survey of apparently healthy U.S. adults. *Curr Dev Nutr.* 2026;10:107704.

Pfeiffer CM, Caudill SP, Gunter EW, Osterloh J, Sampson EJ. Biochemical indicators of B vitamin status in the U.S. population after folic acid fortification: results from the National Health and Nutrition Examination Survey 1999–2000. *Am J Clin Nutr.* 2005;82:442–450.

Pfeiffer CM, Johnson CL, Jain RB, Yetley EA, Picciano MF, Rader JI, et al. Trends in blood folate and vitamin B12 concentrations in the United States, 1988–2004. *Am J Clin Nutr.* 2007;86:718–727.

Pfeiffer CM, Osterloh JD, Kennedy-Stephenson J, Picciano MF, Yetley EA, Rader JI, et al. Trends in circulating concentrations of total homocysteine among US adolescents and adults: findings from the 1991–1994 and 1999–2004 National Health and Nutrition Examination Surveys. *Clin Chem.* 2008;54:801–813.

Pfeiffer CM, Hughes JP, Lacher DA, Bailey RL, Berry RJ, Zhang M, Yetley EA, Rader JI, Sempos CT, Johnson CL. Estimation of trends in serum and RBC folate in the U.S. population from pre- to postfortification using assay-adjusted data from the NHANES 1988–2010. *J Nutr.* 2012;142:886–893.

Pfeiffer CM, Hughes JP, Durazo-Arvizu RA, Lacher DA, Sempos CT, Zhang M, Yetley EA, Johnson CL. Changes in measurement procedure from a radioassay to a microbiologic assay necessitates adjustment of serum and RBC folate concentrations in the U.S. population from the NHANES 1988–2010. *J Nutr.* 2012; 142:894–900.

Pfeiffer CM, Sternberg MR, Schleicher RL, Rybak ME. Dietary supplement use and smoking were strongly related to biomarkers of water-soluble vitamin status after adjusting for socioeconomic and lifestyle factors in a representative sample of US adults. *J Nutr.* 2013;143:957S–965S.

Pfeiffer CM, Sternberg MR, Fazili Z, Yetley EA, Lacher DA, Bailey RL, Johnson CL. Unmetabolized folic acid is detected in nearly all serum samples from US children, adolescents, and adults. *J Nutr.* 2015;145:520–531.

Pfeiffer CM, Sternberg MR, Fazili Z, Lacher DA, Zhang M, Johnson CL, Hamner HC, Bailey RL, Rader JI, Yamini S, Berry RJ, and Yetley EA. Folate status and concentrations of serum folate forms in the US population: National Health and Nutrition Examination Survey 2011–2. *Br J Nutr.* 2015;113:1965–1977.

Pfeiffer CM, Sternberg MR, Zhang M, Fazili Z, Storandt RJ, Crider KS, Yamini S, Gahche JJ, Juan WY, Wang C-Y, Potischman N, Williams J, LaVoie DJ. Folate status in the US population 20 years after the introduction of folic acid fortification. *Am J Clin Nutr.* 2019;110(5):1088–1097.

Powers CD, Sternberg MR, Patel SB, Pfeiffer CM, Storandt RJ, Schleicher RL. Serum vitamin C status in US adults: NHANES 2017–2018. *J Appl Lab Med.* 2023;08(02):272–284.

Qi YP, Do AN, Hamner HC, Pfeiffer CM, Berry RJ. The prevalence of low serum vitamin B-12 status in the absence of anemia or macrocytosis did not increase among older U.S. adults after mandatory folic acid fortification. *J Nutr.* 2014;144:170–176.

Schleicher RL, Carroll MD, Ford ES, Lacher DA. Serum vitamin C and the prevalence of vitamin C deficiency in the United States: 2003–2004 National Health and Nutrition Examination Survey (NHANES). *Am J Clin Nutr.* 2009;90:1252–1263.

Selhub J, Jacques PF, Rosenberg IH, Rogers G, Bowman BA, Gunter EW. Serum total homocysteine concentrations in the Third National Health and Nutrition Examination Survey (1991–1994): population reference ranges and contribution of vitamin status to high serum concentrations. *Ann Intern Med.* 1999;131:331–339.

Tinker SC, Hamner HC, Berry RJ, Bailey LB, Pfeiffer CM. Does obesity modify the association of supplemental folic acid with folate status among nonpregnant women of childbearing age in the United States? *Birth Defect Res A Clin Mol Teratol.* 2012;94(10):749–755.

U.S. Centers for Disease Control and Prevention. Folate status in women of childbearing age, by race/ethnicity—United States, 1999–2000. *Morb Mortal Wkly Rep.* 2002;51:808–810.

Wang A, Rose CE, Qi YP, Williams JL, Pfeiffer CM, Crider KS. Impact of voluntary fortification of corn masa flour on RBC folate concentrations in the U.S. (NHANES 2011–2018). *Nutrients*. 2021;13:1325.

Wang A, Yeung LF, Burrows NR, Rose CE, Fazili Z, Pfeiffer CM, Crider KS. Reduced kidney function is associated with increasing red blood cell folate concentration and changes in folate form distributions (NHANES 2011–2018). *Nutrients*. 2022;14:1054.

Wang A, Fothergill A, Yeung LF, Crider KS, Williams JL. Update on the impact of voluntary folic acid fortification of corn masa flour on red blood cell folate concentrations—National Health and Nutrition Examination Survey, 2011–March 2020. *Birth Def Res*. 2024;116:e2321.

Wang A, Zauche LH, Crider KS, Mai CT, Qi YP, Yeung LF, Williams JL. Trends and prevalence of modifiable risk factors for birth defects among U.S. women of reproductive age: National Health and Nutrition Examination Survey 2007 to March 2020. *Am J Prev Med*. 2025;69:107947.

Yang Q-H, Carter HK, Mulinare J, Berry RJ, Friedman JM, Erickson JD. Race-ethnicity differences in folic acid intake in women of childbearing age in the United States after folic acid fortification: findings from the National Health and Nutrition Examination Survey, 2001–2002. *Am J Clin Nutr*. 2007;85:1409–1416.

Yang Q-H, Botto LD, Gallagher M, Friedman JM, Sanders CL, Koontz D, et al. Prevalence and effects of gene-gene and gene-nutrient interactions on serum folate and serum total homocysteine concentrations in the United States: findings from the third National Health and Nutrition Examination Survey DNA Bank. *Am J Clin Nutr*. 2008;88:232–246.

Yang Q, Cogswell ME, Hamner HC, Carriquiri A, Bailey LB, Pfeiffer CM, et al. Folic acid source, usual intake, and folate and vitamin B12 status in US adults: National Health and Nutrition Examination Survey (NHANES) 2003–2006. *Am J Clin Nutr*. 2010;91:64–72.

Yetley EA, Pfeiffer CM, Phinney KW, Fazili Z, Lacher DA, Bailey RL, Blackmore S, et al. Biomarkers of folate status in the National Health and Nutrition Examination Survey (NHANES): a roundtable summary. *Am J Clin Nutr*. 2011; 94:303S–312S.

Yetley EA, Pfeiffer CM, Phinney KW, Bailey RL, Blackmore S, Bock JL, Brody LC, et al. Biomarkers of vitamin B-12 status in the National Health and Nutrition Examination Survey (NHANES): a roundtable summary. *Am J Clin Nutr*. 2011;94:313S–321S.

Yeung L, Yang Q, Berry RJ. Contributions of total daily intake of folic acid to serum folate concentrations. *J Am Med Assoc*. 2008;300:2486–2487.

Yeung LF, Cogswell ME, Carriquiry AL, Bailey LB, Pfeiffer CM, Berry RJ. Contributions of enriched cereal-grain products ready-to-eat cereals and supplements to folic acid and vitamin B-12 usual intake and folate and vitamin B-12 status in US children: National Health and Nutrition Examination Survey (NHANES) 2003–2006. *Am J Clin Nutr.* 2011;93:172–185.

Zhang M, Sternberg MR, Yeung LF, Pfeiffer CM. Population RBC folate concentrations can be accurately estimated from measured whole blood folate, measured hemoglobin, and predicted serum folate, National Health and Nutrition Examination Survey 1988-2010. *Am J Clin Nutr.* 2020;111:601–612.

Zhou Y, Wang A, Yeung LF, Qi YP, Pfeiffer CM, Crider KS. Folate and vitamin B12 usual intake and biomarker status by intake source in U.S. adults (≥ 19 y): National Health and Nutrition Examination Survey (NHANES) 2007-2018. *Am J Clin Nutr.* 2023;118(1):241–254.

Fat-Soluble Vitamins and Nutrients

Couch CA, Williams AM, Stierman B, Storandt R, Leachman J, Mishra S, Juan W, Gahche JJ, Wambogo E, Mineva E, Pfeiffer CM, Ogden CL. Trends in vitamin D status in the United States, 2007-2023: a cross-sectional analysis of data from the National Health and Nutrition Examination Survey. *Am J Clin Nutr.* 2026;123:101078.

Ford ES, Liu S, Mannino DM, Giles WH, Smith SJ. C-reactive protein concentration and concentrations of blood vitamins, carotenoids, and selenium among United States adults. *Eur J Clin Nutr.* 2003;57:1157–1163.

Ford ES, Ajani UA, Mokdad AH. Brief communication: the prevalence of high intake of vitamin E from the use of supplements among U.S. adults. *Ann Intern Med.* 2005;143:143–145.

Ford ES, Schleicher RL, Mokdad AH, Ajani UA, Liu S. Distribution of serum concentrations of α -tocopherol and γ -tocopherol in the U.S. population. *Am J Clin Nutr.* 2006;84:375–383.

Gillespie C, Ballew C, Bowman BA, Donehoo R, Serdula MK. Intraindividual variation in serum retinol concentrations among participants in the third National Health and Nutrition Examination Survey, 1988–1994. *Am J Clin Nutr.* 2004;79:625–632.

Herrick KA, Storandt RJ, Afful J, Pfeiffer CM, Schleicher RL, Gahche JJ, Potischman N. Vitamin D status in the United States, 2011–2014. *Am J Clin Nutr.* 2019;110:150–157.

Looker AC, Dawson-Hughes B, Calvo MS, Gunter EW, Sahyoun NR. Serum 25-hydroxyvitamin D status of adolescents and adults in two seasonal subpopulations from NHANES III. *Bone.* 2002;30:771–777.

Looker AC, Pfeiffer CM, Lacher DA, Schleicher RL, Picciano MF, Yetley EA. Serum 25-hydroxyvitamin D status of the US population: 1988–1994 versus 2000–2004. *Am J Clin Nutr.* 2008;88:1519–1527.

Looker AC, Lacher DA, Pfeiffer CM, Schleicher RL, Picciano MF, Yetley EA. Data advisory with regard to NHANES serum 25-hydroxyvitamin D data. *Am J Clin Nutr.* 2009;90:695.

Nesby-O'Dell S, Scanlon KS, Cogswell ME, Gillespie C, Hollis BW, Looker AC, et al. Hypovitaminosis D prevalence and determinants among African American and white women of reproductive age: Third National Health and Nutrition Examination Survey, 1988–1994. *Am J Clin Nutr.* 2002;76:187–192.

Powers CD, Sternberg MR, Mineva EM, Clifton CC, Leachman J, Wambogo E, Pfeiffer CM. Over half of United States population had an undesirably low omega-3 index based on erythrocyte membrane measurements: results from the cross-sectional NHANES from August 2021 to August 2023. *Curr Dev Nutr.* 2026;10:107715.

Schleicher RL, Sternberg MR, Pfeiffer CM. Race-ethnicity is a strong correlate of circulating fat-soluble nutrient concentrations in a representative sample of the U.S. population. *J Nutr.* 2013;143:966S–976S.

Schleicher RL, Sternberg MR, Looker AC, Yetley EA, Lacher DA, Sempos CT, Taylor CL, Durazo-Arvizu RA, Maw KL, Chaudhary-Webb M, Johnson CL, Pfeiffer CM. National estimates of serum total 25-hydroxyvitamin D and metabolite concentrations measured by liquid chromatography-tandem mass spectrometry in the U.S. population during 2007–2010. *J Nutr.* 2016;146:1051–1061.

Schleicher RL, Sternberg MR, Lacher DA, Sempos CT, Looker AC, Durazo-Arvizu RA, Yetley EA, Chaudhary-Webb M, Maw KL, Pfeiffer CM, Johnson CL. The vitamin D status of the U.S. population from 1988 to 2010 using standardized serum concentrations of 25-hydroxyvitamin D shows recent modest increases. *Am J Clin Nutr.* 2016;104:454–461.

Schleicher RL, Sternberg MR, Maw KL, Storandt RJ, Herrick KA, Gahche JJ, Potischman NA, Pfeiffer CM. The relationship between total serum 25-hydroxyvitamin D and dietary and supplemental intake of vitamin D in the adult US population during 2003–2014. *J Nutr.* 2021;151:2446–2454.

Yetley EA, Pfeiffer CM, Schleicher RL, Phinney KW, Lacher DA, Christakos S, Eckfeldt JH, et al. NHANES monitoring of serum 25-hydroxyvitamin D: A roundtable summary. *J Nutr.* 2010;140:2030S–2045S.

Iron Indicators

Cogswell ME, Looker AC, Pfeiffer CM, Cook JD, Lacher DA, Beard JL, Lynch SR, et al. Assessment of iron deficiency in US preschool children and nonpregnant females of childbearing age: National Health and Nutrition Examination Survey 2003–2006. *Am J Clin Nutr.* 2009;89:1334–1342.

Jefferds MED, Mei Z, Addo Y, Hamner HC, Perrine CG, Flores-Ayala R, Pfeiffer CM, Sharma AJ. Iron deficiency in the United States: limitations and guidelines, data, and monitoring of disparities. *Am J Pub Health*. 2022;112(S8):S826–S835.

Looker AC, Gunter EW, Cook JD, Green R, Harris JW. Comparing serum ferritin values from different population surveys. National Center for Health Statistics. *Vital Health Stat Series No 2(111)*, 1991.

Looker AC, Gunter EW, Johnson CL. Methods to assess iron status in various NHANES surveys. *Nutr Rev*. 1995;53:246–254.

Looker AC, Dallman PR, Carroll M, Gunter EW, Johnson CL. Prevalence of iron deficiency in the United States. *J Am Med Assoc*. 1997;277:973–975.

Looker AC, Cogswell ME, Gunter EW. Iron deficiency—United States, 1999–2000. *Morb Mortal Wkly Rep*. 2002;51:897–899.

Mei Z, Cogswell ME, Looker AC, Pfeiffer CM, Cusick SE, Lacher DA, et al. Assessment of iron status in US pregnant women from the National Health and Nutrition Examination Survey 1999–2006. *Am J Clin Nutr*. 2011;93:1–9.

Mei Z, Pfeiffer CM, Looker AC, Flores-Ayala RC, Lacher DA, Mirel LB, Grummer-Strawn LM. Serum soluble transferrin receptor concentrations in US preschool children and non-pregnant women of childbearing age from the National Health and Nutrition Examination Survey 2003–2010. *Clin Chim Acta*. 2012;413:1479–1484.

Michels Blanck H, Cogswell ME, Gillespie C, Reyes M. Iron supplement use and iron status among US adults: results from the Third National Health and Nutrition Examination Survey. *Am J Clin Nutr*. 2005;82:1024–1031.

Pfeiffer CM, Sternberg MR, Caldwell KL, Pan Y. Race-ethnicity is related to biomarkers of iron and iodine status after adjusting for sociodemographic and lifestyle factors in a representative sample of US adults. *J Nutr*. 2013;143:977S–985S.

Ramakrishnan U, Frith-Terhune A, Cogswell M, Kettel Khan L. Dietary intake does not account for differences in low iron stores among Mexican American and non-Hispanic white women: Third National Health and Nutrition Examination Survey, 1988–1994. *J Nutr*. 2002;132:996–1001.

Trace Elements

Caldwell KL, Jones R, Hollowell JG. Urinary iodine concentration: United States National Health and Nutrition Examination Survey 2001–2002. *Thyroid*. 2005;15:692–699.

Caldwell KL, Miller GA, Wang RY, Jain RB, Jones RL. Iodine status of the U.S. population, National Health and Nutrition Examination Survey 2003–2004. *Thyroid*. 2008;18:1207–1214.

Caldwell KL, Makhmudov A, Ely E, Jones RL, Wang R. Iodine status of the U.S. population, National Health and Nutrition Examination Survey, 2005–2006 and 2007–2008. *Thyroid*. 2011;21:419–427.

Caldwell KL, Pan Y, Mortensen ME, Makhmudov A, Merrill L, Moye J. Iodine status in pregnant women in the National Children’s study and in U.S. women (15–44 years), National Health and Nutrition Examination Survey 2005–2010. *Thyroid*. 2013;23:927–937.

Hollowell JG, Staehling NW, Hannon WH, Flanders DW, Gunter EW, Maberly GF, et al. Iodine nutrition in the United States. Trends and public health implications: iodine excretion data from National Health and Nutrition Examination Surveys I and III (1971–1974 and 1988–1994). *J Clin Endocrinol Metab*. 1998;83:3401–3408.

Pan Y, Caldwell KL, Li Y, Caudill SP, Mortensen ME, Makhmudov A, Jones RL. Smoothed urinary iodine percentiles for the US population and pregnant women: National Health and Nutrition Examination Survey, 2001–2010. *Eur Thyroid J*. 2013;2(2):127–134.

Perrine CG, Herrick K, Serdula MK, Sullivan KM. Some subgroups of reproductive age women in the United States may be at risk for iodine deficiency. *J Nutr*. 2010;140:1489–1494.

Phytoestrogens

Rybak ME, Sternberg MR, Pfeiffer CM. Sociodemographic and lifestyle variables are compound- and class-specific correlates of urine phytoestrogen concentrations in the US population. *J Nutr*. 2013;143:986S–994S.

U.S. Centers for Disease Control and Prevention. Fourth National Report on Human Exposure to Environmental Chemicals. Atlanta (GA): CDC; 2009. [cited 2011]. Available from: <http://www.cdc.gov/exposurereport>.

Valentin-Blasini L, Blount BC, Caudill SP, Needham LL. Urinary and serum concentrations of seven phytoestrogens in a human reference population subset. *J Expo Anal Environ Epidemiol*. 2003;13:276–282.

Valentin-Blasini L, Sadowski MA, Walden D, Calabiano L, Needham LL, Barr DB. Urinary phytoestrogen concentrations in the U.S. population (1999–2000). *J Expo Anal Environ Epidemiol*. 2005;15:509–523.

Caffeine and Metabolites

Rybak ME, Sternberg MR, Pao C-I, Ahluwalia N, Pfeiffer CM. Urine excretion of caffeine and select caffeine metabolites is common in the US population and associated with caffeine intake. *J Nutr*. 2015;145:766–774.

Acrylamide Hemoglobin Adducts

Vesper HW, Caudill SP, Osterloh JD, Meyers T, Scott D, Myers GL. Exposure of the U.S. population to acrylamide in the National Health and Nutrition Examination Survey 2003–2004. *Environ Health Perspect*. 2010;118:278–283.

Vesper HW, Sternberg MR, Pfeiffer CM. Among ten sociodemographic and lifestyle factors, smoking was most strongly associated with biomarkers of acrylamide exposure in a representative sample of the US population. *J Nutr*. 2013;143:995S-1000S.

Trans Fatty Acids

Vesper HW, Caudill SP, Kuiper HC, Yang Q, Ahluwalia N, Lacher DA, Pirkle JL. Plasma trans-fatty acid concentrations in fasting adults declined from NHANES 1999–2000 to 2009–2010. *Am J Clin Nutr*. 2017;105:1063–1069.