

CDC's Early Childhood Nutrition Report 2025

DATA TO SUPPORT HEALTHY GROWTH AND DEVELOPMENT FOR CHILDREN 5 YEARS AND YOUNGER

This report was first published in June 2025 and updated in August 2025. The update replaced data on four breastfeeding indicators among children born in 2021 with data from children born in 2022, National Immunization Survey-Child.



What is the Early **Childhood Nutrition** Report?

CDC's Early Childhood Nutrition Report contains information on nutrition-related behaviors, practices, and policies that can impact the health and development of children 5 years and younger. This report provides national and state data on:

- Breastfeeding
- Supports for Infant Feeding
- · First Solid Foods
- Healthy Eating
- Affordability of Nutritious Foods

What is already known?

Infancy and early childhood are periods of rapid growth and development, and healthy nutrition plays a vital role. Healthy nutrition supports brain development, strengthens bones, and supports optimal growth and development. In addition, starting healthy nutrition routines early can establish habits that promote health and prevent chronic diseases.

Early feeding decisions, such as breastfeeding, timing of solid food introduction, and what foods and drinks are provided, can play a critical role in early childhood. Policies and practices in places where children and caregivers spend their time can affect these decisions.

What's new?

CDC has published individual reports, such as the Breastfeeding Report Card, Early Care and Education State Indicator Report, Early Care and Education State Licensing Scorecards, and the State Indicator Report on Fruits and Vegetables. But this is the first time that data on several early child nutrition topics are showcased in a single report. This report and its corresponding individual state reports provide a more complete picture of nutrition in early childhood and related practices and supports at the state level.

How can this report be used?

The Early Childhood Nutrition Report provides data for national partners, state public health practitioners, policymakers, and other interested parties to identify gaps and areas for improvement, as well as opportunities for policies and activities that support healthy nutrition in early childhood. The individual state reports include opportunities for action to improve nutrition for young children.





WHAT ARE THE KEY FINDINGS FOR THE NATION?



Breastfeeding

Only 2 in 5 infants born in 2022 were still being breastfed on their first birthday. This means most infants do not receive the full health benefits that breast milk provides.



Supports for Infant Feeding

10 states and the District of Columbia fully met, and 38 states partially met, the standard to encourage and support breastfeeding in their licensing regulations for early care and education centers in 2023. Meeting, or partially meeting, the standard can help infants receive breast milk for longer.



First Solid Foods

1 in 10 infants born during 2019–2021 was given foods other than breast milk or infant formula before they were 4 months old. This is too early.



Healthy Eating

Only 49% of children 1–5 years in 2022–2023 ate vegetables at least once daily. Eating a variety of vegetables daily is needed to support this critical time of growth and development.

More than half of children 1–5 years in 2022–2023 drank a sugary drink one or more times weekly. Sugary drinks are the greatest source of added sugars in this age group and are not needed.



Affordability of Nutritious Foods

69% of children 1–5 years lived in households that were always able to afford nutritious foods in 2022–2023. It is important for all families to be able to afford healthy foods, so they can make healthier choices.

Breastfeeding

Breastfeeding is the best source of nutrition for most infants. It helps protect babies and moms from some illnesses and diseases. It is recommended that infants are fed only breast milk for about the first 6 months. Infants can be introduced to solid foods at about 6 months and can continue to be fed breast milk for 1 to 2 years, or longer. Many mothers stop breastfeeding earlier than they planned. Breastfed infants who receive infant formula within 2 days of birth typically do not breastfeed for as long as those who do not. The table below provides data on breastfeeding rates.



| State | Ever Breastfed, 2022 (%) ^a | Breastfed at 12 Months, 2022 (%) ^a | Exclusive Breastfeeding Through 6 Months, 2022 (%) ^a | Breastfed Infants Fed Formula Before 2 Days, 2022 (%) ^a |
|----------------------|--|--|--|--|
| National | 85.7 | 40.8 | 27.9 | 22.9 |
| Alabama | 77.9 | 34.8 | 26.4 | 22.7 |
| Alaska | 91.5 | 48.5 | 31.0 | 18.7 |
| Arizona | 86.2 | 36.4 | 22.4 | 26.9 |
| Arkansas | 83.4 | 30.8 | 25.1 | 19.7 |
| California | 90.1 | 41.5 | 28.6 | 20.5 |
| Colorado | 91.4 | 53.4 | 34.7 | 16.2 |
| Connecticut | 91.1 | 48.1 | 25.4 | 32.0 |
| Delaware | 80.2 | 35.7 | 24.0 | 25.2 |
| District of Columbia | 90.2 | 43.8 | 30.1 | 24.0 |
| Florida | 86.6 | 36.7 | 25.9 | 25.8 |
| Georgia | 81.1 | 35.0 | 19.5 | 26.5 |
| Hawaii | 90.7 | 53.5 | 33.8 | 22.0 |
| Idaho | 90.7 | 45.3 | 33.1 | 16.5 |
| Illinois | 86.0 | 41.7 | 29.2 | 23.2 |
| Indiana | 86.0 | 41.0 | 31.9 | 14.8 |
| Iowa | 87.5 | 44.7 | 33.3 | 19.7 |
| Kansas | 87.8 | 42.3 | 28.4 | 21.5 |
| Kentucky | 73.9 | 33.3 | 24.1 | 18.5 |
| Louisiana | 72.4 | 29.6 | 27.4 | 19.3 |
| Maine | 89.4 | 46.3 | 30.0 | 14.8 |
| Maryland | 90.8 | 51.0 | 35.7 | 21.2 |
| Massachusetts | 87.9 | 46.3 | 25.1 | 21.9 |
| Michigan | 81.2 | 43.5 | 29.3 | 24.5 |
| Minnesota | 92.0 | 47.0 | 32.8 | 16.2 |
| Mississippi | 72.5 | 30.9 | 24.0 | 20.3 |
| Missouri | 89.4 | 36.0 | 26.6 | 25.5 |
| Montana | 91.9 | 41.2 | 33.9 | 14.2 |
| | | | | |

| State | Ever Breastfed, 2022 (%) ^a | Breastfed at 12 Months, 2022 (%) ^a | Exclusive Breastfeeding Through 6 Months, 2022 (%) ^a | Breastfed Infants Fed Formula Before 2 Days, 2022 (%) ^a |
|----------------|--|--|--|--|
| Nebraska | 91.2 | 53.2 | 36.7 | 13.9 |
| Nevada | 82.6 | 37.1 | 23.9 | 29.2 |
| New Hampshire | 89.7 | 49.7 | 35.3 | 13.0 |
| New Jersey | 88.5 | 48.0 | 26.1 | 26.3 |
| New Mexico | 87.4 | 43.4 | 35.3 | 21.3 |
| New York | 88.4 | 44.3 | 26.4 | 30.3 |
| North Carolina | 84.6 | 40.5 | 30.8 | 19.1 |
| North Dakota | 86.2 | 49.7 | 31.3 | 23.9 |
| Ohio | 84.2 | 38.1 | 28.0 | 19.3 |
| Oklahoma | 81.6 | 35.7 | 28.0 | 20.7 |
| Oregon | 91.4 | 43.1 | 35.9 | 17.2 |
| Pennsylvania | 85.0 | 38.4 | 26.0 | 25.3 |
| Rhode Island | 86.0 | 43.1 | 25.3 | 28.1 |
| South Carolina | 78.6 | 33.0 | 22.7 | 24.2 |
| South Dakota | 89.4 | 44.0 | 30.3 | 16.7 |
| Tennessee | 78.5 | 33.7 | 26.9 | 19.6 |
| Texas | 83.4 | 37.5 | 26.4 | 27.6 |
| Utah | 94.2 | 56.8 | 31.1 | 20.0 |
| Vermont | 95.4 | 55.0 | 35.7 | 14.4 |
| Virginia | 85.5 | 42.9 | 29.9 | 27.4 |
| Washington | 87.9 | 46.2 | 32.8 | 17.3 |
| West Virginia | 71.7 | 30.8 | 19.6 | 24.1 |
| Wisconsin | 88.8 | 48.7 | 30.7 | 16.9 |
| Wyoming | 89.4 | 48.4 | 36.3 | 12.8 |

^a Among children born in 2022. Breastfeeding indicators are the percentage of infants breastfeeding at the specified time points, calculated among all infants. The rate for infants receiving formula before 2 days of age is calculated among breastfed infants.

Supports for Infant Feeding

Practices and policies can impact how infants and children are fed in places such as hospitals and early care and education (ECE) centers. The table below provides data on maternity care practices in hospitals, ECE licensing standards for breastfeeding support, and state paid family and medical leave legislation. CDC tracks maternity care practices that support optimal infant feeding through the Maternity Practices in Infant Nutrition and Care (mPINC) survey. State ECE licensing regulations and state paid family and medical leave programs can be used to support infant feeding after the birth hospitalization.



| State | Total mPINC Score, 2024 (Out of 100) ^a | Weeks of Paid Family and Medical Leave Available for the Care of a New Child, 2025 ^b | Encourage and Support Breastfeeding: State Center- Based ECE Licensing Standard, 2023 ^d |
|----------------------|---|---|---|
| National | 82 | - - | - |
| Alabama | 76 | - | Partially met |
| Alaska | 77 | - | Partially met |
| Arizona | 76 | - | Partially met |
| Arkansas | 71 | - | Partially met |
| California | 89 | 8 weeks | Fully met |
| Colorado | 86 | 12 weeks | Partially met |
| Connecticut | 88 | 12 weeks | Partially met |
| Delaware | 93 | 12 weeks ^c | Fully met |
| District of Columbia | - | 12 weeks | Fully met |
| Florida | 85 | - | Partially met |
| Georgia | 78 | - | Fully met |
| Hawaii | 78 | - | Partially met |
| Idaho | 79 | - | Did not address |
| Illinois | 81 | - | Fully met |
| Indiana | 83 | - | Partially met |
| Iowa | 76 | - | Partially met |
| Kansas | 81 | - | Partially met |
| Kentucky | 74 | - | Partially met |
| Louisiana | 85 | - | Fully met |
| Maine | 82 | 12 weeks ^c | Partially met |
| Maryland | 83 | 12 weeks ^c | Fully met |
| Massachusetts | 89 | 12 weeks | Partially met |
| Michigan | 79 | - | Partially met |
| Minnesota | 81 | 12 weeks ^c | Partially met |
| Mississippi | 87 | - | Fully met |
| Missouri | 77 | - | Did not address |
| Montana | 84 | - | Partially met |
| | | | |

| State | Total mPINC Score, 2024 (Out of 100) ^a | Weeks of Paid Family and Medical Leave Available for the Care of a New Child, 2025 b | Encourage and Support Breastfeeding: State Center- Based ECE Licensing Standard, 2023 ^d |
|----------------|---|--|---|
| Nebraska | 75 | - | Partially met |
| Nevada | 76 | - | Partially met |
| New Hampshire | 90 | - | Partially met |
| New Jersey | 84 | 12 weeks | Partially met |
| New Mexico | 88 | - | Partially met |
| New York | 84 | 12 weeks | Partially met |
| North Carolina | 84 | - | Fully met |
| North Dakota | 78 | - | Partially met |
| Ohio | 85 | - | Partially met |
| Oklahoma | 79 | - | Partially met |
| Oregon | 87 | 12 weeks | Partially met |
| Pennsylvania | 83 | - | Partially met |
| Rhode Island | 87 | 7 weeks | Partially met |
| South Carolina | 80 | - | Partially met |
| South Dakota | 75 | - | Partially met |
| Tennessee | 76 | - | Partially met |
| Texas | 81 | - | Fully met |
| Utah | 78 | - | Partially met |
| Vermont | 90 | - | Fully met |
| Virginia | 82 | - | Partially met |
| Washington | 86 | 12 weeks | Partially met |
| West Virginia | 76 | - | Partially met |
| Wisconsin | 85 | - | Partially met |
| Wyoming | 73 | - | Partially met |

^a Possible mPINC scores are 0 to 100, with higher scores indicating better maternity care practices and policies. Scores are not reported for District of Columbia because of small sample sizes but are included in the US national total mPINC score.

^b Among states with enacted legislation, the number of weeks presented are those that can be claimed by eligible employees for the care of a new child by birth, adoption, or foster care. Weeks may also be used for other family and medical leave events as specified by the state (e.g., a serious health condition of a qualified family member); in some states, additional weeks of benefits may be available for other needs. Employees must meet eligibility requirements to claim benefits; requirements vary across state programs. Voluntary paid family and medical leave programs in New Hampshire and Vermont are not included.

State has enacted legislation to create a paid family and medical leave insurance program, but the program is not currently paying benefits. Paid leave benefits for eligible employees in Delaware, Maine, Maryland, and Minnesota will be available in 2026.

^d States can include high-impact obesity prevention standards in their ECE licensing and administrative regulations. Family child care homes are not included. This indicator reports whether states fully met, partially met, did not address, or contradicted, the standard Encourage and Support Breastfeeding: Encourage and support breastfeeding and feeding of breast milk by making arrangements for mothers to feed their children on-site.

First Solid Foods

It is recommended that infants be introduced to foods other than breast milk or infant formula when they are about 6 months old. Solid foods can provide important nutrients to support growth and development. However, introducing solid foods before 4 months of age is too early because children might not be developmentally ready, and it could increase their risk of overweight and obesity. The table below provides data on the percentage of children introduced to solid foods too early and the related state ECE licensing regulations, which can support appropriate timing of first solid foods.



| State | Early Solid Foods, 2019–2021 (%) ^a | Timely Introduction of Solid Foods: State Center-Based ECE Licensing Standard, 2023 ^b |
|----------------------|---|---|
| National | 10.1 | - |
| Alabama | 13.4 | Did not address |
| Alaska | 11.4 | Fully met |
| Arizona | 11.8 | Did not address |
| Arkansas | 11.3 | Fully met |
| California | 11.0 | Fully met |
| Colorado | 4.8 | Fully met |
| Connecticut | 7.0 | Fully met |
| Delaware | 9.5 | Fully met |
| District of Columbia | 10.7 | Fully met |
| Florida | 7.8 | Fully met |
| Georgia | 10.1 | Fully met |
| Hawaii | 3.8 | Fully met |
| Idaho | 6.3 | Did not address |
| Illinois | 13.3 | Fully met |
| Indiana | 13.4 | Did not address |
| Iowa | 5.9 | Fully met |
| Kansas | 8.9 | Did not address |
| Kentucky | 11.3 | Did not address |
| Louisiana | 10.3 | Fully met |
| Maine | 6.3 | Did not address |
| Maryland | 8.3 | Fully met |
| Massachusetts | 10.4 | Did not address |
| Michigan | 7.2 | Fully met |
| Minnesota | 7.2 | Fully met |
| Mississippi | 12.5 | Partially met |
| Missouri | 11.9 | Did not address |
| Montana | 9.7 | Fully met |
| | | |

| State | Early Solid Foods, 2019–2021 (%) ^a | Timely Introduction of Solid Foods: State Center-Based ECE Licensing Standard, 2023 ^b |
|----------------|---|---|
| Nebraska | 9.2 | Fully met |
| Nevada | 13.3 | Fully met |
| New Hampshire | 7.3 | Fully met |
| New Jersey | 6.2 | Fully met |
| New Mexico | 11.9 | Fully met |
| New York | 10.4 | Fully met |
| North Carolina | 12.4 | Fully met |
| North Dakota | 6.5 | Partially met |
| Ohio | 10.0 | Partially met |
| Oklahoma | 8.7 | Did not address |
| Oregon | 10.5 | Contradicted |
| Pennsylvania | 10.3 | Did not address |
| Rhode Island | 14.4 | Fully met |
| South Carolina | 9.8 | Fully met |
| South Dakota | 6.5 | Did not address |
| Tennessee | 15.6 | Fully met |
| Texas | 11.7 | Fully met |
| Utah | 9.0 | Fully met |
| Vermont | 10.5 | Fully met |
| Virginia | 11.9 | Fully met |
| Washington | 5.7 | Fully met |
| West Virginia | 13.3 | Fully met |
| Wisconsin | 6.1 | Fully met |
| Wyoming | 6.8 | Did not address |

^a Data include children ages 1–3 years born in 2019–2021. Introducing foods before 4 months of age was considered early solid food introduction.

^b States can include high-impact obesity prevention standards in their ECE licensing and administrative regulations. Family child care homes are not included. This indicator reports whether states fully met, partially met, did not address, or contradicted the standard Timely Introduction of Solid Foods: Introduce age-appropriate solid foods no sooner than 4 months of age, and preferably around 6 months of age.

Healthy Eating

Once children start eating solid foods, they need a variety of foods from different food groups to meet their nutrient needs. This includes fruits and vegetables. Children younger than 2 years should also avoid foods and beverages with added sugars. Children 2 years and older should limit calories from added sugars. The table below provides data on the percentage of children who ate fruits and vegetables daily and drank a sugary drink one or more times weekly, as well as related state ECE licensing regulations, which can support healthy eating.



| State | Ate Fruit at Least Once Daily, 2022–2023 (%) ^a | Serve Fruits of Several Varieties: State Center-Based ECE Licensing Standard, 2023 b | Ate Vegetables at Least Once Daily, 2022–2023 (%) a | Serve Vegetables: State Center-Based ECE Licensing Standard, 2023 b | Drank a Sugary Drink One or More Times Weekly, 2022–2023 (%) a | Avoid Sugar: State Center- Based ECE Licensing Standard, 2023 b |
|----------------------|---|---|---|---|---|---|
| National | 66.6 | - | 49.1 | - | 57.0 | - |
| Alabama | 60.1 | Partially met | 44.4 | Partially met | 67.1 | Did not address |
| Alaska | 70.5 | Partially met | 59.4 | Partially met | 52.0 | Partially met |
| Arizona | 70.5 | Partially met | 47.2 | Partially met | 61.1 | Partially met |
| Arkansas | 59.1 | Partially met | 45.9 | Partially met | 72.6 | Partially met |
| California | 70.5 | Fully met | 53.3 | Partially met | 49.9 | Fully met |
| Colorado | 71.7 | Partially met | 51.7 | Partially met | 55.7 | Partially met |
| Connecticut | 75.2 | Partially met | 55.6 | Partially met | 43.1 | Partially met |
| Delaware | 61.9 | Partially met | 44.6 | Partially met | 55.5 | Partially met |
| District of Columbia | 70.2 | Partially met | 54.1 | Partially met | 41.2 | Partially met |
| Florida | 56.1 | Partially met | 38.5 | Partially met | 50.3 | Partially met |
| Georgia | 67.1 | Partially met | 54.4 | Partially met | 65.7 | Partially met |
| Hawaii | 60.6 | Partially met | 44.7 | Partially met | 50.7 | Partially met |
| Idaho | 64.5 | Did not address | 49.7 | Did not address | 66.7 | Did not address |
| Illinois | 71.9 | Partially met | 47.4 | Partially met | 57.2 | Fully met |
| Indiana | 64.6 | Did not address | 45.1 | Did not address | 60.3 | Partially met |
| Iowa | 69.8 | Partially met | 53.1 | Partially met | 61.4 | Partially met |
| Kansas | 68.4 | Partially met | 48.6 | Partially met | 57.3 | Did not address |
| Kentucky | 61.7 | Partially met | 49.9 | Partially met | 63.9 | Did not address |
| Louisiana | 51.4 | Partially met | 39.7 | Partially met | 68.3 | Partially met |
| Maine | 79.4 | Partially met | 62.1 | Partially met | 39.0 | Partially met |
| Maryland | 67.1 | Partially met | 47.3 | Partially met | 53.6 | Partially met |
| Massachusetts | 68.7 | Did not address | 49.9 | Did not address | 47.8 | Did not address |
| Michigan | 71.9 | Partially met | 53.4 | Partially met | 52.1 | Partially met |
| Minnesota | 75.1 | Partially met | 56.5 | Partially met | 49.3 | Partially met |
| Mississippi | 50.1 | Fully met | 38.2 | Fully met | 74.3 | Partially met |
| Missouri | 70.1 | Partially met | 57.4 | Partially met | 58.3 | Did not address |
| Montana | 78.9 | Partially met | 54.8 | Partially met | 59.4 | Partially met |

| State | Ate Fruit at Least Once Daily, 2022–2023 (%) a | Serve Fruits of Several Varieties: State Center-Based ECE Licensing Standard, 2023 b | Ate Vegetables at Least Once Daily, 2022–2023 (%) a | Serve Vegetables: State Center-Based ECE Licensing Standard, 2023 b | Drank a Sugary Drink One or More Times Weekly, 2022–2023 (%) a | Avoid Sugar: State Center- Based ECE Licensing Standard, 2023 b |
|----------------|--|---|---|---|--|---|
| Nebraska | 67.6 | Partially met | 50.6 | Partially met | 63.1 | Partially met |
| Nevada | 64.3 | Partially met | 48.9 | Partially met | 63.1 | Partially met |
| New Hampshire | 78.9 | Partially met | 57.3 | Partially met | 35.2 | Partially met |
| New Jersey | 68.9 | Fully met | 46.9 | Fully met | 44.6 | Partially met |
| New Mexico | 66.0 | Fully met | 52.5 | Fully met | 69.0 | Partially met |
| New York | 62.6 | Partially met | 45.1 | Partially met | 51.7 | Partially met |
| North Carolina | 66.6 | Partially met | 51.9 | Partially met | 60.4 | Partially met |
| North Dakota | 70.2 | Did not address | 53.3 | Did not address | 59.9 | Did not address |
| Ohio | 66.8 | Partially met | 49.1 | Fully met | 53.3 | Did not address |
| Oklahoma | 63.0 | Partially met | 44.3 | Partially met | 69.9 | Partially met |
| Oregon | 76.6 | Partially met | 54.0 | Partially met | 51.7 | Partially met |
| Pennsylvania | 69.5 | Fully met | 49.3 | Fully met | 55.6 | Did not address |
| Rhode Island | 71.0 | Partially met | 50.4 | Partially met | 42.2 | Partially met |
| South Carolina | 66.1 | Partially met | 47.1 | Partially met | 59.8 | Partially met |
| South Dakota | 69.4 | Did not address | 50.1 | Did not address | 59.0 | Did not address |
| Tennessee | 65.9 | Partially met | 48.0 | Partially met | 61.9 | Fully met |
| Texas | 59.0 | Partially met | 44.1 | Partially met | 69.2 | Fully met |
| Utah | 69.6 | Partially met | 47.6 | Partially met | 65.9 | Partially met |
| Vermont | 75.8 | Partially met | 62.6 | Partially met | 41.8 | Partially met |
| Virginia | 70.1 | Partially met | 54.2 | Partially met | 55.9 | Partially met |
| Washington | 81.1 | Partially met | 59.5 | Partially met | 49.2 | Partially met |
| West Virginia | 63.4 | Partially met | 54.4 | Partially met | 68.6 | Fully met |
| Wisconsin | 73.7 | Partially met | 58.4 | Partially met | 54.2 | Partially met |
| Wyoming | 69.3 | Did not address | 51.6 | Did not address | 66.1 | Did not address |

^a Data include children 1–5 years.

^b States can include high-impact obesity prevention standards in their ECE licensing and administrative regulations. Family child care homes are not included. This indicator reports whether states fully met, partially met, did not address, or contradicted the respective standards: (Serve Fruit: Serve fruits of several varieties, especially whole fruits; Serve Vegetables: Serve vegetables, specifically, dark green, orange, and deep yellow vegetables; and root vegetables such as potatoes and viandas; Avoid Sugar: Avoid sugar, including concentrated sweets such as candy, sodas, sweetened drinks, fruit nectars, and flavored milk).

Affordability of Nutritious Foods

When healthy foods are inaccessible or unaffordable, families might feed their children foods that are higher in calories and lower in important nutrients. It is important for all families to be able to afford healthy foods so they can make healthier choices. The table below provides data on the percentage of children 1–5 years of age living in households that were always able to afford nutritious foods during the past year.



| State | Children 1–5 years living in households that were always able to afford nutritious foods, 2022–2023 (%) | | |
|----------------------|---|--|--|
| National | 69.3 | | |
| Alabama | 64.7 | | |
| Alaska | 68.4 | | |
| Arizona | 65.9 | | |
| Arkansas | 59.2 | | |
| California | 77.1 | | |
| Colorado | 73.8 | | |
| Connecticut | 74.0 | | |
| Delaware | 67.9 | | |
| District of Columbia | 87.2 | | |
| Florida | 61.0 | | |
| Georgia | 69.0 | | |
| Hawaii | 65.2 | | |
| Idaho | 65.9 | | |
| Illinois | 74.2 | | |
| Indiana | 67.0 | | |
| Iowa | 67.9 | | |
| Kansas | 68.7 | | |
| Kentucky | 59.8 | | |
| Louisiana | 60.8 | | |
| Maine | 72.3 | | |
| Maryland | 76.7 | | |
| Massachusetts | 79.3 | | |
| Michigan | 70.6 | | |
| Minnesota | 77.4 | | |
| Mississippi | 58.8 | | |
| Missouri | 63.4 | | |
| Montana | 69.7 | | |
| Nebraska | 64.0 | | |
| Nevada | 63.7 | | |
| | | | |

| State | Children 1–5 years living in households that were always able to afford nutritious foods, 2022–2023 (%) |
|----------------|---|
| New Hampshire | 74.8 |
| New Jersey | 74.5 |
| New Mexico | 68.0 |
| New York | 72.7 |
| North Carolina | 69.2 |
| North Dakota | 72.3 |
| Ohio | 70.5 |
| Oklahoma | 59.4 |
| Oregon | 72.7 |
| Pennsylvania | 70.1 |
| Rhode Island | 76.6 |
| South Carolina | 70.0 |
| South Dakota | 69.5 |
| Tennessee | 67.9 |
| Texas | 59.9 |
| Utah | 73.0 |
| Vermont | 78.4 |
| Virginia | 72.3 |
| Washington | 74.8 |
| West Virginia | 62.0 |
| Wisconsin | 70.3 |
| Wyoming | 71.6 |

INDICATORS, DATA SOURCES, AND METHODOLOGY

| Breastfeeding | | |
|---|--|--|
| Indicators | Data Sources & Methodology | |
| Ever breastfed (%) Breastfed at 12 months (%) Exclusive breastfeeding through 6 months (%) Breastfed infants fed formula before 2 days (%) | Source: CDC National Immunization Survey – Child (NIS-Child) 2023 and 2024, among children born in 2022. https://www.cdc.gov/breastfeeding-data/survey/ The NIS-Child provides current national, state, and selected urban-area estimates of vaccination coverage rates for U.S. children and is used to estimate national and state breastfeeding rates. Because breastfeeding data are obtained by parent or guardian recall when children are between 19 and 35 months of age, breastfeeding rates are analyzed by when children were born rather than by the survey year. This report presents the proportion of children who were ever breastfed or given breast milk, the proportion of children who continued breastfeeding or received breast milk through 12 months of age or older, the proportion that were fed only breast milk through 6 months of age or older and the proportion of breastfed infants who were given infant formula before 2 days of age. | |



Supports for Infant Feeding

| Indicators | Data Sources & Methodology |
|--|--|
| mPINC total score | Source: Maternity Practices in Infant Nutrition and Care (mPINC) Survey, 2024. https://www.cdc.gov/breastfeeding-data/mpinc/index.html CDC's national survey of Maternity Practices in Infant Nutrition and Care (mPINC) assesses and reports on maternity care practices and policies to encourage hospitals to make improvements that better support breastfeeding. mPINC national and state total scores represent the overall level of maternity care practices and policies that support optimal infant feeding during the previous year. Possible scores are from 0 to 100, with higher scores indicating better maternity care practices and policies supportive of optimal infant feeding. Scores are not reported for District of Columbia because of small sample sizes but are included in the US national total mPINC score. Read more about the mPINC scoring algorithm at https://www.cdc.gov/breastfeeding-data/mpinc/scoring.html. |
| Weeks of paid family and medical leave available for the care of a new child | Source: Congressional Research Service Report: Paid Family and Medical Leave in the United States as of March 26, 2025. (Donovan, SA). Paid Family and Medical Leave in the United States (CRS Report No. R44835). https://www.congress.gov/crs_external_products/R/PDF/R44835/R44835.30.pdf Paid family and medical leave (PFML) refers to partially or fully compensated time away from work for specific and generally significant family caregiving needs, such as the arrival of a new child or serious illness of a close family member, or an employee's own serious medical needs. Among states with enacted legislation, the PFML indicator identifies the number of weeks of paid parental leave benefits for the care of a new child by birth, adoption, or foster care available to eligible employees within a benefit year. As of March 2025, Delaware, Maine, Maryland, and Minnesota had enacted legislation to create a paid family and medical leave insurance program, but the program was not yet paying benefits. Paid leave benefits for eligible employees will be available in these states in 2026. Voluntary paid family and medical leave programs in New Hampshire and Vermont are not included. |
| Encourage and support breastfeeding: state center-based ECE licensing standard | Source: University of Colorado College of Nursing. 2023 Supplement: Achieving a State of Healthy Weight. State Profile Pages: Child Care Centers. University of Colorado Anschutz Medical Campus; 2024 https://nursing.cuanschutz.edu/research/healthy-weight States can include high-impact obesity prevention standards in their ECE licensing regulations. This indicator reports whether states' licensing fully met, partially met, did not address, or contradicted standard IA1 for child care centers: Encourage and support breastfeeding and feeding of breast milk by making arrangements for mothers to feed their children on-site. Family child care homes are not included. |



| | ~ = | | | |
|--------|-----|----|-----|-----|
| First: | Sol | hi | F.U | ode |

| Indicators | Data Sources & Methodology |
|--|--|
| Early solid foods (%) | Source: National Survey of Children's Health (NSCH) 2021–2023 Survey Years, among children born in 2019–2021. https://www.childhealthdata.org/learn-about-the-nsch/NSCH NSCH is a national survey that provides data on multiple aspects of children's health and well-being. Caregivers were asked their child's age at which they first fed anything other than breast milk or infant formula. Introducing foods before 4 months of age was considered early solid food introduction. Only children 1–3 years of age were included to improve the reliability of recall. Children were excluded if the reported age that they were first given solid foods was unlikely to be correct (after >12 months of age) or the combination of breastfeeding duration, infant formula introduction, and solid food introduction suggested the child had no source of nutrition for ≥2 months. Data are analyzed by when children were born rather than survey year. |
| Timely introduction of solid foods: state center-based ECE licensing standard | Source: University of Colorado College of Nursing. 2023 Supplement: Achieving a State of Healthy Weight. State Profile Pages: Child Care Centers. University of Colorado Anschutz Medical Campus; 2024 https://nursing.cuanschutz.edu/research/healthy-weight States can include high-impact obesity prevention standards in their ECE licensing regulations. This indicator reports whether states' licensing fully met, partially met, did not address, or contradicted, standard IC2 for child care centers: Introduce age-appropriate solid foods no sooner than 4 months of age, and preferably around 6 months of age. Family child care homes are not included. |



| Healthy Eating | | | | |
|---|--|--|--|--|
| Indicators | Data Sources & Methodology | | | |
| Ate fruit at least once daily (%) | Source: National Survey of Children's Health (NSCH) 2022–2023 Survey Years. https://www.childhealthdata.org/learn-about-the-nsch/NSCH . NSCH is a national survey that provides data on multiple aspects of children's health and well-being. Respondents with children 1–5 years were asked about how many times in the past week did the child eat fruit*? This report presents the percentage of children 1–5 years who ate fruits at least once daily (≥1 time per day in preceding week). *Fruit includes fresh, frozen, or canned. It does not include juice. | | | |
| Serve fruits of several varieties: state center-based ECE licensing standard | Source: University of Colorado College of Nursing. 2023 Supplement: Achieving a State of Healthy Weight. State Profile Pages: Child Care Centers. University of Colorado Anschutz Medical Campus; 2024 https://nursing.cuanschutz.edu/research/healthy-weight States can include high-impact obesity prevention standards in their ECE licensing regulations. This indicator reports whether states' licensing fully met, partially met, did not address, or contradicted standard NB3 for child care centers: Serve fruits of several varieties, especially whole fruits. Family child care homes are not included. | | | |
| Ate vegetables at least once daily (%) | Source: National Survey of Children's Health (NSCH) 2022–2023 Survey Years. https://www.childhealthdata.org/learn-about-the-nsch/NSCH . NSCH is a national survey that provides data on multiple aspects of children's health and well-being. Respondents with children 1–5 years were asked about how many times in the past week did the child eat vegetables*? This report presents the percentage of children 1–5 years who ate vegetables at least once daily (≥1 time per day in preceding week). *Vegetables include fresh, frozen, or canned. It does not include french fries, fried potatoes, or potato chips. | | | |
| Serve vegetables: state center-based ECE licensing standard | Source: University of Colorado College of Nursing. 2023 Supplement: Achieving a State of Healthy Weight. State Profile Pages: Child Care Centers. University of Colorado Anschutz Medical Campus; 2024. https://nursing.cuanschutz.edu/research/healthy-weight States can include high-impact obesity prevention standards in their ECE licensing regulations. This indicator reports whether states' licensing fully met, partially met, did not address, or contradicted standard NB2 for child care centers: Serve vegetables, specifically, dark green, orange, and deep yellow vegetables; and root vegetables such as potatoes and viandas. Family child care homes are not included. | | | |
| Drank a sugary drink one or more times weekly (%) | Source: National Survey of Children's Health (NSCH) 2022–2023 Survey Years. https://www.childhealthdata.org/learn-about-the-nsch/NSCH NSCH is a national survey that provides data on multiple aspects of children's health and well-being. Respondents with children 1–5 years were asked how many times in the past week did the child drink sugary drinks such as soda, fruit drinks, sports drinks, or sweet tea? This report presents the percentage of children 1–5 years who drank a sugary drink one or more times weekly (≥1 time in preceding week). | | | |
| Avoid sugar: state center-based ECE licensing standard | Source: University of Colorado College of Nursing. 2023 Supplement: Achieving a State of Healthy Weight. State Profile Pages: Child Care Centers. University of Colorado Anschutz Medical Campus; 2024 https://nursing.cuanschutz.edu/research/healthy-weight States can include high-impact obesity prevention standards in their ECE licensing regulations. This indicator reports whether states' licensing fully met, partially met, did not address, or contradicted standard NG2 for child care centers: Avoid sugar, including concentrated sweets such as candy, sodas, sweetened drinks, fruit nectars, and flavored milk. Family child care homes are not included. | | | |

Affordability of Nutritious Foods

| Indicators | Data Sources & Methodology |
|--|---|
| Children 1–5 years living in households that could always afford to eat nutritious foods (%) | Source: National Survey of Children's Health (NSCH) 2022–2023 Survey Years. https://www.childhealthdata.org/learn-about-the-nsch/NSCH NSCH is a national survey that provides data on multiple aspects of children's health and well-being. Respondents from households with children 1–5 years were asked to describe their ability to afford and eat nutritious foods during the past 12 months. This report presents the percentage of children 1–5 years living in households that could always afford to eat nutritious foods. |



For more information please contact

Division of Nutrition, Physical Activity, and Obesity National Center for Chronic Disease Prevention and Health Promotion Centers for Disease Control and Prevention

www.cdc.gov/nutrition/php/resources 800-CDC-INFO (800-232-4636)

This report was first published in June 2025 and updated in August 2025. The update replaced data on four breastfeeding indicators among children born in 2021 with data from children born in 2022, National Immunization Survey-Child.

Centers for Disease Control and Prevention. Early Childhood Nutrition Report 2025. Atlanta, Georgia. 2025.

