

Menveo | Overview of Two Presentations

Menveo is a MenACWY vaccine product approved to prevent invasive meningococcal disease caused by *Neisseria meningitidis* serogroups A, C, W, and Y.

Menveo is available in two presentations:

- **Two-vial** (requires reconstitution)
For ages **2 months and older**
- **Single-vial** (ready-to-use)
For ages **10 years and older**
NOT for use in individuals before their 10th birthday

To extend existing stocks of MenACWY vaccine products, administer specific presentations of age-appropriate vaccines:

Ages 2 months through 23 months	Ages 2 years through 9 years	Ages 10 years and older
Menveo Two-vial (requires reconstitution)	MenQuadfi (single-dose vial)	Menveo Single-vial (ready-to-use) or MenQuadfi (single-dose vial)

	Important Differences Between Menveo Presentations	
	Two-vial (vaccine + vaccine)	Single-vial (ready-to-use)
		
Ages Indications	Ages 2 months and older	Ages 10 years and older
Presentation	Supplied as two vials Requires reconstitution • Orange cap vial contains MenA lyophilized conjugate • Gray cap vial contains MenCYW-135 liquid conjugate	Supplied as one ready-to-use vial Reconstitution not required • Pink cap vial
Packaging	Each carton contains 10 vials for 5 doses Five vials of MenA lyophilized conjugate component  Five vials of MenCWY-135 liquid conjugate component 	Each carton contains 10 vials for 10 doses  Ten single vials of ready-to-use Menveo
Storage	Storage and handling is the same for both presentations: Store refrigerated between 36°F and 46°F (2°C and 8°C) Protect from light Do not freeze — frozen/previously frozen product should be discarded	
Preparation	• Reconstitution required • Only reconstitute using vials supplied by the manufacturer • Mix vaccine vials before administering	• Ready-to-use • Reconstitution not required
	See Menveo package insert for preparation guidance	
Dosage/Route	• 0.5 mL/intramuscular (IM) injection • Use immediately after reconstituted	• 0.5 mL/intramuscular (IM) injection

